

A summary list of fossil spiders and their relatives

compiled by

Jason A. Dunlop (Berlin), David Penney (Manchester) & Denise Jekel (Berlin)

with additional contributions from Lyall I. Anderson, Simon J. Braddy,
James C. Lamsdell, Paul A. Selden & O. Erik Tetlie

Suggested citation:

Dunlop, J. A., Penney, D. & Jekel, D. 2013. A summary list of fossil spiders and their relatives. *In* Platnick, N. I. (ed.) The world spider catalog, version 14.0 American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html>

Last updated: 21.06.2013

INTRODUCTION

Fossil spiders have not been fully cataloged since Bonnet's *Bibliographia Araneorum* and are not included in the current Catalog. Since Bonnet's time there has been considerable progress in our understanding of the fossil record of spiders – and other arachnids – and numerous new taxa have been described. For an overview see Dunlop & Penney (2012). Spiders remain the single largest fossil group, but our aim here is to offer a summary list of all fossil Chelicerata in their current systematic position; as a first step towards the eventual goal of combining fossil and Recent data within a single arachnological resource.

To integrate our data as smoothly as possible with standards used for living spiders, our list for Araneae follows the names and sequence of families adopted in the Platnick Catalog. For this reason some of the family groups proposed in Wunderlich's (2004, 2008, 2012) monographs of amber and copal spiders are not reflected here, and we encourage the reader to consult these studies for details and alternative opinions. Extinct families have been inserted in the position which we hope best reflects their probable affinities. For other arachnid groups we have largely followed the nomenclature and family sequences adopted in other online or printed summaries; for example Victor Fet *et al.*'s work on scorpions, Mark Harvey's catalogues of pseudoscorpions and the 'minor' orders – all of which also list the fossils – Adriano Kury's harvestman overviews and the third edition of the Manual of Acarology for mites. For all groups, genus and species names were compiled from established lists and cross-referenced against the primary literature.

We aim to reflect the latest published opinions on the taxonomy of fossil species. A caveat here is that some synonymies and transfers proposed in the literature were only provisional or tentative in nature. At times we were forced to interpret whether a formal nomenclatural change had actually been made, and we have tried to accommodate these difficulties as best as possible. We should also stress that many historical fossil types require revision. Older species names assigned to common, modern genera such as *Araneus*, *Clubiona* or *Linyphia* among the spiders, should be treated with caution. The list has been extended to include Recent species – particularly some spiders and numerous oribatid mites – found as (sub)fossils. These are generally specimens of Quaternary age found in copal, or recovered from peats or archeological sites.

We have provided references for the first descriptions of all the fossil species, and where possible we have added the relevant taxonomic literature for all the taxon names which we mention here. We should, however, note that for some groups (especially mites) recovering the correct author and date for higher taxa proved challenging, and we hope in future releases to be able to clarify these names and augment the reference list accordingly. Formal synonymy lists for the fossil species are being compiled and that which we have for individual taxa can be made available upon request upon a 'fair use' basis. As with any project of this size, we cannot guarantee the accuracy of all these entries and we encourage readers to forward omissions or corrections to <jason.dunlop@mfn-berlin.de> or <David.Penney@manchester.ac.uk>.

PRINCIPAL CHANGES SINCE THE LAST UPDATE

A significant proposal comes from the paper of Lamsdell (2013), who argued that the fossils traditionally assigned to the synziphosurines may in fact include both stem horseshoe crabs and stem chelicerates. The section on Xiphosura has thus been reorganised to try and accommodate this hypothesis, and a category Xiphosura *sensu stricto* is recognised. A number of major clades forming the base of Chelicerata have also been proposed. Other new discoveries from 2013 include two new sea spiders from the Ordovician and Devonian respectively, a new Devonian horseshoe crab from China, a few new Silurian eurypterid genera plus one higher taxon, and a *Tityus* scorpion from Dominican amber. There are also phoretic mesostigmatid mites from a number of families recorded on beetles and new labidostommatids among the acariform mites. For spiders, there is the discovery of a Triassic mygalomorph spider – one of the few examples of a Triassic spider and the oldest putative record of Atypoidea – as well as new records of jumping spiders from Dominican amber.

ACKNOWLEDGMENTS

We are, as ever, especially grateful to Norman Platnick for agreeing to host this list as an appendix to the Catalog, to Paul Selden for encouragement and support and to those colleagues who have advised us on oversights and/or provided further literature.

EXPLANATIONS

- † indicates an entirely extinct genus, family or other higher taxon
- all species listed assumed to be extinct unless marked **[Recent]**
- * indicates the type species of (fossil) genera

Stratigraphical abbreviations:

pЄ = Precambrian, Є = Cambrian, O = Ordovician, S = Silurian,

D = Devonian, C = Carboniferous, P = Permian

Tr = Triassic, J = Jurassic, K = Cretaceous

Pa = Palaeogene, Ne = Neogene, Qt = Quaternary

PYCNOGONIDA

11 currently valid species of fossil sea spider

- note that in some modern phylogenies the Palaeozoic genera resolve *within* the crown group

PYCNOGONIDA Latreille, 1810 Cambrian – Recent

= ARACHNOPODA Dana, 1853

- † **Cambropycnogon Waloszek & Dunlop, 2002** **Cambrian**
1. *Cambropycnogon klausmuelleri* Waloszek & Dunlop, 2002* € 'Orsten', Sweden
Pycnogonid affinities questioned by Bamber (2007)
- † **Haliestes Siveter, Sutton, Briggs & Siveter, 2004** **Silurian**
2. *Haliestes dasos* Siveter, Sutton, Briggs & Siveter, 2004* S Herefordshire Lgst.
- † **Flagellopantopus Poschmann & Dunlop, 2006** **Devonian**
3. *Flagellopantopus blocki* Poschmann & Dunlop, 2006* D Hünsruckschiefer
- † **Palaeomarachne Rudkin, Cuggy, Young & Thompson, 2013** **Ordovician**
4. *Palaeomarachne granulata* Rudkin, Cuggy, Young & Thompson, 2013* O Manitoba, Canada
- † **Pentapantopus Kühl, Poschmann & Rust, 2013** **Devonian**
5. *Pentapantopus vogteli* Kühl, Poschmann & Rust, 2013* D Hünsruckschiefer
- † **PALAEOISOPODIDAE Dubinin, 1957** **Devonian**
- † **Palaeoisopus Broili, 1928** **Devonian**
6. *Palaeoisopus problematicus* Broili, 1928* D Hünsruckschiefer
- † **PALAEOPANTOPODIDAE Broili, 1930** **Devonian**
- † **Palaeopantopus Broili, 1928** **Devonian**
7. *Palaeopantopus maucheri* Broili, 1928* D Hünsruckschiefer
- PANTOPODA Gerstaecker, 1863** **Devonian – Recent**
- = PEGMATA Fry, 1978
- family uncertain
- † **Palaeothea Bergström, Stürmer & Winter, 1980** **Devonian**
8. *Palaeothea devonica* Bergström, Stürmer & Winter, 1980* D Hünsruckschiefer
- AUSTRODECIDAE Stock, 1954** **Recent**
- no fossil record
- PYCNOGONIDAE Wilson, 1878** **Recent**
- no fossil record

COLOSSENDEIDAE Hoek, 1881 **?Jurassic – Recent**

= PASITHOIDAE Sars, 1891

= RHOPALORHYNCHIDAE Fry, 1978

† **Colossopantopodus Charbonnier, Vannier & Riou, 2007** **Jurassic**

9. *Colossopantopodus boissinensis* Charbonnier, Vannier & Riou, 2007* . J La Voulte-sur-Rhône
tentative referal

AMMOTHEIDAE Dohrn, 1881 **?Jurassic – Recent**

= EURYCIDIDAE Sars, 1891

= OORHYNCHIDAE Schimkewitsch, 1913

= TANYSTYLIDAE Schimkewitsch, 1913

= AMMOTHELLIDAE Fry, 1978

= EPHYROGYMNIDAE Fry, 1978

= PARANYMPHONIDAE Fry, 1978

= SERICOSURIDAE Fry, 1978

= TRYGAEIDAE Fry, 1978

† **Palaeopycnogonides Charbonnier, Vannier & Riou, 2007** **Jurassic**

10. *Palaeopycnogonides gracilis* Charbonnier, Vannier & Riou, 2007* J La Voulte-sur-Rhône
tentative referal

CALLIPALLENIDAE Hilton, 1942 **Recent**= PALLENIDAE Wilson, 1878 [*Pallene* is a preoccupied genus]

= CHEILAPALLENIDAE Fry, 1978

= CLAVIGEROPALLENIDAE Fry, 1978

= HANNONIDAE Fry, 1978

= METAPALLENIDAE Fry, 1978

= QUEUBIDAE Fry, 1978

= STYLOPALLENIDAE Fry, 1978

no fossil record

NYMPHONIDAE Wilson, 1878 **Recent**

no fossil record

PALLENOPSISAE Fry, 1978 **Recent**

no fossil record

ENDEIDAE Norman, 1904 **?Jurassic – Recent**† **Palaeoendeis Charbonnier, Vannier & Riou, 2007** **Jurassic**

11. *Palaeoendeis elmii* Charbonnier, Vannier & Riou, 2007* J La Voulte-sur-Rhône
tentative referal

PHOXICHILIDIIDAE Sars, 1891 **Recent**

= ANOPLODACTYLIDAE Fry, 1978

= PHOXIPHILYRIDAE Fry, 1978

no fossil record

RHYNCHOTHORACIDAE Thompson, 1909 **Recent**

no fossil record

MISIDENTIFICATIONS

1. *Palpipes cursor* Roth, 1854 [crustacean] J Solnhofen
2. *Pentapalaeopycnon inconspicua* Hedgpeth, 1978 [crustacean] J Solnhofen
3. *Phalangites multipes* Münster, 1851 [crustacean] J Solnhofen
4. *Phalangites priscus* Münster, 1839 [crustacean] J Solnhofen
5. *Pycnogonites uncinatus* Quenstedt, 1852 [crustacean] J Solnhofen

c. 1,300 Recent species

EUCHELICERATA

4 currently valid, but unplaced euchelicerate fossil species

- *Offacolus* has been described in detail from reconstructions based on serial sections, and was resolved in some phylogenies to a basal position within Euchelicerata
- *Dibasterium* was described as a horseshoe crab, albeit one with multiple biramous appendages
- the other listed taxa are mostly poor or incomplete specimens which have been treated as either xiphosurans, chasmataspidids or eurypterids
- resting impressions imply that Chasmataspidida were probably present in the late Cambrian

EUCHELICERATA Weygoldt & Paulus, 1979 ?Cambrian – Recent

STEM-EUCHELICERATA?

- † *Offacolus* Orr, Siveter, Briggs, Siveter & Sutton, 2000 **Silurian**
1. *Offacolus kingi* Orr, Siveter, Briggs, Siveter & Sutton, 2000* S Herefordshire Lgst.
- † *Dibasterium* Briggs, Siveter, Siveter, Sutton, Garwood & Legg, 2012 **Silurian**
2. *Dibasterium durgae* Briggs, Siveter, Siveter, Sutton, Garwood & Legg, 2012* S Herefordshire Lgst.

EUCHELICERATA INCERTAE SEDIS

- † *Polystomurum* Novojilov, 1958 **Devonian**
3. *Polystomurum stormeri* Novojilov, 1958* D Voroneje, Siberia
- † *Thurandina* Størmer, 1974 **Devonian**
4. *Thurandina waterstoni* Størmer, 1974* D Alken an der Mosel

XIPHOSURA *s. lat.*

103 currently valid species traditionally assigned to horseshoe crabs, of which 82 are unequivocal Xiphosura

- Lamsdell (2013) argued that Xiphosura may not be monophyletic and that a number of fossils traditionally placed as stem-group (synziphosurine) horseshoe crabs are actually stem-group euchelicerates. The list below attempts to reflect this position, whereby it should be noted that in this scheme the Planaterga clade would also include Chasmataspidida, Eurpeterida and Arachnida and Planaterga is nested within Prosomapoda.

PROSOMAPODA Lamsdell, 2013a	Siliurian – Recent
FAMILY UNSPECIFIED	
† <i>Anderella</i> Moore, McKenzie & Lieberman, 2007	Carboniferous
1. <i>Anderella parva</i> Moore, McKenzie & Lieberman, 2007*	C Bear Gulch
† <i>Borchgrevinkium</i> Novojilov, 1959	Devonian
2. <i>Borchgrevinkium taimyrensis</i> Novojilov, 1959*	D Taimyr, Siberia
† <i>Camanchia</i> Moore, Briggs, Braddy & Shultz, 2011	Silurian
3. <i>Camanchia grovensis</i> Moore, Briggs, Braddy & Shultz, 2011*	S Scotch Grove, Iowa
† <i>Legrandella</i> Eldredge, 1974	Devonian
4. <i>Legrandella lombardii</i> Eldredge, 1974*	D Cochabamba, Bolivia
† <i>Venustulus</i> Moore, 2005 in Moore et al.	Silurian
5. <i>Venustulus waukeshaensis</i> Moore, 2005 in Moore et al.*	S Waukesha Lgst.
† WEINBERGINIDAE Richter & Richter, 1929	Devonian
† <i>Weinbergina</i> Richter & Richter, 1929	Devonian
6. <i>Weinbergina opitzi</i> Richter & Richter, 1929*	D Hünsruckschiefer
PLANATERGA Lamsdell, 2013a	Siliurian – Recent
FAMILY UNSPECIFIED	
† <i>Bembicosoma</i> Laurie, 1899	Silurian
7. <i>Bembicosoma pomphicus</i> Laurie, 1899*	S Pentland hills
† <i>Cyamocephalus</i> Currie, 1927	Silurian
8. <i>Cyamocephalus loganensis</i> Currie, 1927*	S Lesmahagow
† <i>Pseudoniscus</i> Nieszkowski, 1859	Silurian
= † <i>Neolimulus</i> Woodward, 1868a	
9. <i>Pseudoniscus aculeatus</i> Nieszkowski, 1859*	S Saaremaa
10. <i>Pseudoniscus clarkei</i> Ruedemann, 1916	S Pittsford, New York
11. <i>Pseudoniscus falcatus</i> (Woodward, 1868a)	S Lesmahagow
12. <i>Pseudoniscus roosevelti</i> Clarke, 1902	S 'Bertie Waterlime'
† <i>Bunaia</i> Clarke, 1919	Silurian

13. '*Bunaia*' *heintzi* Størmer, 1934a S Spitsbergen
14. *Bunaia woodwardi* Clarke, 1919* S 'Bertie Waterlime'
- † **BUNODIDAE Packard, 1896** **Silurian**
- † ***Bunodes* Eichwald, 1854** **Silurian**
 = † *Exapinurus* Nieszkowski, 1859
15. *Bunodes lunula* Eichwald, 1854* S Saaremaa
 i. = *Bunodes rugosus* Eichwald, 1854 S Saaremaa
 ii. = *Exapinurus schrenki* Nieszkowski, 1859 S Saaremaa
- † ***Limuloides* Woodward, 1865** **Silurian**
 = † *Hemiaspis* Woodward, 1864 [preoccupied]
16. *Limuloides limuloides* (Woodward, 1865) S Ludlow
17. *Limuloides horridus* (Woodward, 1872a) S Ludlow
18. *Limuloides salweyi* (Woodward, 1872a) S Ludlow
 i. = *Hemiaspis tuberculatus* (Salter in Woodward, 1872a) S Ludlow
19. *Limuloides speratus* Woodward, 1872a S Ludlow
 i. = *Hemiaspis optatus* (Salter in Woodward, 1872a) S Ludlow
- † ***Pasternakevia* Selden & Drygant, 1987** **Silurian**
20. *Pasternakevia podolica* Selden & Drygant, 1987* S Podolia

Planaterga *sensu* Lamsdell (2013a) also includes chasmataspids, eurypterids and arachnids

XIPHOSURA Latreille, 1802 **Ordovician – Recent**
 = MEROSTOMATA Dana, 1852

FAMILY UNSPECIFIED

- † ***Kiaeria* Størmer, 1934b** **Silurian**
21. *Kiaeria limuloides* Størmer, 1934b* S Ringerike
- † ***Maldybulakia* Tesakov & Alekseev, 1998** **Devonian**
 = † *Lophodesmus* Tesakov & Alekseev, 1992 [preoccupied]
- NB: Originally described as possible myriapods
22. *Maldybulakia angusi* Edgecombe, 1998 D New South Wales
23. *Maldybulakia malcomi* Edgecombe, 1998 D New South Wales
24. *Maldybulakia mirabilis* (Tesakov & Alekseev, 1992)* D Kazakhstan
- † ***Willwerathia* Størmer, 1969** **Devonian**
25. *Willwerathia laticeps* (Størmer, 1936a)* D Willwerath
- † **KASIBELINURIDAE Pickett, 1993** **Devonian**
- † ***Kasibelinurus* Pickett, 1993** **Devonian**
26. *Kasibelinurus amicorum* Pickett, 1993* D New South Wales
27. *Kasibelinurus yueya* Lamsdell, Xue & Selden, 2013 D Yunann, China
- possible kasibelinurids?

28. '*Belinurus' alleghenyensis* Eller, 1938a D New York State
29. '*Belinurus' carterae* Eller, 1940 D Pennsylvania
30. '*Prestwichia' randalli* Beecher, 1902 D Pennsylvania
- † **ELLERIDAE Raymond, 1944** **Devonian**
- † ***Elleria* Raymond, 1944** **Devonian**
31. *Elleria morani* (Eller, 1938b)* D Pennsylvania
- XIPHOSURIDA Latreille, 1802** **Ordovician – Recent**
- family uncertain
- † ***Lunataspis* Rudkin, Young & Nowlan, 2008** **Ordovician**
32. *Lunataspis aurora* Rudkin, Young & Nowlan, 2008 O Manitoba
- † **BELLINURINA Zittel & Eastman, 1913** **Carboniferous**
- † **BELLINURIDAE Zittel & Eastman, 1913** **Carboniferous**
- † ***Bellinurus* Pictet, 1846** **Carboniferous**
- = † *Belinurus* König, 1851
- = † *Steropsis* Baily, 1869
- = † *Koenigiella* Raymond, 1944
- NB: Pictet's 1846 name *Bellinurus* [sic] was based on a misspelling of *Belinurus* from König's unpublished plates, which themselves only became available posthumously as of 1851
33. *Bellinurus arcuatus* Baily, 1863 C Coal Measues
34. *Bellinurus baldwini* Woodward, 1907b C Coal Measues
35. *Bellinurus bellulus* Pictet, 1846 C Coalbrookdale, UK
36. *Bellinurus carwayensis* Dix & Pringle, 1929 C South Wales, UK
37. *Bellinurus concinnus* Dix & Pringle, 1929 C South Wales, UK
38. *Bellinurus grandaevus* Jones & Woodward, 1899 C Nova Scotia
39. *Bellinurus iswariensis* (Chernyshev, 1928) C Donetz Basin
40. *Bellinurus kiltorkensis* Baily, 1869 C Coal Measues
41. *Bellinurus koenigianus* Woodward, 1872a C Coal Measues
42. *Bellinurus lacoey* Packard, 1885 C Mazon Creek
43. *Bellinurus longicaudatus* Woodward, 1907b C Coal Measues
44. *Bellinurus lunatus* (Martin, 1809) C Mansfield, UK
45. *Bellinurus metschetensis* (Chernyshev, 1928) C Donetz Basin
46. *Bellinurus morgani* Dix & Pringle, 1930 C South Wales, UK
47. *Bellinurus pustulosus* Dix & Pringle, 1929 C South Wales, UK
48. *Bellinurus reginae* Baily, 1863 C Coal Measues
49. *Bellinurus stepanovi* (Chernyshev, 1928) C Donetz Basin
50. *Bellinurus trechmanni* Woodward, 1918 C Coal Measues
51. *Bellinurus trilobitoides* (Buckland, 1837)* C Coalbrookdale, UK
52. *Bellinurus truemani* Dix & Pringle, 1929 C South Wales, UK

† EUPROOPIIDAE Eller, 1938b

= † LIOMESASPIDIDAE Raymond, 1944

- † *Anacontium* Raymond, 1944 **Permian**
53. *Anacontium brevis* Raymond, 1944 P Oklahoma
54. *Anacontium carpenteri* Raymond, 1944 P Oklahoma
- † *Euproops* Meek, 1867 **Carbon. – ?Permian**
- = † *Prestwichia* Woodward, 1867 [preoccupied]
- = † *Prestwichianella* Cockerell, 1905 [replacement name for *Prestwichia*]
55. *Euproops anthrax* (Prestwich, 1840) C Coal Measures
56. *Euproops bifidus* Siegfried, 1972 C Coal Measures
57. *Euproops cambrensis* Dix & Pringle, 1929 C Coal Measures
58. *Euproops danae* (Meek & Worthen, 1865)* C Coal Measures
- i. = *Euproops amiae* Woodward, 1918 C Coal Measures
- ii. = *Euproops darrahi* Raymond, 1944 C Coal Measures
- iii. = *Euproops graigolae* Dix & Pringle, 1929 C South Wales
- iv. = *Euproops gventi* Dix & Pringle, 1929 C South Wales
- v. = *Euproops islwyni* Dix & Pringle, 1929 C South Wales
- vi. = *Euproops kilmersdonensis* Ambrose & Romano, 1972 C Kilmersdon, UK
- vii. = *Euproops laevicula* Raymond, 1944 C Coal Measures
- viii. = *Euproops laticephalus* Raymond, 1944 C Coal Measures
- ix. = *Euproops packardi* Willard & Jones, 1935 C Coal Measures
- x. = *Prestwichia (Euproops) scheeleana* Ebert, 1892 C Coal Measures
- xi. = *Euproops thompsoni* Raymond, 1944 C Coal Measures
59. *Euproops longispina* Packard, 1885 C Mazon Creek
60. *Euproops mariae* Crônier & Courville, 2005 C Massif Central
61. *Euproops meeki* Dix & Pringle, 1929 C South Wales
62. *Euproops nitida* Dix & Pringle, 1929 C South Wales
63. *Euproops orientalis* Kobayashi, 1933 ?P Korea
64. *Euproops rotundatus* Prestwich, 1840 C Coal Measures
- Euproops* sp. in Brauckmann (1982) C Piesberg, Germany
- † *Liomesaspis* Raymond, 1944 **Carbon. – Permian**
- = † *Pringlia* Raymond, 1944
- = † *Palatinaspis* Malz & Poschmann, 1993
65. ?*Liomesaspis birtwelli* (Woodward, 1872a) C Coal Measures
66. *Liomesaspis laevis* Raymond, 1944* C Coal Measures
- i. = *Palatinaspis beimbaueri* Malz & Poschmann, 1993 C Saar-Nahe Basin
- ii. = *Pringlia bispinosa* Raymond, 1944 C Coal Measures
- iii. = *Pringlia demaisterei* Vandenbergh, 1961 C Coal Measures
- iv. = *Pringlia fritschi* Remy & Remy, 1959 C Coal Measures
67. *Liomesaspis leonardensis* (Tasch, 1961) P Annelly, Kansas
- † *Prolimulus* Frič, 1899 **Carboniferous**
68. *Prolimulus woodwardi* Frič, 1899* C Nýřany

UNNAMED TAXON

- † **Bellinuroopsis Chernyshev, 1933** **Carboniferous**
 = † *Neobelinuroopsis* Eller, 1938a
 69. *Bellinuroopsis rossicus* Chernyshev, 1933* C Coal Measures
- † **ROLFEIIDAE Selden & Siveter, 1987** **Carboniferous**
- † **Rolfeia Waterston, 1985** **Carboniferous**
 70. *Rolfeia fouldenensis* Waterston, 1985* C Fouldon, Scotland
- LIMULINA Richter & Richter, 1929** **Carbon. – Recent**
 Unanmed specimen *in* Krause *et al.* (2009) Tr Ohrdruf, Germany
- † **PALEOLIMULOIDEA Raymond, 1944** **Carbon. – Jurassic**
- † **PALEOLIMULIDAE Raymond, 1944** **Carbon. – Jurassic**
 = † MESOLIMULIDAE (Størmer, 1952) [in part; see Reik & Gill 1971]
 = † DUBBOLIMULIDAE Pickett, 1984
- † **Limulitella Størmer, 1952** **Triassic – Jurassic**
 = † *Limulites* Schimper, 1853 [preoccupied]
 Limulitella sp. *in* Hauschke *et al.* (2004) Tr Madagascar
 ? *Limulitella* sp. *in* Hauschke & Wilde (2008) Tr Dallau, Germany
 ? *Limulitella* sp. *in* Hauschke *et al.* (2009) Tr Winterswijk
 71. *Limulitella bronniei* (Schimper, 1853)* Tr Grés à Voltzia
 i. = *Limulus sandbergeri* Kirchner, 1923 Tr Germany
 72. *Limulitella henkeli* Fritsch, 1906 Tr Halle, Germany
 73. ? *Limulitella liasokeuperensis* (Braun, 1860) J Germany
 74. *Limulitella vicensis* (Bleicher, 1897) Tr Lorraine
 75. *Limulitella volgensis* Ponomarenko, 1985 Tr Moscow
- † **Paleolimulus Dunbar, 1923** **Carbon. – Triassic**
 = † *Dubbolimulus* Pickett, 1984
 ? *Palaeolimulus* sp. *in* Hauschke & Wilde (2000) Tr Harz, Germany
 76. *Paleolimulus fuchsbergensis* Hauschke & Wilde, 1987 Tr northwest Germany
 77. *Paleolimulus jakovlevi* Glushenko *in* Glushenko & Ivanov, 1961 P Novoselovka, Ukraine
 78. ? *Paleolimulus juresanensis* Chernyshev, 1933 C Ural region
 79. *Paleolimulus longispinus* Schram, 1979 C Bear Gulch, Montana
 80. *Paleolimulus peetae* (Pickett, 1984) Tr New South Wales
 81. *Paleolimulus signatus* (Beecher, 1904) C–P Kansas, Illinois
 i. = *Paleolimulus avitus* Dunbar, 1923* P Kansas
- MORAVURIDAE Příbyl, 1967** **Carboniferous**
- † **Moravurus Příbyl, 1967** **Carboniferous**
 82. *Moravurus rehoi* Příbyl, 1967 C Ostrava-Karviná

- † *Xaniopyramis* Siveter & Selden, 1987 **Carboniferous**
 83. *Xaniopyramis linseyi* Siveter & Selden, 1987* C Weardale, UK
- LIMULOIDEA Zittel, 1885** **Carbon. – Recent**
 unnamed specimen *in* Hauschke & Wilde (1989) P Korbacher Bucht
- † *Alanops* Racheboeuf *et al.*, 2002 **Carboniferous**
 84. *Alanops magnifica* Racheboeuf *et al.*, 2002 C Montceau-les-Mines
- † *Casterolimulus* Holland, Erickson & O'Brien, 1975 **Cretaceous**
 85. *Casterolimulus kletti* Holland, Erickson & O'Brien, 1975* K North Dakota
- † *Panduralimulus* Allen & Feldman, 2005 **Permian**
 86. *Panduralimulus babcocki* Allen & Feldman, 2005 P Texas
- † *Valloisella* Racheboeuf, 1992 **Carboniferous**
 87. *Valloisella lievinensis* Racheboeuf, 1992* C northern France
- † **AUSTROLIMULIDAE Riek, 1955** **Triassic**
- † *Austrolimulus* Riek, 1955 **Triassic**
 88. *Austrolimulus fletcheri* Riek, 1955* Tr New South Wales
- LIMULIDAE Zittel, 1885** **Triassic – Recent**
 = † MESOLIMULIDAE (Størmer, 1952) [in part; see Reik & Gill 1971]
 ?Limulidae gen. et sp. indet *in* Hauschke *et al.* (1992) Tr Rüdersdorf, Germany
- Crenatolimulus** Feldmann, Schweitzer, Dattilo & Farlow, 2011 **Cretaceous**
 89. *Crenatolimulus paluxyensis* Feldmann, Schweitzer, Dattilo & Farlow,
 2011* K Texas
- Limulus** Müller, 1785 **Triassic – Recent**
 90. *Limulus coffini* Reeside & Harris, 1952 K Colorado
 91. "*Limulus*" *decheni* Zinken, 1862 Pa Teuchern, Germany
 [NB: Hauschke & Wilde (2004) considered this intermediate between *Limulus* and *Tachypleus*]
 92. *Limulus priscus* Münster, 1839 Tr Rottweil, Germany
 93. *Limulus woodwardi* Watson, 1909 J Northamptonshire
- † **Mesolimulus** Størmer, 1952 **Triassic – Cretaceous**
Mesolimulus sp. *in* Ross & Vannier (2002) J southern England
94. *Mesolimulus crespelli* Via Boada, 1987 Tr Tarragona, Spain
 95. *Mesolimulus sibiricus* Ponomarenko, 1985 J Siberia
 96. ?*Mesolimulus syriacus* (Woodward, 1879) K Lebanon
 97. *Mesolimulus walchi* (Desmarest, 1822)* J Solnhofen, etc.
 i. = *Limulus brevicauda* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 ii. = *Limulus brevispina* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 iii. = *Limulus intermedius* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 iv. = *Limulus ornatus* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 v. = *Limulus sulcatus* Münster *in* v. d. Hoeven, 1838 J Solnhofen
 vi. = *Limulus giganteus* Münster, 1840 J Solnhofen

NB: not entirely clearly that all these names have been formally synonymised

- † **Psammolimulus Lange, 1923** **Triassic**
 98. *Psammolimulus gottingensis* Lange, 1923* Tr Göttingen, Germany
- Tachypleus Leach, 1819** **Triassic – Recent**
 = † *Heterolimulus* Via Boada & Villalta, 1966
 99. *Tachypleus gadeai* (Via Boada & Villalta, 1966) Tr Tarragona, Spain
- † **Tarracolimulus Romero & Via Boada, 1977** **Triassic**
 100. *Tarracolimulus rieki* Romero & Via Boada, 1977* Tr Tarragona, Spain
- † **Victalimulus Riek & Gill, 1971** **Cretaceous**
 101. *Victalimulus mcqueeni* Riek & Gill, 1971* K Koonwarra
- † **Yunnanolimulus Zhang, Hu, Zhou, Iv & Bai, 2009** **Triassic**
 102. *Yunnanolimulus luopingensis* Zhang, Hu, Zhou, Iv & Bai, 2009* Tr Luoping, China

INCERTAE SEDIS

- † **Belinuropsis Matthew 1910**
 103. *Belinuropsis wigudensis* Matthew, 1910 C Coal Measures

NOMEN DUBIUM

1. *Limulus nathorsti* Jackson, 1906 J southern Sweden

NOMINA NUDA

1. *Euproops rotunda major* (Woodward, 1907) C Sparth Bottoms
 2. *Veltheimia bicorns* Beyschlag & von Fritsch, 1899 C? Rotliegend

MISIDENTIFICATIONS

1. *Belinurus carterae* Eller, 1940 [synonym of *P. eriensis*; see below]
 2. *Bifarius compta* Tasch, 1961 [insect] P Kansas
 3. *Eolimulus alatus* Moberg, 1892 [doubtful xiphosuran] C Öland, Sweden
 4. *Elmocephalus carltonensis* (Tasch, 1963) [?crustacean] P Kansas
 5. *Hemiaspis tunnecliffei* Chapman, 1932 [trilobite] S Victoria
 6. *Hypatocephala rugosa* Tasch, 1961 [insect] P Kansas
 7. *Lemoneites ambiguus* Flower, 1969 [Echinodermata] O Texas
 8. *Lemoneites gomphocaudatus* Flower, 1969 [Echinodermata] O Texas
 9. *Lemoneites mirabilis* Flower, 1969 [Echinodermata] O Texas
 10. *Lemoneites simplex* Flower, 1969 [Echinodermata] O Texas
 11. *Pincombella belmontensis* Chapman, 1932 [insect – Hemiptera] P New South Wales
 12. *Permolimulinella raris* Tasch, 1963 [insect] P Kansas
 13. *Strongylocephalus charactis* Tasch, 1961 [insect] P Kansas
 14. *Protolimulus eriensis* [Xiphosuran trace fossil: see *Selenichnites*]

CHASMATASPIDIDA

8 currently valid species of fossil chasmataspidid

- there are some doubts about the monophy of Chasmataspidida

- † **CHASMATASPIDIDA Caster & Brooks, 1956** ?Camb. – Devonian
 = † DIPLOASPIDIDA Simonetta & Delle Cave, 1978
- † **CHASMATASPIDIDAE Caster & Brooks, 1956** ?Camb. – Ordovician
- † ***Chasmataspis* Caster & Brooks, 1956** ?Camb. – Ordovician
- ? *Chasmataspis* sp. resting traces in Dunlop *et al.* (2004) C Texas
1. *Chasmataspis laurencii* Caster & Brooks, 1956* O Tennessee
- † **DIPLOASPIDIDAE Størmer, 1972** Silurian – Devonian
 = † HETEROASPIDIDAE Størmer, 1972
- † ***Achanarraspis* Anderson, Dunlop & Trewin, 2000** Devonian
2. *Achanarraspis reedi* Anderson, Dunlop & Trewin, 2000* D Achanarras, Scotland
- † ***Diploaspis* Størmer, 1972** Devonian
 = † *Heteroaspis* Størmer, 1972
3. *Diploaspis casteri* Størmer, 1972* D Alken an der Mosel
 i. = *Heteroaspis novojilovi* Størmer, 1972 D Alken an der Mosel
4. *Diploaspis muelleri* Poschmann, Anderson & Dunlop, 2005 D Hombach, Germany
- † ***Forfarella* Dunlop, Anderson & Braddy, 1999** Devonian
5. *Forfarella mitchelli* Dunlop, Anderson & Braddy, 1999* D Arbroath, Scotland
- † ***Loganamaraspis* Tetlie & Braddy, 2004a** Silurian
6. *Loganamaraspis dunlopi* Tetlie & Braddy, 2004a* S Lesmahagow
- † ***Octoberaspis* Dunlop, 2002** Devonian
7. *Octoberaspis ushakovi* Dunlop, 2002* D October Rev. Is.
- DIPLOASPIDIDAE INCERTAE SEDIS
- † **'*Eurypterus*'**
8. '*Eurypterus*' *stoermeri* Novojilov, 1959 D Taimyr, Siberia

no Recent species

EURYPTERIDA

249 currently valid species of fossil sea scorpion

- Tollerton (1989) suggested removing Hibbertopteroidea from Euryperida s.s., but this has not been adopted by subsequent workers and they are treated here as derived stylonurid eurypterids

† EURYPTERIDA Burmeister, 1843	Ordovician – Permian
= † GIGANTOSTRACA Haeckel, 1866	
= † CYRTOCTENIDA Størmer & Waterston, 1968	
† STYLONURINA Diener, 1924	Ordovician – Permian
= † WOODWARDOPTERINA Kjellesvig-Waering, 1959	
= † HIBBERTOPTERINA Størmer, 1974	
† RHENOPTEROIDEA Størmer, 1951	Ordovician – Devonian
= † BRACHYOPTERELLOIDEA Tollerton, 1989	
† RHENOPTERIDAE Størmer, 1951	Ordovician – Devonian
= † BRACHYOPTERELLIDAE Tollerton, 1989	
= † ALKENOPTERIDAE Poschmann & Tetlie, 2004	
† Alkenopterus Størmer, 1974	Devonian
1. <i>Alkenopterus brevitelson</i> Størmer, 1974*	D Alken an der Mosel
2. <i>Alkenopterus burglahrensis</i> Poschmann & Tetlie, 2004	D Westerwald, Germ.
† Brachyopterella Kjellesvig-Waering, 1966a	Silurian
3. <i>Brachyopterella pentagonalis</i> (Størmer, 1934b)*	S Ringerike, Norway
4. <i>Brachyopterella ritchiei</i> Waterston, 1979	S Slot Burn, Scotland
† Brachyopterus Størmer, 1951	Ordovician
5. <i>Brachyopterus stubblefieldi</i> Størmer, 1951*	O Montgomeryshire
† Kiaeropterus Waterston, 1979	Silurian
6. <i>Kiaeropterus cyclophthalmus</i> (Laurie, 1892)	S Pentland Hills, Scotl.
7. <i>Kiaeropterus ruedemanni</i> (Størmer, 1934b)*	S Ringerike, Norway
† Leiopterella Lamsdell, Braddy, Loeffler & Dineley, 2010	Devonian
8. <i>Leiopterella tetliei</i> Lamsdell, Braddy, Loeffler & Dineley, 2010	D Nunavut, Canada
† Rhenopterus Størmer, 1936a	Devonian
9. <i>Rhenopterus diensti</i> Størmer, 1936a*	D Willwerath, Germ.
i. = <i>Rhenopterus latus</i> Størmer, 1936a	D Willwerath, Germ.
10. <i>Rhenopterus macrotuberculatus</i> Størmer, 1974	D Alken an der Mosel
11. <i>Rhenopterus tuberculatus</i> Størmer, 1936a	D Overath, Germ.
† STYLONUROIDEA Kjellesvig-Waering, 1959	Silurian – Devonian
† PARASTYLONURIDAE Waterston, 1979	Silurian – Devonian

- † **Parastylonurus Kjellesvig-Waering, 1966a** **Silurian**
12. *Parastylonurus hendersoni* Waterston, 1979 S Pentland Hills, Scotl.
13. *Parastylonurus ornatus* (Laurie, 1892)* S Scotland
14. ?*Parastylonurus sigmoidalis* Kjellesvig-Waering, 1971 S Shropshire, UK
- † **Stylonurella Kjellesvig-Waering, 1966a** **Silurian – Devonian**
15. *Stylonurella ?arnoldi* (Ehlers, 1935) D Pennsylvania, USA
16. *Stylonurella ?beecheri* (Hall, 1884c) D Pennsylvania, USA
17. *Stylonurella spinipes* (Page, 1859)* S Kip Burn, Scotland
- i. = *Stylonurus logani* Woodward, 1872 S Kip Burn, Scotland
- † **STYLONURIDAE Diener, 1924** **Silurian–Devonian**
- = † LAURIEIPTERIDAE Kjellesvig-Waering, 1966a
- = † PAGEIDAE Kjellesvig-Waering, 1966a
- † **Ctenopterus Clarke & Ruedemann, 1912** **Silurian**
18. *Ctenopterus cestrotus* (Clarke, 1907)* S Otisville, New York
- † **Laurieipterus Kjellesvig-Waering, 1966a** **Silurian**
19. *Laurieipterus elegans* (Laurie, 1899)* S Pentland Hills, Scotl.
- † **Pagea Waterston, 1962** **Devonian**
20. *Pagea plotnicki* Lamsdell, Braddy, Loeffler & Dineley, 2010 D Nunavut, Canada
21. *Pagea sturrocki* Waterston, 1962* D Old Red Sandstone
22. *Pagea symondsii* (Salter, 1859) D Old Red Sandstone
- † **Stylonurus Page, 1856** **Devonian**
23. *Stylonurus powriensis* Page, 1856* D Mid. Valley Scotland
- i. = *Stylonurus ensiformis* Woodward, 1864 D Mid. Valley Scotland
24. ?*Stylonurus shaffneri* Willard, 1933 D Pennsylvania
- † **KOKOMOPTEROIDEA Kjellesvig-Waering, 1966a** **Silurian**
- † **KOKOMOPTERIDAE Kjellesvig-Waering, 1966a** **Silurian**
- † **Kokomopterus Kjellesvig-Waering, 1966a** **Silurian**
25. *Kokomopterus longicaudatus* (Clarke & Ruedemann, 1912)* S Kokomo, Indiana
- † **Lamontopterus Waterston, 1979** **Silurian**
26. *Lamontopterus knoxae* (Lamont, 1955)* S Pentland Hills, Scotl.
- † **HARDIEOPTERIDAE Tollerton, 1989** **Silurian – Devonian**
- † **Hallipterus Kjellesvig-Waering, 1963a** **Devonian**
27. *Hallipterus excelsior* (Hall, 1884a)* D New York
- i. = *Dolichocephala lacoana* Claypole, 1883 D Pennsylvania
- † **Hardieopterus Waterston, 1979** **Silurian**
28. ?*Hardieopterus lanarkensis* Waterston, 1979 S Patrick Burn, Scotl.
29. *Hardieopterus macrophthalmus* (Laurie, 1892)* S Pentland Hills, Scotl.
30. *Hardieopterus megalops* (Salter, 1859) S Herefordshire, Engl.
31. *Hardieopterus myops* (Clarke, 1907) S eastern USA

- † **Tarsopterella Størmer, 1951** **Devonian**
 32. *Tarsopterella scotica* (Woodward, 1872)* D Mid. Valley Scotland
 i. = *?Eriopteris brewsteri* Woodward, 1864 D Mid. Valley Scotland
 ii. = *Stylonurus armatus* Page, 1867 D Mid. Valley Scotland
- † **HIBBERTOPTEROIDEA Kjellesvig-Waering, 1959** **Devonian – Permian**
 † **DREPTOPTERIDAE Kjellesvig-Waering, 1966a** **Silurian – Devonian**
 † **Drepanopterus Laurie, 1892** **Silurian – Devonian**
 33. *Drepanopterus abonensis* Simpson, 1951 D Portishead, England
 34. *Drepanopterus odontospathus* Lamsdell, 2013b D NW Territories, Can.
 35. *Drepanopterus pentlandicus* Laurie, 1892* S Pentland Hills, Scotl.
- † **HIBBERTOPTERIDAE Kjellesvig-Waering, 1959** **Devonian – Permian**
 = † **CYRTOCTENIDAE Waterston, Oelofsen & Oosthuizen, 1985**
- † **Campylocephalus Eichwald, 1860** **Carboniferous – Perm.**
 36. *Campylocephalus oculus* (Kutorga, 1838)* P Dourasovo, Russia
 37. *Campylocephalus permianus* (Ponomarenko, 1985) P Komi, Russia
 38. *?Campylocephalus salmi* Stur, 1877 C Ostrava, Czech Rep.
- † **Cyrtoctenus Størmer & Waterston, 1968** **Devonian – Carbon.**
 39. *Cyrtoctenus caledonicus* (Salter, 1863) C East Lothian, Scotl.
 40. *Cyrtoctenus dewalquei* (Fraipont, 1889) D Pont-de-Bonne, Belg.
 i. = *Eurypteris dewalquei* var. *longimanus* Fraipont,
 1889 D Pont-de-Bonne, Belg.
 41. *Cyrtoctenus dicki* (Peach, 1883) C Thurso, Scotland
 42. *Cyrtoctenus ostraviensis* (Augusta & Přibyl, 1951) C Ostrava, Czech Rep.
 43. *Cyrtoctenus peachi* Størmer & Waterston, 1968* C Berwickshire, Scotl.
 44. *Cyrtoctenus wittebergensis* Waterston, Oelofsen & Oosthuizen, 1985 ... C Cape Province
- † **Dunsophterus Waterston, 1968** **Carboniferous**
 45. *Dunsophterus stevensoni* (Etheridge Jr, 1877)* C Berwickshire, Scotl.
- † **Hastimima White, 1908** **Permian**
 46. *Hastimima whitei* White, 1908* P Brazil
- † **Hibbertopterus Kjellesvig-Waering, 1959** **Carboniferous – Perm.**
 47. *?Hibbertopterus hibernicus* (Baily, 1872) C Kiltorcan, Ireland
 48. *Hibbertopterus scouleri* (Hibbert, 1836)* C West Lothian, Scotl.
- † **Vernonopterus Waterston, 1957** **Carboniferous**
 49. *Vernonopterus minutisculptus* (Peach, 1907)* C Lanarkshire, Scotland
- † **MYCTEROPIDAE Cope, 1886** **Carboniferous – Perm.**
 = † **WOODWARDOPTERIDAE Kjellesvig-Waering, 1959**
- † **Megarachne Hünicken, 1980** **Carboniferous – Perm.**
 50. *Megarachne servinei* Hünicken, 1980* C–P Santa Rosa, Argen.
- † **Mycterops Cope, 1886** **Carboniferous**

51. ?*Mycterops blairi* Waterston, 1968 C Loanhead, Scotland
 52. *Mycterops matthieui* Pruvost, 1924 C Charleroi, Belgium
 53. *Mycterops ordinatus* Cope, 1886* C Channelton, PA
 54. ?*Mycterops whitei* Schram, 1984 C Crescent, Iowa
 † **Woodwardopterus Kjellesvig-Waering, 1959** **Carboniferous**
 55. *Woodwardopterus scabrosus* (Woodward, 1887)* C Glencartholm, Scotl.
- STYLONURINA *incertae sedis***
- † ***Stylonuroides* Kjellesvig-Waering, 1966a** **Silurian – Devonian**
 56. *Stylonuroides dolichopteroides* (Størmer, 1934b)* S Ringerike, Norway
 57. *Stylonuroides orientalis* Shpinev, 2012 D Lake Shunet, Siberia
- † **EURYPTERINA Burmeister, 1843** **Ordovician – Permian**
 † **ONYCHOPTERELLOIDEA Lamsdell, 2011** **Ordovician–Silurian**
 † **ONYCHOPTERELLIDAE Lamsdell, 2011** **Ordovician–Silurian**
 † ***Onychopterella* Størmer, 1951** **Ordovician–Silurian**
 58. *Onychopterella augusti* Braddy, Aldridge & Theron, 1995 O Soom Shale, S. Afr.
 59. *Onychopterella kokomoensis* (Miller & Gurley, 1896)* S Kokomo, Indiana
 i. = *Eurypterus ranilarva* Clarke & Ruedemann, 1912 S Kokomo, Indiana
 60. ?*Onychopterella pumilus* (Savage, 1916) S Essex, Illinois
 † ***Tylopterella* Størmer, 1951** **Silurian**
 61. *Tylopterella boylei* (Whiteaves, 1884) S Ontario, Canada
 62. ?*Tylopterella menneri* (Novojilov, 1959) D Taimyr, Russia
- † **MOSELOPTEROIDEA Lamsdell, Braddy & Tetlie, 2010** **Silurian – Devonian**
 † **MOSELOPTERIDAE Lamsdell, Braddy & Tetlie, 2010** **Devonian**
 † ***Moselopterus* Størmer, 1974** **Devonian**
 63. *Moselopterus ancyloptelson* Størmer, 1974* D Alken an der Mosel
 64. *Moselopterus elongatus* Størmer, 1974 D Alken an der Mosel
 65. *Moselopterus lancmani* (Delle, 1937) D Plavinas, Latvia
 † ***Stoermeropterus* Lamsdell, 2011** **Silurian**
 66. *Stoermeropterus conicus* (Laurie, 1892)* S Pentland Hills
 i. = *Drepanopterus bembycoides* Laurie, 1899 S Pentland Hills
 ii. = *Drepanopterus lobatus* Laurie, 1899 S Pentland Hills
 67. *Stoermeropterus latus* (Størmer, 1934b) S Ringerike, Norway
 68. *Stoermeropterus nodosus* (Kjellesvig-Waering & Leutze, 1966) S Bass, West Virginia
 † ***Vinetopterus* Poschmann & Tetlie, 2004** **Devonian**
 69. *Vinetopterus martini* Poschmann & Tetlie, 2004 D Westerwald, Germ.
 70. *Vinetopterus struvei* (Størmer, 1974)* D Alken an der Mosel
- † **MEGALOGRAPTOIDEA Caster & Kjellesvig-Waering, 1955** **Ordovician**
 † **MEGALOGRAPTIDAE Caster & Kjellesvig-Waering, 1955** **Ordovician**

- † ***Echinognathus* Walcott, 1882** **Ordovician**
71. *Echinognathus clevelandi* Walcott, 1882* O New York
- † ***Megalograptus* Miller, 1874** **Ordovician**
72. *Megalograptus alveolatus* (Shuler, 1915) O Virginia
73. *Megalograptus ohioensis* Caster & Kjellesvig-Waering, 1955 O Ohio
74. *Megalograptus shideleri* Caster & Kjellesvig-Waering, 1964 O Ohio
75. *Megalograptus welchi* Miller, 1874* O Ohio
76. *Megalograptus williamsae* Caster & Kjellesvig-Waering, 1964 O Ohio
- † **'EURYPTEROIDEA' Burmeister, 1843** **Ordovician – Devonian**
- NB: Lamsdell *et al.* (2013) questioned the monophyly of this superfamily
- Family uncertain
- † ***Pentlandopterus* Lamsdell, Hoşgör & Selden, 2013** **Ordovician**
77. *Pentlandopterus minor* (Laurie, 1899)* S Pentland Hills, Scotl.
- † ***Paraeurypterus* Lamsdell, Hoşgör & Selden, 2013** **Ordovician**
78. *Paraeurypterus anatoliensis* Lamsdell, Hoşgör & Selden, 2013* O Şort Tepe, Turkey
- † **DOLICHOPTERIDAE Kjellesvig-Waering & Størmer, 1952** **Silurian – Devonian**
- † ***Clarkeipterus* Kjellesvig-Waering, 1966 [a/b?]** **Silurian**
79. *Clarkeipterus ?otisius* (Clarke, 1907) S eastern USA
80. *Clarkeipterus testudineus* (Clarke & Ruedeman, 1912)* S New York
- † ***Dolichopterus* Hall, 1859** **Silurian**
81. *Dolichopterus gotlandicus* Kjellesvig-Waering, 1979 S Gotland, Sweden
82. *Dolichopterus jewetti* Caster & Kjellesvig-Waering, 1956 S New York
83. *Dolichopterus macrocheirus* Hall, 1859* S New York / Canada
84. *Dolichopterus siluriceps* Clarke & Ruedemann, 1912 S New York / Canada
- † ***Ruedemannipecterus* Kjellesvig-Waering, 1966** **Silurian**
85. *Ruedemannipecterus stylonuroides* (Clarke & Ruedemann, 1912)* S Otisville, New York
- † **EURYPTERIDAE Burmeister, 1843** **Silurian**
- † ***Eurypterus* de Kay, 1825** **Silurian**
- = † *Baltoeurypterus* Størmer, 1973
86. *?Eurypterus cephalaspis* Salter, 1856 S Herefordshire, Engl.
87. *Eurypterus dekayi* Hall, 1859 S New York / Ontario
88. *Eurypterus flintstonensis* Swartz, 1923 S eastern USA
89. *Eurypterus hankeni* Tetlie, 2006a S Ringerike, Norway
90. *Eurypterus henningsmoeni* (Tetlie, 2002) S Bærum, Norway
91. *Eurypterus laculatus* Kjellesvig-Waering, 1958 S New York / Ontario
92. *Eurypterus lacustris* Harlan, 1834 S New York / Ontario
- i. = *Eurypterus pachycheirus* Hall, 1859 S New York / Ontario
- ii. = *Eurypterus robustus* Hall, 1859 S New York / Ontario
93. *Eurypterus leopoldi* Tetlie, 2006a S Somerset Is., Canada

94. *Eurypterus megalops* Clarke & Ruedemann, 1912 S New York
95. *Eurypterus ornatus* Leutze, 1958 S Fayette, Ohio
96. *Eurypterus pittsfordensis* Sarle, 1903 S Pittsford, New York
97. *Eurypterus quebecensis* Kjellesvig-Waering, 1958 S Québec, Canada
98. *Eurypterus remipes* DeKay, 1825* S New York / Ontario
- i. = *Carcinosoma trigona* (Ruedemann, 1916)..... S New York
99. *Eurypterus serratus* (Jones & Woodward, 1888) S Gotland, Sweden
100. *Eurypterus tetragonophthalmus* Fischer, 1839 S Saaremaa, Estonia
- i. = *Eurypterus fischeri* Eichwald, 1854 S Estonia / Ukraine
- ii. = *Eurypterus fischeri* var. *rectangularis* Schmidt, 1883...S Saaremaa, Estonia
- † **ERIEOPTERIDAE Tollerton, 1989** **Silurian – Devonian**
- † ***Erieopterus* Kjellesvig-Waering, 1958** **Silurian – Devonian**
101. *Erieopterus eriensis* (Whitfield, 1882)..... S Ohio
102. *Erieopterus hypsophthalmus* Kjellesvig-Waering, 1958..... S Ohio
103. ?*Erieopterus limuloides* (Kjellesvig-Waering, 1948a) S Kokomo, Indiana
104. *Erieopterus microphthalmus* (Hall, 1859)*..... D New York / Canada
105. ?*Erieopterus phillipsensis* Copeland, 1971..... S Cornwallis Is. Canada
106. ?*Erieopterus statzi* Størmer, 1936a D Siegburg, Germany
107. ?*Erieopterus turgidus* Stumm & Kjellesvig-Waering, 1962 S Michigan
- † **STROBILOPTERIDAE Lamsdell & Selden, 2013** **Silurian – Devonian**
- † ***Buffalopterus* Kjellesvig-Waering & Heubusch, 1962** **Silurian**
108. *Buffalopterus pustulosus* (Hall, 1859)*..... S New York / Ontario
- i. = *Eurypterus giganteus* Pohlman, 1882..... S New York / Ontario
- ii. = *Pterygotus globicaudatus* Pohlman, 1882..... S New York / Ontario
- † ***Strobilopterus* Ruedemann, 1935** **Silurian – Devonian**
- = † *Syntomopterus* Kjellesvig-Waering, 1961 [preoccupied]
- = † *Syntomopterella* Tetlie, 2007 [replacement name]
109. *Strobilopterus laticeps* (Schmidt, 1883) S Saaremaa, Estonia
- i. = *Dolichopterus stoermeri* Caster & Kjellesvig-Waering,
 1956 S Saaremaa, Estonia
110. *Strobilopterus princetonii* (Ruedemann, 1934)* D Wyoming, USA
- i. = *Erieopterus latus* Ruedemann, 1935 D Wyoming, USA
111. *Strobilopterus proteus* Lamsdell & Selden, 2013 D Wyoming, USA
112. *Strobilopterus richardsoni* (Kjellesvig-Waering, 1961a*) D Ohio
- † **DIPLOPERCULATA Lamsdell, Hoşgör & Selden, 2013** **Ordovician – Devonian**
- † **MIXOPTEROIDEA Caster & Kjellesvig-Waering, 1955** **Ordovician – Devonian**
- CARCINOSOMATIDAE Størmer, 1934b** **Ordovician – Devonian**
- Carcinosoma* Claypole, 1890b** **Silurian**
- = † *Eurysoma* Claypole, 1890a [preoccupied]

113. ?*Carcinosoma harleyi* Kjellesvig-Waering, 1961*b* S England
114. *Carcinosoma libertyi* Copeland & Bolton, 1960 S Manitoulin I., Canada
115. *Carcinosoma newlini* (Claypole, 1890*a*)* S Kokomo, Indiana
 i. = *Carcinosoma ingens* Claypole, 1894 S Kokomo, Indiana
116. ?*Carcinosoma punctatum* (Salter in Huxley & Salter, 1859) S England
117. *Carcinosoma scorpioides* (Woodward, 1868)..... S Lesmahagow
 i. = *Pterygotus raniceps* Woodward, 1868 S Lesmahagow
118. *Carcinosoma scoticus* (Laurie, 1899)..... S Pentland Hills, Scotl.
119. ?*Carcinosoma spiniferum* Kjellesvig-Waering & Heubusch, 1962 S Pittsford, New York
- † ***Eocarcinosoma* Caster & Kjellesvig-Waering, 1964** **Ordovician**
120. *Eocarcinosoma batrachophthalmus* Caster & Kjellesvig-Waering,
 1964* O Ohio
- † ***Eusarcana* Strand, 1942** **Silurian – Devonian**
 = † *Eusarcus* Grote & Pitt, 1875 [preoccupied]
 = † *Paracarcinosoma* Caster & Kjellesvig-Waering, 1964
121. *Eusarcana acrocephalus* (Semper, 1898)..... S–D Barrandian area
122. *Eusarcana obesus* (Woodward, 1868)..... S Lesmahagow
123. *Eusarcana scorpionis* (Grote & Pitt, 1875)* S New York / Ontario
- † ***Rhinocarcinosoma* Novojilov, 1962** **Silurian**
124. *Rhinocarcinosoma cicerops* (Clarke, 1907) S Otisville, New York
125. *Rhinocarcinosoma dosonensis* Braddy, Selden & Doan Nhat, 2002 S Dô Son, Vietnam
126. *Rhinocarcinosoma vaningeni* (Clarke & Ruedemann, 1912)* S Clinton, New York
- † **MIXOPTERIDAE Caster & Kjellesvig-Waering, 1955** **Silurian**
 = † LANARKOPTERIDAE Tollerton, 1989
- † ***Lanarkopterus* Ritchie, 1968** **Silurian**
127. *Lanarkopterus dolichoschelus* (Størmer, 1936*b*)* S Scotland
- † ***Mixopterus* Ruedemann, 1921** **Silurian**
128. *Mixopterus kiaeri* Størmer, 1934*b* S Ringerike, Norway
129. *Mixopterus multispinosus* (Clarke & Ruedemann, 1912)* S New York
130. *Mixopterus simonsoni* Schmidt, 1883 S Saaremaa, Estonia
- † **‘WAERINGOPTEROIDEA’** **Silurian – Devonian**
- NB: Superfamily name appears to be derived from a thesis; a family Waeringopteridae has not been formally published
- † ***Grossopterus* Størmer, 1934*c*** **Devonian**
131. *Grossopterus overathi* (Gross, 1933)* D Overath
132. *Grossopterus inexpectans* (Ruedemann, 1921) D Gilboa
- † ***Orcanopterus* Stott, Tetlie, Braddy, Nowlan, Glasser & Devereux, 2005** **Ordovician**
133. *Orcanopterus manitoulinensis* Stott, Tetlie, Braddy, Nowlan, Glasser
 & Devereux, 2005* O Manitoulin I., Canada
- † ***Waeringopterus* Leutze, 1961** **Silurian**

134. *Waeringopterus apfeli* Leutze, 1961 S New York / Ontario
135. *Waeringopterus cumberlandicus* (Swartz, 1923)* S West Virginia
 i. = *Eurypterus swartzi* Kjellesvig-Waering, 1958 S West Virginia
- † **ADELOPHTHALMOIDEA Tollerton, 1989** **Devonian – Permian**
- † **ADELOPHTHALMIDAE Tollerton, 1989** **Devonian – Permian**
- † ***Adelophthalmus* Jordan in Jordan & von Mayer, 1854** **Devonian – Permian**
 = † *Lepidoderma* Reuss, 1855
 = † *Anthraconectes* Meek & Worthen, 1868 [a/b?]
 = † *Polyzosternites* Goldenberg, 1873
 = † *Glyptoscorpius* Peach, 1882
136. *Adelophthalmus approximatus* (Hall & Clarke, 1888) C Pennsylvania, USA
137. *Adelophthalmus asturica* (Melendez, 1971) C d'Ablana, Spain
138. *Adelophthalmus bradorensis* (Bell, 1922) C N. Campbelltown
139. *Adelophthalmus cambieri* (Pruvost, 1930) C Charleroi, Belgium
140. ?*Adelophthalmus carbonarius* (Chernyshev, 1933) C Donetsk, Ukraine
141. *Adelophthalmus chinensis* (Grabau, 1920) C–P Zhaozezhuang
142. *Adelophthalmus corneti* (Pruvost, 1939) C Quaregnon, Belgium
143. *Adelophthalmus douvillei* (de Lima, 1890) P Bussaco, Portugal
144. *Adelophthalmus dumonti* (Stainier, 1917) C Mechelen-sur-Meuse
145. *Adelophthalmus granosus* Jordan in Jordan & von Meyer, 1854* C Saarbrücken, Germ.
146. *Adelophthalmus imhofi* (Reuss, 1855) C Vlkys, Czech Rep.
147. *Adelophthalmus irinae* Shpinev, 2006 C Krasnoyarsk, Russia
148. *Adelophthalmus kidstoni* (Peach, 1888) C Radstock, England
149. ?*Adelophthalmus lohesti* (Dewalque in Fraipont 1889) D Pont de Bonne, Belg.
150. *Adelophthalmus luceroensis* Kues & Kietzke, 1981 P New Mexico
151. *Adelophthalmus mansfieldi* (Hall, 1877) C Pennsylvania
 i. = *Eurypterus stylus* Hall, 1884 C Pennsylvania
152. *Adelophthalmus mazonensis* (Meek & Worthen, 1868) C Illinois
153. *Adelophthalmus moyseyi* (Woodward, 1907a) C Ilkeston, Blaengarw
 i. = *Eurypterus derbiensis* Woodward, 1907a C Ilkeston, England
154. *Adelophthalmus nebraskensis* (Barbour, 1914) P Nebraska
155. *Adelophthalmus pennsylvanicus* (Hall, 1877) C Pennsylvania
156. ?*Adelophthalmus perornatus* (Peach, 1882) C Glencartholm, Scotl.
157. *Adelophthalmus pruvosti* Kjellesvig-Waering, 1948b C Lens, France
158. ?*Adelophthalmus raniceps* Goldenberg, 1873 C Saarbrücken, Germ.
159. *Adelophthalmus sellardsi* (Dunbar, 1924) P Elmo, Kansas
160. *Adelophthalmus sievertsi* (Størmer, 1969) D Willwerath, Germ.
 i. = ?*Eurypterus trapezoides* Størmer, 1974 D Nellenköpfchen, Ger.
161. *Adelophthalmus waterstoni* (Tetlie et al., 2004) D Kimberley, Australia
162. *Adelophthalmus wilsoni* (Woodward, 1888) C Radstock, England
163. *Adelophthalmus zadrai* Přibyl, 1952 C Moravo-Silesia

- † **Bassipterus Kjellesvig-Waering & Leutze, 1966** **Silurian**
 164. *Bassipterus virginicus* Kjellesvig-Waering & Leutze, 1966* S Bass, West Virginia
- † **Esyslopterus Tetlie & Poschmann, 2008** **Silurian**
 165. *Esyslopterus patteni* (Størmer, 1934d) S Saaremaa, Estonia
- † **Nanahughmilleria Kjellesvig-Waering, 1961b** **Silurian – Devonian**
 166. *Nanahughmilleria clarkei* Kjellesvig-Waering, 1964b S Otisville, New York
 167. *Nanahughmilleria norvegica* (Kiær, 1911)* S Ringerike, Norway
 i. = *Eurypterus minutus* Kiær, 1911 S Ringerike, Norway
 168. *Nanahughmilleria notosiberica* Shpinev, 2012 D Krasnoyarsk, Siberia
 169. ?*Nanahughmilleria prominens* (Hall, 1884b) S Cayuga, New York
 170. *Nanahughmilleria pygmaea* (Salter, 1859) S Herefordshire, Engl.
 171. ?*Nanahughmilleria schiraensis* (Pirozhnikov, 1957) D Khakassia, Russia
- † **Parahughmilleria Kjellesvig-Waering, 1961b** **Silurian – Devonian**
 172. *Parahughmilleria bellistriata* (Kjellesvig-Waering, 1950a) S West Virginia
 173. *Parahughmilleria hefteri* Størmer, 1973 D Rhenish Massif, Ge.
 174. *Parahughmilleria longa* Shpiney, 2012 D Lake Shunet, Siberia
 175. *Parahughmilleria maria* (Clarke, 1907) S New York
 176. *Parahughmilleria matarakensis* (Pirozhnikov, 1957) D Khakassia, Russia
 177. *Parahughmilleria salteri* Kjellesvig-Waering, 1961b* S Herefordshire, Engl.
- † **Pittsfordipterus Kjellesvig-Waering & Leutze, 1966** **Silurian**
 178. *Pittsfordipterus phelpsae* (Ruedemann, 1921)* S Pittsford, New York
- † **PTERYGOTIOIDEA Clarke & Ruedemann, 1912** **Silurian – Devonian**
- † **HUGHMILLERIIDAE Kjellesvig-Waering, 1951** **Silurian**
- † **Herefordopterus Tetlie, 2006b** **Silurian**
 179. *Herefordopterus banksii* (Salter, 1856)* S Herefordshire, Engl.
 i. = *Eurypterus acuminatus* Salter, 1859a S Herefordshire, Engl.
- † **Hughmilleria Sarle, 1903** **Silurian**
 180. *Hughmilleria shawangunk* Clarke, 1907 S eastern USA
 181. *Hughmilleria socialis* Sarle, 1903* S Pittsford, New York
 i. = *Hughmilleria robusta* Sarle, 1903 S Pittsford, New York
 182. *Hughmilleria wangi* Tetlie, Selden & Ren, 2007 S Hunan, China
- † **SLIMONIDAE Novojilov, 1968** **Silurian**
- † **Salteropterus Kjellesvig-Waering, 1951** **Silurian**
 183. *Salteropterus abbreviatus* (Salter, 1859)* S Herefordshire, Engl.
- † **Slimonia Page, 1856** **Silurian**
 184. *Slimonia acuminata* Salter, 1856* S Lesmahagow
 i. = *Himantopterus maximus* Salter, 1856 S Lesmahagow
 185. *Slimonia boliviana* Kjellesvig-Waering, 1973 S Cochambamba, Bol.
 186. *Slimonia dubia* Laurie, 1899 S Pentland Hills, Scotl.

- † **PTERYGOTIDAE Clarke & Ruedemann, 1912** **Silurian – Devonian**
 = † JAEKELOPTERIDAE Størmer, 1974
- † **Acutiramus Ruedemann, 1935** **Silurian – Devonian**
187. *Acutiramus bohemicus* (Barrande, 1872) S Barrandian area
 i. = *Pterygotus comes* Barrande, 1872 S Barrandian area
 ii. = *Pterygotus mediocris* Barrande, 1872 S Barrandian area
 iii. = *Pterygotus blahai* Semper, 1898 S Barrandian area
 iv. = *Pterygotus fissus* Seemann, 1906 S Barrandian area
188. *Acutiramus cummingsi* (Grote & Pitt, 1875) S USA / Canada
 i. = *Pterygotus acuticaudatus* Pohlman, 1882 S New York
 ii. = *Pterygotus buffaloensis* Pohlman, 1881 S New York
 iii. = *Pterygotus quadraticaudatus* Pohlman, 1882 S New York
189. *Acutiramus floweri* Kjellesvig-Waering & Caster, 1955 S Kenwood, New York
190. *Acutiramus macrophthalmus* (Hall, 1859)* S USA / Canada
 i. = *Pterygotus osborni* Hall, 1859 S New York
 ii. = *Pterygotus cobbi* var. *juvenis* Clarke & Ruedemann,
 1912 S New York
191. *Acutiramus perneri* Chlupáč, 1994 D Barrandian area
192. *Acutiramus perryensis* Leutze, 1958 S Ohio
193. *Acutiramus suwanneensis* Kjellesvig-Waering, 1955 S? Florida
- † **Ciurcopteris Tetlie & Briggs, 2009** **Silurian**
194. *Ciurcopteris sarlei* (Ciurca & Tetlie, 2007) S Pittsford, New York
195. *Ciurcopteris ventricosus* (Kjellesvig-Waering, 1948a)* S Kokomo, Indiana
- † **Erettopteris Salter in Huxley & Salter, 1859** **Silurian – Devonian**
 = † *Truncatiramus* Kjellesvig-Waering, 1961b
196. *Erettopteris bilobus* (Salter, 1856)* S Lesmahagow
 i. = *Eurypterus perornatus* Salter, 1856 S Lesmahagow
 ii. = *Pterygotus bilobus* var. *acidens* Woodward, 1878 S Lesmahagow
 iii. = *Pterygotus bilobus* var. *crassus* Woodward, 1878 S Lesmahagow
 iv. = *Pterygotus bilobus* var. *inornatus* Woodward, 1878... S Lesmahagow
 v. = *Pterygotus bilobus* var. *perornatus* Woodward, 1878. S Lesmahagow
 vi. = *Pterygotus perornatus* var. *plicatissimus* Salter in
 Huxley & Salter, 1859 S Lesmahagow
197. *Erettopteris brodiei* Kjellesvig-Waering, 1961b S Herefordshire, Engl.
198. *Erettopteris canadensis* (Dawson, 1879) S Ontario, Canada
199. *Erettopteris exophthalmus* Kjellesvig-Waering & Leutze, 1966 S Bass, West Virginia
200. *Erettopteris gigas* Salter in Huxley & Salter, 1859 S Herefordshire, Engl.
201. *Erettopteris globiceps* Clarke & Ruedemann, 1912 S eastern USA
202. *Erettopteris grandis* Pohlman, 1881 S New York
203. *Erettopteris holmi* (Størmer, 1934b) S Ringerike, Norway
204. *Erettopteris laticauda* Schmidt, 1883 S Saaremaa, Estonia

205. *Erettopterus marstoni* Kjellesvig-Waering, 1961*b* S England
206. *Erettopterus megalodon* Kjellesvig-Waering, 1961*b* S England
207. *Erettopterus osiliensis* Schmidt, 1883 S Saaremaa, Estonia
208. *Erettopterus saetiger* Kjellesvig-Waering, 1964*a* S Pennsylvania
209. *Erettopterus serratus* Kjellesvig-Waering, 1961*b* D Ohio
210. *Erettopterus spatulatus* Kjellesvig-Waering, 1961*b* S Herefordshire, Engl.
211. ?*Erettopterus vogti* Størmer, 1934*a* D Spitsbergen
212. *Erettopterus waylandsmithi* Kjellesvig-Waering & Caster, 1955 S Kenwood, New York
- † **Jaekelopterus Waterston, 1964** **Devonian**
213. *Jaekelopterus howelli* Kjellesvig-Waering & Størmer, 1952 D Wyoming
- i. = *Pterygotus mcgrewi* Kjellesvig-Waering & Richardson
 In Kjellesvig-Waering (1986) [*nomen nudum*] D Wyoming
214. *Jaekelopterus rhenaniae* (Jaekel, 1914)* D Rhenish Massif, Ger.
- † **Necrogammarus Woodward, 1870** **Silurian**
215. *Necrogammarus salweyi* Woodward, 1870 S Herefordshire, Engl.
- † **Pterygotus Agassiz, 1839** **Silurian – Devonian**
- = † *Curviramus* Reudemann, 1935
216. *Pterygotus anglicus* Agassiz, 1844* D Scotland, Canada
- i. = *Pterygotus atlanticus* Clarke & Ruedemann, 1912 D New Brunswick, Can.
- ii. = *Pterygotus minor* Woodward, 1864 D Scotland
217. *Pterygotus arcuatus* Salter *in* Huxley & Salter, 1859 S Herefordshire, Engl.
218. ?*Pterygotus australis* McCoy, 1899 S Melbourne, Australia
219. *Pterygotus barrandeii* Semper, 1898 S Barrandian area
- i. = *Pterygotus beraunensis* Semper, 1898 S Barrandian area
220. *Pterygotus bolivianus* Kjellesvig-Waering, 1964*a* D Belen, Bolivia
221. *Pterygotus carmani* Kjellesvig-Waering, 1961 D Ohio
222. *Pterygotus cobbi* Hall, 1859 S New York / Canada
223. *Pterygotus denticulatus* Kjellesvig-Waering, 1961*b* S Herefordshire, Engl.
224. *Pterygotus floridanus* Kjellesvig-Waering, 1950*b* D Florida
225. *Pterygotus gaspesiensis* Russell, 1953 D Québec, Canada
226. ?*Pterygotus grandidentatus* Kjellesvig-Waering, 1961*b* S England
227. ?*Pterygotus impacatus* Kjellesvig-Waering, 1964*a* S Saaremaa, Estonia
228. *Pterygotus kopaninensis* Barrande, 1872 S Barrandian area, Cz.
229. *Pterygotus lanarkensis* Kjellesvig-Waering, 1964*a* S Lesmahagow, Scotl.
230. *Pterygotus lightbodyi* Kjellesvig-Waering, 1961*b* S England
231. *Pterygotus ludensis* Salter *in* Huxley & Salter, 1859 S Herefordshire, Engl.
232. *Pterygotus marylandicus* Kjellesvig-Waering, 1964*a* S Maryland
233. *Pterygotus monroensis* Sarle 1902 S New York

EURYPTERIDA *incertae sedis*

- † **Dorfopterus Kjellesvig-Waering, 1955** **Devonian**
234. *Dorfopterus angusticollis* Kjellesvig-Waering, 1955* D Wyoming

† ?*Dolichopterus*

235. ?*Dolichopterus asperatus* Kjellesvig-Waering, 1961 [a/b?] D Ohio
 236. ?*Dolichopterus bulbosus* Kjellesvig-Waering, 1961*b* S Herefordshire, Engl.
 237. ?*Dolichopterus herkimereensis* Caster & Kjellesvig-Waering, 1956 S New York / Canada

† ?*Eurypterus*

238. ?*Eurypterus loi* Chang, 1957 [non eurypterid?] S Hubei, China
 239. ?*Eurypterus podolicus* Chernyshev, 1947 S Ukraine
 240. ?*Eurypterus satpaevi* Simorin, 1956 C Karaganda, Kazakh.
 241. ?*Eurypterus styliformis* Chang, 1957 [non eurypterid?] S Hubei, China
 242. ?*Eurypterus tschernyschevi* Simorin, 1956 C Karaganda, Kazakh.
 243. ?*Eurypterus yangi* Chang, 1957 [non eurypterid?] S Hubei, China

† *Holmipterus* Kjellesvig-Waering, 1979 **Silurian**

244. *Holmipterus suecicus* Kjellesvig-Waering, 1979 S Gotland, Sweden

† *Marsupipterus* Caster & Kjellesvig-Waering, 1955 **Silurian**

245. *Marsupipterus sculpturatus* Caster & Kjellesvig-Waering, 1955* S Herefordshire, Engl.

† ?*Nanahughmilleria*

246. ?*Nanahughmilleria lanceolata* Salter, 1856 S Lesmahagow
 i. = *Eurypterus chartarius* Salter, 1859 S Lesmahagow
 ii. = *Eurypterus linearis* Salter, 1859 S Lesmahagow

† ?*Salteropterus*

247. ?*Salteropterus longilabium* Kjellesvig-Waering, 1961*b* S Welsh Borderlands

† ?*Stylonurus*

248. ?*Stylonurus perspicillum* Størmer, 1969 D Willwerath, Germany

† *Unionopterus* Chernyshev, 1948 **Carboniferous**

249. *Unionopterus anastasiae* Chernyshev, 1948* C Kazakhstan

NOMINA DUBIA

1. *Bunodella horrida* Matthew, 1888 [non Xiphosura] S New Brunswick
 2. ?*Dunsophterus wrightianus* Dawson 1881 D New York
 3. *Eurypterella ornata* Matthew, 1888 C 'Fern Ledges'
 4. *Eurypterus potens* Hall, 1884 C Pennsylvania
 5. *Eurypterus pulicaris* Salter, 1863 D New Brunswick
 6. *Hastimima sewardi* Strand, 1926 D South Africa
 7. ?*Pterygotus formosus* Dawson, 1871 D Gaspé, Canada
 8. *Pterygotus nobilis* Barrande, 1872 S Barrandian area
 9. *Pterygotus siemiradzki* Strand, 1926 D Podolia, Ukraine
 10. *Pterygotus taurinus* Salter, 1868 S Ewyas Harold, Engl.
 11. ?*Slimonia stylops* Salter in Huxley & Salter, 1859 S Herefordshire, Engl.

NOMINA NUDA

1. *Baltoeurypterus latus* Hanken & Størmer, 1975 S Ringerike, Norway

NOMINA VANA

1. *Pterygotus problematicus* Agassiz, 1844 S United Kingdom

MISIDENTIFICATIONS

1. *Buffalopectus verrucosus* Kjellesvig-Waering & Heubusch, 1962 [crustacean] ... O New York
2. *Carcinosoma ?logani* (Williams, 1915) [crustacean] S Ontario, Canada
3. *Eurypterus (Stylonurus?) macCarthyi* Kjellesvig-Waering, 1934 [cephalopod] D Ludlowville, New York
4. *Eurypterus pugio* Barrande, 1872 [crustacean] S Barrandian area
5. *Eurypterus thomasi* Walter, 1924 [aglaspidid] C Wisconsin
6. *Kockurus grandis* Chlupáč, 1995 [aglaspidid] C central Bohemia
7. *Kodymirus vagans* Chlupáč & Havlíček, 1965 [aglaspidid] C central Bohemia
8. *Mazonipterus cyclophthalmus* Kjellesvig-Waering, 1963b [plant] C Mazon Creek
9. *Melbournopterus crossotus* Caster & Kjellesvig-Waering, 1953 [brachiopod] ... S Melbourne, Australia
10. *Pterygotus expectatus* Barrande, 1872 [crustacean] S Barrandian area
11. *Pterygotus (Curviramus) elleri* Ruedemann, 1935 [crustacean] D New York
12. *Pterygotus (Curviramus) montanensis* Ruedemann, 1935 [crustacean] D Montana
13. *Pterygotus (Leptocheles) leptodactylum* M'Coy, 1849 [crustacean] S Herefordshire, Engl.

PSEUDOFOSILS

1. *Brachyopterella magna* (Clarke & Ruedemann, 1912) O New York
2. *?Carcinosoma linguata* (Clarke & Ruedemann, 1912) O New York
3. *?Carcinosoma longiceps* (Clarke & Ruedemann, 1912) O New York
4. *Dolichopterus antiquus* Ruedemann, 1942 O New York
5. *Dolichopterus frankfortensis* (Clarke & Ruedemann, 1912) O New York
6. *Dolichopterus insolitus* Ruedemann, 1926 O New York
7. *?Dolichopterus stellatus* (Clarke & Ruedemann, 1912) O New York
8. *?Drepanopterus ruedemanni* (O'Connell, 1916) O New York
9. *?Eocarcinosoma breviceps* (Ruedemann, 1926) O New York
10. *Eocarcinosoma ruedemanni* (Flower, 1945) O New York
11. *Eocarcinosoma triangulatus* (Clarke & Ruedemann, 1912) O New York
12. *Erettopterus walcotti* (Ruedemann, 1926) O New York
13. *Erieopterus chadwicki* (Clarke & Ruedemann, 1912) O New York
14. *Erieopterus hudsonicus* (Ruedemann, 1934) O New York
15. *?Eurypterus decepiens* (Ruedemann, 1942) O New York
16. *Eurypterus indicus* Dubey, 1985 pC M. Pradesh, India
17. *?Eurypterus pristinus* (Clarke & Ruedemann, 1912) O New York
18. *Eurypterus vermai* Dubey, 1985 pC M. Pradesh, India
19. *Hughmilleria chiplonkari* Dubey, 1985 pC M. Pradesh, India
20. *Hughmilleria kilfoylei* Ruedemann, 1934 O New York
21. *Hughmilleria prisca* Ruedemann, 1934 O New York
22. *Hughmilleria uticana* Ruedemann, 1926 O New York
23. *Parastylonurus rusti* (Ruedemann, 1926) O New York

24. *Pterygotus deepkillensis* Ruedemann, 1934 O New York
25. *Pterygotus nasutus* Clarke & Ruedemann, 1912 O New York
26. ?*Pterygotus normanskillensis* Clarke & Ruedemann, 1912 O New York
27. *Ruedemannipterus breviceps* (Clarke & Ruedemann, 1912) O New York
28. *Ruedemannipterus latifrons* (Clarke & Ruedemann, 1912) O New York
29. *Stylonurella modestus* (Clarke & Ruedemann, 1912) O New York
30. *Stylonuroides limbatus* (Clarke & Rudemann, 1912) O New York
31. ?*Waeringopterus pristinus* (Ruedemann, 1942) O New York
32. *Waeringopterus prolificus* (Clarke & Ruedemann, 1912) O New York

no Recent species

SCORPIONES

117 currently valid species of fossil scorpion

SCORPIONES C. L. Koch, 1851	Silurian – Recent
† Pelson (Family) PROSCORPIIDAE Scudder, 1885	Silurian – Carbon.
= † ARCHAEOCTONIDAE Petrunkevitch, 1949	
= † HYDROSCORPIONIDAE Kjellesvig-Waering, 1986	
= † LABRIOSCORPIONIDAE Kjellesvig-Waering, 1986	
= † STOERMEROSCORPIONIIDAE Kjellesvig-Waering, 1986	
= † WAERINGOSCORPIONIDAE Størmer, 1970	
† Archaeoctonus Pocock, 1911	Carboniferous
1. <i>Archaeoctonus glaber</i> (Peach, 1883)*	C Glencartholm
† Hydroscorpius Kjellesvig-Waering, 1986	Devonian
2. <i>Hydroscorpius denisoni</i> Kjellesvig-Waering, 1986*	D Wyoming
† Labriscorpio Leary, 1980	Carboniferous
3. <i>Labriscorpio alliedensis</i> Leary, 1980*	C Illinois
† Proscorpius Whitfield, 1885b	Silurian
= † <i>Archaeophonus</i> Kjellesvig-Waering, 1966b	
= † <i>Stoermeroscorpio</i> Kjellesvig-Waering, 1986	
4. <i>Proscorpius osborni</i> (Whitfield, 1885a)*	S ‘Bertie Waterlime’
i. = <i>Archaeophonus eurypteroides</i> Kjellesvig-Waering,	
1966b*	S ‘Bertie Waterlime’
ii. = <i>Stoermeroscorpio delicatus</i> Kjellesvig-Waering, 1986	S ‘Bertie Waterlime’
† Pseudoarchaeoctonus Kjellesvig-Waering, 1986	Carboniferous
5. <i>Pseudoarchaeoctonus denticulatus</i> Kjellesvig-Waering, 1986*	C Glencartholm
† Waeringoscorpio Størmer, 1970	Devonian
6. <i>Waeringoscorpio hefteri</i> Størmer, 1970*	D Alken an der Mosel
7. <i>Waeringoscorpio westerwaldensis</i> Poschmann, Dunlop, Kamenz & Scholtz, 2008	D Westerwald
† BILOBOSTERNINA Kjellesvig-Waering, 1986 (suborder)	Silurian – Devonian
† BRANCHIOSCORPIONOIDEA Kjellesvig-Waering, 1986	Devonian
† BRANCHIOSCORPIONIIDAE Kjellesvig-Waering, 1986	Devonian
† Branchioscorpio Kjellesvig-Waering, 1986	Devonian
8. <i>Branchioscorpio richardsoni</i> Kjellesvig-Waering, 1986*	D Wyoming
† DOLICHOPHONIIDAE Petrunkevitch, 1953	Silurian
† Dolichophonus Petrunkevitch, 1949	Silurian

9. *Dolichophonus loudonensis* (Laurie, 1899)* S Pentland Hills
- † **HOLOSTERNINA Kjellesvig-Waering, 1986** **Devonian**
- † **ACANTHOSCORPIONOIDEA Kjellesvig-Waering, 1986** **Devonian**
- † **ACANTHOSCORPIONIIDAE Kjellesvig-Waering, 1986** **Devonian**
- † ***Acanthoscorpio* Kjellesvig-Waering, 1986** **Devonian**
10. *Acanthoscorpio mucronatus* Kjellesvig-Waering, 1986* D Wyoming
- † **STENOSCORPIONIIDAE Kjellesvig-Waering, 1986** **Triassic**
- † ***Stenoscorpio* Kjellesvig-Waering, 1986** **Triassic**
11. *Stenoscorpio gracilis* (Wills, 1910)* Tr Keuper sandstone
12. *Stenoscorpio pseudogracilis* (Wills, 1947) Tr Keuper sandstone
- † **ALLOPALAEOPHONOIDEA Kjellesvig-Waering, 1986** **Silurian**
- † **ALLOPALAEOPHONIDAE Kjellesvig-Waering, 1986** **Silurian**
- † ***Allopalaeophonus* Kjellesvig-Waering, 1986** **Silurian**
13. *Allopalaeophonus caledonicus* (Hunter, 1886)* S Logan Water
- i. = *Palaeophonus hunteri* Pocock, 1901 S Logan Water
- † **EOCTONOIDEA Kjellesvig-Waering, 1986** **Carboniferous**
- † **ALLOBUTHISCORPIIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † ***Aspiscorpio* Kjellesvig-Waering, 1986** **Carboniferous**
14. *Aspiscorpio eageri* Kjellesvig-Waering, 1986* C Sparth Bottoms
- Aspiscorpio* sp. in Poschmann (2009) C Saar
- † **ANTHRACOSCORPIONIDAE Frič, 1904** **Carboniferous**
- † ***Allobuthus* Kjellesvig-Waering, 1986** **Carboniferous**
15. *Allobuthus pescei* (Vachon & Heyler, 1985)* C Montceau-les-Mines
- † ***Anthracoscorpio* Kušta, 1885** **Carboniferous**
16. *Anthracoscorpio dunlopi* Pocock, 1911 C Airdrie
17. *Anthracoscorpio juvenis* Kušta, 1885* C Rakovník
- † **BUTHISCORPIIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † ***Buthiscorpius* Petrunkevitch, 1953** **Carboniferous**
18. *Buthiscorpius lemayeri* Kjellesvig-Waering, 1986 C Illinois
- † **EOCTONIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † ***Eoctonus* Petrunkevitch, 1913** **Carboniferous**
19. *Eoctonus miniatus* Petrunkevitch, 1913* C Mazon Creek
- † **GARNETTIIDAE Dubinin, 1962** **Carboniferous**
- † ***Garnettius* Petrunkevitch, 1953** **Carboniferous**

20. *Garnettius hungerfordi* (Elias, 1936)* C Garnett, Kansas
- † **GIGANTOSCORPIONOIDEA Kjellesvig-Waering, 1986** **Devonian – Carbon.**
- † **GIGANTOSCORPIONIDAE Kjellesvig-Waering, 1986** **Devonian – Carbon.**
 = † PETALOSCORPIONIDAE Kjellesvig-Waering, 1986
- † ***Gigantoscopus* Størmer, 1963** **Carboniferous**
 21. *Gigantoscopus willsi* Størmer, 1963* C Glencartholm
- † ***Petaloscopus* Kjellesvig-Waering, 1986** **Devonian**
 22. *Petaloscopus bureaui* Kjellesvig-Waering, 1986* D Miguasha, Quebec
- † **MESOPHONOIDEA Wills, 1910** **Carbon. – Triassic**
- † **CENTROMACHIDAE Petrunkevitch, 1953** **Carboniferous**
 = † ANTHRACOCOAERILIDAE Kjellesvig-Waering, 1986
 = † PHOXISCORPIONIDAE Kjellesvig-Waering, 1986
- † ***Anthracochaerilus* Kjellesvig-Waering, 1986** **Carboniferous**
 23. *Anthracochaerilus palustris* Kjellesvig-Waering, 1986* C Glencartholm
- † ***Centromachus* Thorell & Lindström, 1885** **Carboniferous**
 24. *Centromachus euglyptus* (Peach, 1883)* C Glencartholm
- † ***Phoxiscopus* Kjellesvig-Waering, 1986** **Carboniferous**
 25. *Phoxiscopus peachi* Kjellesvig-Waering, 1986* C Dalmeny, Edinburgh
- † ***Pulmonoscorpio* Jeram, 1994a** **Carboniferous**
 26. *Pulmonoscorpius kirktonensis* Jeram, 1994a* C East Kirkton
- † **GALLIOSCORPIONIDAE Lourenço & Gall, 2004** **Triassic**
- † ***Gallioscorpio* Lourenço & Gall, 2004** **Triassic**
 27. *Gallioscorpio voltzi* Lourenço & Gall, 2004* Tr Vosges, France
- † **HELOSCORPIONIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † ***Heloscorpio* Kjellesvig-Waering, 1986** **Carboniferous**
 28. *Heloscorpio sutcliffei* (Woodward, 1907b)* C Sparth Bottoms
- † **MAZONIIDAE Petrunkevitch, 1913** **Carboniferous**
- † ***Mazonia* Meek & Worthen, 1868b** **Carboniferous**
 29. *Mazonia wardingleyi* (Woodward, 1907b) C Sparth Bottoms
 30. *Mazonia woodiana* Meek & Worthen, 1868b* C Mazon Creek
- † **MESOPHONIDAE Wills, 1910** **Triassic**
- † ***Mesophonus* Wills, 1910** **Triassic**
 31. *Mesophonus perornatus* Wills, 1910* Tr Keuper sandstone
 i. = *Mesophonus opisthophthalmus* Wills, 1947 Tr Keuper sandstone
 32. ?*Mesophonus pulcherrimus* Wills, 1910 Tr Keuper sandstone
 33. ?*Mesophonus pulcherrimus immaculatus* Wills, 1947 Tr Keuper sandstone

- † **WILLSISCORPIONIDAE** Kjellesvig-Waering, 1986 **Triassic**
- † *Willsiscorpio* Kjellesvig-Waering, 1986 **Triassic**
34. *Willsiscorpio bromsgroviensis* (Wills, 1910)* Tr Keuper sandstone
- † **PALAEOSCORPOIDEA** Lehmann, 1944 **Devonian – Triassic**
- † **PALAEOSCORPIONIDAE** Lehmann, 1944 **Devonian**
- † *Palaeoscorpio* Lehmann, 1944 **Devonian**
35. *Palaeoscorpius devonicus* Lehmann, 1944* D Hünsruckschiefer
- [NB: Kühl *et al.* (2012) simply list the genus unplaced under Protoscorpionina.]
- † **SPONGIOPHONOIDEA** Kjellesvig-Waering, 1986 **Devonian –Triassic**
- † **PRAERCTURIDAE** Kjellesvig-Waering, 1986 **Devonian**
- † *Praearcturus* Woodward, 1871a **Devonian**
36. *Praearcturus gigas* Woodward, 1871a* D Rowlestone
- † **SPONGIOPHONIDAE** Kjellesvig-Waering, 1986 **Triassic**
- † *Spongiophonus* Wills, 1947 **Triassic**
37. *Spongiophonus pustulosus* Wills, 1947* Tr Keuper sandstone
- † **MERISTOSTERNINA** Kjellesvig-Waering, 1986 **Carboniferous**
- † **CYCLOPHTHALMOIDEA** Thorell & Lindström, 1885 **Carboniferous**
- † **CYCLOPHTHALMIDAE** Thorell & Lindström, 1885 **Carboniferous**
- † *Cyclophthalmus* Corda, 1835 **Carboniferous**
38. *Cyclophthalmus senior* Corda, 1835* C Cholme
39. *Cyclophthalmus robustus* Kjellesvig-Waering, 1986 C Coseley
40. ?*Cyclophthalmus sibiricus* Novojilov & Størmer, 1963 C Kemerov Region
- † **MICROLABIIDAE** Kjellesvig-Waering, 1986 **Carboniferous**
- † *Microlabis* Corda, 1839 **Carboniferous**
41. *Microlabis sternbergii* Corda, 1839* C Cholme
- † **PALAEOBUTHOIDEA** Kjellesvig-Waering, 1986 **Carboniferous**
- † **PALAEOBUTHIDAE** Kjellesvig-Waering, 1986 **Carboniferous**
- † *Palaeobuthus* Petrunkevitch, 1913 **Carboniferous**
- = † *Mazoniscorpio* Wills, 1960
42. *Palaeobuthus distinctus* Petrunkevitch, 1913* C Mazon Creek
- i. = *Mazoniscorpio mazonensis* Wills, 1960 C Mazon Creek
- † **LOBOSTERNINA** Pocock, 1911 **Silurian – Carbon.**
- † **ISOBUTHOIDEA** Petrunkevitch, 1913 **Carboniferous**
- † **EOBUTHIDAE** Kjellesvig-Waering, 1986 **Carboniferous**

† <i>Eobuthus</i> Frič, 1904	Carboniferous
43. <i>Eobuthus cordai</i> Kjellesvig-Waering, 1986	C Kralupy Hill
44. <i>Eobuthus holti</i> Pocock, 1911	C Sparth Bottoms
45. <i>Eobuthus rakovnicensis</i> Frič, 1904*	C Rakovník
† EOSCORPIIDAE Scudder, 1884	Carboniferous
† <i>Eoscorpius</i> Meek & Worthen, 1868a	Carboniferous
= † <i>Alloscorpius</i> Petrunkevitch, 1949	
= † <i>Europhthalmus</i> Petrunkevitch, 1949	
= † <i>Lichnophthalmus</i> Petrunkevitch, 1949	
= † <i>Trigonoscorpio</i> Petrunkevitch, 1913	
= † <i>Typhloscorpius</i> Petrunkevitch, 1949	
46. <i>Eoscorpius bornaensis</i> Sterzel, 1918	C Chemnitz–Borna
47. <i>Eoscorpius carbonarius</i> Meek & Worthen, 1868a*	C Mazon Creek
i. = <i>Eoscorpius typicus</i> Petrunkevitch, 1913	C Mazon Creek
ii. = <i>Eoscorpius granulatus</i> Petrunkevitch, 1913	C Mazon Creek
iii. = <i>Trigonoscorpio americanus</i> Petrunkevitch, 1913	C Mazon Creek
48. <i>Eoscorpius casei</i> Kjellesvig-Waering, 1986	C Nova Scotia
49. <i>Eoscorpius distinctus</i> (Petrunkevitch, 1949)	C Coseley
50. <i>Eoscorpius mucronatus</i> Kjellesvig-Waering, 1986	C Barnsley
51. <i>Eoscorpius pulcher</i> (Petrunkevitch, 1949)	C Barnsley
i. = <i>Europhthalmus longimanus</i> Petrunkevitch, 1949	C Barnsley
52. <i>Eoscorpius sparthenensis</i> Baldwin & Sutcliffe, 1904	C Sparth Bottoms
† <i>Eskioscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
53. <i>Eskioscorpio parvus</i> Kjellesvig-Waering, 1986*	C Glencartholm
† <i>Trachyscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
54. <i>Trachyscorpio squarrosus</i> Kjellesvig-Waering, 1986*	C Fouldon
† ISOBUTHIDAE Petrunkevitch, 1913	Carbon. – Triassic
† <i>Boreoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
55. <i>Boreoscorpio copelandi</i> Kjellesvig-Waering, 1986*	C Nova Scotia
† <i>Bromsgroviscorpio</i> Kjellesvig-Waering, 1986	Triassic
56. <i>Bromsgroviscorpio willsi</i> Kjellesvig-Waering, 1986*	Tr Keuper sandstone
† <i>Feistmantelia</i> Frič, 1904	Carboniferous
57. <i>Feistmantelia ornata</i> Frič, 1904*	C Studnoves
† <i>Isobuthus</i> Frič, 1904	Carboniferous
58. <i>Isobuthus kralupensis</i> (Thorell & Lindström, 1885)*	C Kralup
59. ? <i>Isobuthus nyransensis</i> Frič, 1904	C Nýřany
† KRONOSCORPIONIDAE Kjellesvig-Waering, 1986	Carboniferous
† <i>Kronoscorpio</i> Kjellesvig-Waering, 1986	Carboniferous
60. <i>Kronoscorpio danielsi</i> (Petrunkevitch, 1913)*	C Mazon Creek

- † **PAREOBUTHIDAE Wills, 1959** **Carboniferous**
- † *Pareobuthus* Wills, 1959 **Carboniferous**
61. *Pareobuthus salopiensis* Wills, 1959* C Shropshire
- † **PARAISOBUTHOIDEA Kjellesvig-Waering, 1986** **Carboniferous**
- † **OPSIEOBUTHIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Opsieobuthus* Kjellesvig-Waering, 1986 **Carboniferous**
62. *Opsieobuthus pottsvillensis* (Moore, 1923)* C Indiana
- † **PARAISOBUTHIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Paraisobuthus* Kjellesvig-Waering, 1986 **Carboniferous**
63. *Paraisobuthus duobicarinatus* Kjellesvig-Waering, 1986 C Shipley
64. *Paraisobuthus frici* Kjellesvig-Waering, 1986 C Kralupy Hill
65. *Paraisobuthus prantli* Kjellesvig-Waering, 1986* C Rakovnik
66. *Paraisobuthus virginiae* Kjellesvig-Waering, 1986 C Mazon Creek
- † **SCOLOPOSCORPIONIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Benniescorpio* Wills, 1960 **Carboniferous**
67. *Benniescorpio tuberculatus* (Peach, 1883)* C Dysart, Fife
- † *Scoloposcorpio* Kjellesvig-Waering, 1986 **Carboniferous**
68. *Scoloposcorpio cramondensis* Kjellesvig-Waering, 1986* C Cramond, Edinburgh
- † **TELMATOSCORPIONIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Telmatoscorpio* Kjellesvig-Waering, 1986 **Carboniferous**
69. *Telmatoscorpio brevipectus* Kjellesvig-Waering, 1986* C Mazon Creek
- † **LOBOARCHAEOCTONOIDEA Kjellesvig-Waering, 1986** **Carboniferous**
- † **LOBOARCHAEOCTONIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Loboarchaeoctonus* Kjellesvig-Waering, 1986 **Carboniferous**
70. *Loboarchaeoctonus squamosus* Kjellesvig-Waering, 1986* C Glencarholm
- † **WATERSTONIIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † *Waterstonia* Kjellesvig-Waering, 1986 **Carboniferous**
71. *Waterstonia airdriensis* Kjellesvig-Waering, 1986* C Airdrie
72. ?*Waterstonia brachistodactyla* Kjellesvig-Waering, 1986 [claw only !] ... C Beith, Ayrshire
- † **PALAEOPHONOIDEA Thorell & Lindström, 1884** **Silurian**
- † **PALAEOPHONIDAE Thorell & Lindström, 1884** **Silurian**
- † *Palaeophonus* Thorell & Lindström, 1884 **Silurian**
73. *Palaeophonus nuncius* Thorell & Lindström, 1884* S Visby, Gotland
74. ?*Palaeophonus lightbodyi* Kjellesvig-Waering, 1954 [claw only !] S Ludford Lane

- ORTHOSTERNINA Pocock, 1911** **Carbon. – Recent**
Orthosternina incertae sedis
- † **Corniops Jeram, 1994b** **Carboniferous**
 75. *Corniops mapesii* Jeram, 1994b* C Lone Star Lake
- SCORPIONIOIDEA Latreille, 1802** **Carbon. – Recent**
- † **PALAEOPISTHACANTHIDAE Kjellesvig-Waering, 1986** **Carboniferous**
- † **Cryptoscorpium Jeram, 1994b** **Carboniferous**
 76. *Cryptoscorpium americanus* Jeram, 1994b* C Lone Star Lake
- † **Palaeopisthacanthus Petrunkevitch, 1913** **Carboniferous**
 77. *Palaeopisthacanthus schucherti* Petrunkevitch, 1913* C Mazon Creek
 78. *Palaeopisthacanthus vogelandurdeni* Jeram, 1994b C Lone Star Lake
- family uncertain**
- † **Compsoscorpium Petrunkevitch 1949** **Carboniferous**
 = † *Allobuthiscorpium* Kjellesvig-Waering, 1986
 = † *Coseleyscorpium* Kjellesvig-Waering, 1986
 = † *Leioscorpium* Kjellesvig-Waering, 1986
 = † *Lichnoscorpium* Petrunkevitch, 1949
 = † *Pseudobuthiscorpium* Kjellesvig-Waering, 1986
 = † *Typhlopisthacanthus* Petrunkevitch, 1949
79. *Compsoscorpium buthiformis* (Pocock, 1911)* C Coal Measures
 i. = *Typhlopisthacanthus anglicus* Petrunkevitch, 1949 ... C Coseley
 ii. = *Lichnoscorpium minutus* Petrunkevitch, 1949 C Coseley
 iii. = *Compsoscorpium elegans* Petrunkevitch 1949 C Coseley
 iv. = *Compsoscorpium elongatus* Petrunkevitch, 1949 C Coseley
 v. = *Buthiscorpium major* Wills, 1960 C Kilburn Coal
 vi. = *Leioscorpium pseudobuthiformis* Kjellesvig-Waering,
 1986 C Coseley
 vii. = *Pseudobuthiscorpium labiosus* Kjellesvig-Waering,
 1986 C Coseley
 viii. = *Coseleyscorpium lanceolatus* Kjellesvig-Waering, 1986 C Coseley
 ix. = *Allobuthus macrostethus* Kjellesvig-Waering, 1986C Coseley
- PSEUDOCHACTIDAE Gromov, 1998** **Recent**
 no fossil record
- BUTHOIDEA C. L. Koch, 1837** **Triassic – Recent**
- family uncertain**
- † **Palaeoburmesebuthus Lourenço, 2002** **Cretaceous**
 80. *Palaeoburmesebuthus grimaldii* Lourenço, 2002* K Myanmar amber

† ARCHAEOBUTHIDAE Lourenço, 2001	Cretaceous
† <i>Archaeobuthus</i> Lourenço, 2001	Cretaceous
81. <i>Archaeobuthus estephani</i> Lourenço, 2001*	K Lebanese amber
† PROTOBUTHIDAE Lourenço & Gall, 2004	Triassic
† <i>Protobuthus</i> Lourenço & Gall, 2004	Triassic
82. <i>Protobuthus elegans</i> Lourenço & Gall, 2004*	Tr Vosges
BUTHIDAE C. L. Koch, 1837	Palaeogene – Recent
= ANDROCTONIDAE C. L. Koch, 1837	
= MICROCHARMIDAE Lourenço, 1996a	
Centruroides Marx, 1890a	Neogene – Recent
83. <i>Centruroides nitidus</i> (Thorell, 1876a) [Recent]	Ne Dominican amber
i. = <i>Centruroides beynai</i> Schawaller, 1979a	Ne Dominican amber
Microcharmum Lourenço, 1995	Quaternary – Recent
84. <i>Microcharmum henderickxi</i> (Lourenço, 2009a)	Qt Madagascar copal
Microtityus Kjellesvig-Waering, 1966c	Neogene – Recent
85. <i>Microtityus ambarensis</i> (Schawaller, 1982a)	Ne Dominican amber
† Palaeoakentrobuthus Lourenço & Weitschat, 2000	Palaeogene
86. <i>Palaeoakentrobuthus knodeli</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
† Palaeoananteris Lourenço & Weitschat, 2001	Palaeogene
87. <i>Palaeoananteris ribnitiodamgartensis</i> Lourenço & Weitschat, 2001*	Pa Baltic amber
88. <i>Palaeoananteris ukrainensis</i> Lourenço & Weitschat, 2009	Pa Rovno amber
89. <i>Palaeoananteris wunderlichi</i> Lourenço, 2004	Pa Baltic amber
† Palaeoisometrus Lourenço & Weitschat, 2005a	Palaeogene
90. <i>Palaeoisometrus elegans</i> Lourenço & Weitschat, 2005a*	Pa Baltic amber
† Palaeogrosphus Lourenço, 2000a	Quaternary
91. <i>Palaeogrosphus copalensis</i> (Lourenço, 1996b)	Qt Copal
92. <i>Palaeogrosphus jacquesi</i> Lourenço & Henderickx, 2002	Qt Copal
† Palaeolychas Lourenço & Weitschat, 1996	Palaeogene
93. <i>Palaeolychas balticus</i> Lourenço & Weitschat, 1996*	Pa Baltic amber
94. <i>Palaeolychas weitschati</i> Lourenço, 2012	Pa Baltic amber
† Palaeoprotobuthus Lourenço & Weitschat, 2000	Palaeogene
95. <i>Palaeoprotobuthus pusillus</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
† Palaeospinobuthus Lourenço, Henderickx & Weitschat, 2005	Palaeogene
96. <i>Palaeospinobuthus cenozoicus</i> Lourenço, Henderickx &	
Weitschat, 2005*	Pa Baltic amber
† Palaeotityobuthus Lourenço & Weitschat, 2000	Palaeogene
97. <i>Palaeotityobuthus longiaculeus</i> Lourenço & Weitschat, 2000*	Pa Baltic amber
Tityus C. L. Koch, 1836	?Palaeogene – Recent
98. <i>Tityus azari</i> Lourenço, 2013	Ne Dominican amber

99. '*Tityus*' *eogenus* Menge, 1869 [presumably misplaced] Pa Baltic amber
100. *Tityus geratus* Santiago-Blay & Poinar, 1988 Ne Dominican amber
101. *Tityus (Brazilotityus) hartkorni* Lourenço, 2009b Ne Dominican amber
- † ***Uintascorpio* Perry, 1995** **Palaeogene**
102. *Uintascorpio halandrasorum* Perry, 1995* Pa Green River
- BUTHIDAE *incertae sedis***
103. '*Scorpio*' *schweiggeri* Holl, 1829 Qt Copal [not amber!]
- BOTHRIURIDAE Simon, 1880** **Recent**
- = TELEGONIDAE Peters, 1861 [based on a generic homonym]
- = ACANTHOCHIROIDAE Karsch, 1880b
- no fossil record
- CHACTOIDEA Pocock, 1893** **Cretaceous – Recent**
- † **PALAEOEUSCORPIDAE Lourenço, 2003** **Cretaceous**
- † ***Palaeoescorpius* Lourenço, 2003** **Cretaceous**
104. *Palaeoescorpius gallicus* Lourenço, 2003* K French amber
- CHACTIDAE Pocock, 1893** **Cretaceous – Recent**
- = BROTEIDAE Simon, 1879a [suppressed for lack of useage]
- † ***Araripescorpius* Campos, 1986** **Cretaceous**
105. *Araripescorpius ligabuei* Campos, 1986* K Crato Formation
- Chactas* Gervais, 1844** **Subrecent – Recent**
106. *Chactas pleistocenicus* Lourenço & Weitschat, 2005b Qt Colombian copal
- AKRAVIDAE Levy, 2007** **Recent**
- no fossil record
- CHAERILIDAE Pocock, 1893** **Cretaceous – Recent**
- Electrochaerilus* Santiago-Blay *et al.*, 2004** **Cretaceous**
107. *Electrochaerilus buckleyi* Santiago-Blay *et al.*, 2004 K Myanmar amber
- DIPLOCENTRIDAE Karsch, 1880b** **Recent**
- no fossil record
- EUSCORPIIDAE Laurie, 1896** **Recent**
- no fossil record
- HETEROSCORPIONIDAE Kraepelin, 1905** **Recent**
- no fossil record
- HEMISCORPIIDAE Pocock, 1893** **Cretaceous – Recent**
- = ISCHNURIDAE Simon, 1879a

- = LIOCHELIDAE Fet & Bechly, 2001
= †PROTOISCHNURIDAE Carvalho & Lourenço, 2001
- † ***Protoischnurus* Carvalho & Lourenço, 2001** **Cretaceous**
 108. *Protoischnurus axelrodorum* Carvalho & Lourenço, 2001* K Crato Formation
- IURIDAE Thorell, 1876b** **Recent**
 no fossil record
- SCORPIONIDAE Latreille, 1802** **Neogene – Recent**
 = PANDINOIDAE Thorell, 1876b
 = HETEROMETRIDAE Simon, 1879a
- † ***Mioscorpio* Kjellesvig-Waering, 1986** **Neogene**
 109. *Mioscorpio zeuneri* (Hadži, 1931)* Ne Swabian Alps
- † ***Sinoscorpius* Hong, 1983a** **Neogene**
 110. *Sinoscorpius shandongensis* Hong, 1983a* Ne Shandong, China
- SUPERSTITIONIIDAE Stahnke, 1940** **Recent**
 no fossil record
- TROGLOTAYOSICIDAE Lourenço, 1998** **Recent**
 no fossil record
- VAEJOVIDAE Thorell, 1876b** **Recent**
 no fossil record
- SCORPIONES *incertae sedis*
Scorpiones incertae sedis in Dunlop & Selden (2013) S Trecastle, Wales
- † ***Brontoscorpio* Kjellesvig-Waering, 1972** **Devonian**
 111. *Brontoscorpio anglicus* Kjellesvig-Waering, 1972 D England
- † ***Gymnoscorpius* Jeram, 1994b** **Carboniferous**
 112. *Gymnoscorpius mutillidigitatus* Jeram, 1994b* C northern England
- † ***Hubeiscorpio* Walossek, Li & Brauckmann, 1990** **Devonian**
 113. *Hubeiscorpio gracilitarsis* Walossek, Li & Brauckmann, 1990* D Hubei, China
- † ***Liasscorpionides* Bode, 1951** **Jurassic**
 114. *Liasscorpionides schmidtii* Bode, 1951* J Hondelage, Germany
- † ***Palaeomachus* Pocock, 1911** **Carboniferous**
 115. *Palaeomachus anglicus* (Woodward, 1876)* C Mansfield
- † ***Titanoscorpio* Kjellesvig-Waering, 1986** **Carboniferous**
 116. *Titanoscorpio douglassi* Kjellesvig-Waering, 1986 C Mazon Creek
- † ***Wattisonia* Wills, 1960** **Carboniferous**
 117. *Wattisonia coseleyensis* Wills, 1960 C Coseley

MISIDENTIFICATIONS

1. ?*Mesophonus maculatus* (Brauer, Redtenbacher & Ganglbauer, 1889)
 [?insect: cockroach] J Siberia
2. *Tiphoscorpio hueberi* Kjellesvig-Waering, 1986 [myriapod: *Eoarthroleura*] D New York

c. 2,000 Recent species

OPILIONES

34 currently valid species of fossil harvestman

OPILIONES Sundevall, 1833	Devonian – Recent
CYPHOPHTHALMI Simon, 1879a (suborder)	Cretaceous – Recent
NEOGOVEIDAE Shear, 1980	Recent
no fossil record	
OGOVEIDAE Shear, 1980	Recent
no fossil record	
PETTALIDAE Shear, 1980	Recent
no fossil record	
SIRONIDAE Simon, 1879a	Palaeogene – Recent
Siro Latreille, 1796	Palaeogene – Recent
1. <i>Siro balticus</i> Dunlop & Mitov, 2011	Pa Baltic amber
2. <i>Siro platypedibus</i> Dunlop & Giribet, 2003	Pa Bitterfeld amber
STYLOCELLIDAE Hansen & Sørensen, 1904	Cretaceous – Recent
† Palaeosiro Poinar, 2008	Cretaceous – Recent
3. <i>Palaeosiro burmanicum</i> Poinar, 2008	K Myanmar amber
NB: Originally described as a sironid, but regarded as a stylocellid by Giribet <i>et al.</i> (2012)	
TROGLOSIRONIDAE Shear, 1993	Recent
no fossil record	
EUPNOI Hansen & Sørensen, 1904 (suborder)	Devonian - Recent
plesion taxa	
† Eophalangium Dunlop, Anderson, Kerp & Hass, 2004	Devonian
4. <i>Eophalangium sheari</i> Dunlop, Anderson, Kerp & Hass, 2004*	D Rhynie chert
† Brigantibunum Dunlop & Anderson, 2005	Carboniferous
5. <i>Brigantibunum listoni</i> Dunlop & Anderson, 2005*	C East Kirkton
† Kustarachne Scudder, 1890b	Carboniferous
6. <i>Kustarachne tenuipes</i> Scudder, 1890b*	C Mazon Creek
i. = <i>Kustarachne exstincta</i> Melander, 1903	C Mazon Creek
ii. = <i>Kustarachne conica</i> Petrunkevitch, 1913	C Mazon Creek
† Macroglyion Garwood <i>et al.</i>, 2011	Carboniferous

7. *Macrogyion cronus* Garwood *et al.* 2011* C Montceau-les-Mines
- CADDOIDEA Banks, 1893** **Palaeogene – Recent**
- CADDIDAE Banks, 1893** **Palaeogene – Recent**
- Caddo Banks, 1892a** **Palaeogene – Recent**
8. *Caddo dentipalpus* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- PHALANGIOIDEA Latreille, 1802** **Palaeogene – Recent**
- family uncertain
- † *Petrunkevitchiana* Mello-Leitão, 1937 [genus *incertae sedis*] **Palaeogene**
9. *Petrunkevitchiana oculata* (Petrunkevitch, 1922)* Pa Florissant
- MONOScutIDAE Forster, 1948** **Recent**
- no fossil record
- NEOPILIONIDAE Lawrence, 1931** **Recent**
- no fossil record
- PHALANGIIDAE Latreille, 1802** **Palaeogene – Recent**
- Dicranopalpus* Doleschall, 1852** **Palaeogene – Recent**
10. *Dicranopalpus ramiger* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- i. = *Opilio corniger* Menge, 1854 Pa Baltic amber
- ii. = *Dicranopalpus palmnickensis* Roewer, 1939 Pa Baltic amber
- † ***Stephanobunus* Dunlop & Mammitzsch, 2010** **Palaeogene**
11. *Stephanobunus mitovi* Dunlop & Mammitzsch, 2010* Pa Baltic amber
- ?Phalangiidae**
12. *Opilio ovalis* C. L. Koch & Berendt, 1854 Pa Baltic amber
- [probably misplaced at genus level]
- SCLEROSOMATIDAE Simon, 1879a** **Jurassic – Recent**
- † ***Amauropilio* Mello-Leitão, 1937** **Palaeogene**
13. *Amauropilio atavus* (Cockerell, 1907) Pa Florissant
14. *Amauropilio lacoei* (Petrunkevitch, 1922) Pa Florissant
- Leiobunum* C. L. Koch, 1839a** **Jurassic – Recent**
15. *Leiobunum longipes* Menge, 1854 Pa Baltic amber
- i. = *Leiobunum saporum* Menge, 1854 [?*lapsus*] Pa Baltic amber
- ii. = *Leiobunum inclusum* Roewer, 1939 Pa Baltic amber
- † ***Mesobunus* Huang, Selden & Dunlop, 2009** **Jurassic**
16. *Mesobunus dunlopi* Giribet, Tourhino, Shih & Ren, 2012 J Daohugou
17. *Mesobunus martensi* Huang, Selden & Dunlop, 2009* J Daohugou

Family uncertain

- † ***Daohugopilio* Huang, Selden & Dunlop, 2009** **Jurassic**
 18. *Daohugopilio sheari* Huang, Selden & Dunlop, 2009* J Daohugou
- DYSPNOI Hansen & Sørensen, 1904 (suborder)** **Carbon. – Recent**
 family uncertain
- † ***Ameticos* Garwood *et al.*, 2011** **Carboniferous**
 19. *Ameticos scolos* Garwood *et al.* 2011* C Montceau-les-Mines
- † ***Echinopustulatus* Dunlop, 2004** **Carboniferous**
 20. *Echinopustulatus samuelnelsoni* Dunlop, 2004* C Missouri
- ISCHYROPSALIDOIDEA Simon, 1879a** **Palaeogene – Recent**
 Tentative assignment, family uncertain
- † ***Piankhi* Dunlop, Bartel & Mitov, 2012** **Palaeogene**
 21. *Piankhi steineri* Dunlop, Bartel & Mitov, 2012* Pa Baltic amber
- CERATOLASMATIDAE Shear, 1986** **Recent**
 no fossil record
- ISCHYROPSALIDIDAE Simon, 1879a** **Recent**
 no fossil record
- SABACONIDAE Dresco, 1970** **Palaeogene – Recent**
***Sabacon* Simon, 1879a** **Palaeogene – Recent**
 22. *Sabacon claviger* (Menge, 1854) Pa Baltic amber
 i. = *Sabacon bachofeni* Roewer, 1939 Pa Baltic amber
- TROGULOIDEA Sundevall, 1833** **Cretaceous – Recent**
 [family uncertain; Shear (2010) suggested it is not an ortholasmatine, but may represent a new family]
- † ***Halitherses* Giribet & Dunlop, 2005** **Cretaceous**
 23. *Halitherses grimaldii* Giribet & Dunlop, 2005* K Myanmar amber
- DICRANOLASMATIDAE Simon, 1879a** **Recent**
 no fossil record
- † **EOTROGULIDAE Petrunkevitch, 1955a** **Carboniferous**
 † ***Eotrogulus* Thevenin, 1901** **Carboniferous**
 24. *Eotrogulus fayoli* Thevenin, 1901* C Commeny
- NEMASTOMATIDAE Simon, 1879a** **Palaeogene – Recent**
***Histicostoma* Kratochvíl, 1958** **Palaeogene – Recent**
 25. ?*Histicostoma tuberculatum* (C. L. Koch & Berendt, 1854) Pa Baltic amber
***Mitostoma* Roewer, 1951** **Palaeogene – Recent**

26. ?*Mitostoma denticulatum* (C. L. Koch & Berendt, 1854) Pa Baltic amber
 i. = *Nemastoma succineum* Roewer, 1939 Pa Baltic amber
Nemastoma C. L. Koch, 1836 **Palaeogene – Recent**
27. ?*Nemastoma incertum* C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **NEMASTOMOIDIDAE Petrunkevitch, 1955a** **Carboniferous**
 † *Nemastomoides* Thevenin, 1901 **Carboniferous**
 = † *Protopilio* Petrunkevitch, 1913
28. *Nemastomoides elaveris* Thevenin, 1901* C Commenry
 29. *Nemastomoides longipes* (Petrunkevitch, 1913) C Mazon Creek
- NIPPONOSALIDIDAE Martens, 1976** **Recent**
 no fossil record
- TROGULIDAE Sundevall, 1833** **Palaeogene – Recent**
Trogulus Latreille, 1802 **Palaeogene – Recent**
 30. *Trogulus longipes* Haupt, 1956 Pa Geiseltal
- LANIATORES Thorell, 1876c (suborder)** **Palaeogene – Recent**
 family uncertain
- Philacarus* Sørensen, 1932 **Neogene – Recent**
 31. *Philacarus hispaniolensis* Cokendolpher & Poinar, 1992 Ne Dominican amber
- INSIDIATORES Loman, 1900 (infraorder)** **Palaeogene – Recent**
TRAVUNIOIDEA Absolon & Kratochvíl, 1932 **Palaeogene – Recent**
CLADONYCHIDAE Hadži, 1935 **Palaeogene – Recent**
 † *Proholoscotolemon* Ubick & Dunlop, 2005 **Palaeogene**
 32. *Proholoscotolemon nemastomoides* (C. L. Koch & Berendt, 1854)* Pa Baltic amber
 ? *Proholoscotolemon* sp. in Ubick & Dunlop (2005) Pa Baltic amber
- PENTANYCHIDAE Briggs, 1971** **Recent**
 no fossil record
- TRAVUNIIDAE Absolon & Kratochvíl, 1932** **Recent**
 no fossil record
- TRIAENONYCHOIDEA Sørensen, 1886** **Recent**
SYNTHETONYCHIIDAE Forster, 1954 **Recent**
 no fossil record
- TRIAENONYCHIDAE Sørensen, 1886** **Recent**
 no fossil record

GRASSATORES Kury, 2002 (infraorder)	Neogene – Recent
SAMOIDEA Sørensen, 1886	Neogene – Recent
BIANTIDAE Thorell, 1889	Recent
no fossil record	
ESCADABIIDAE Kury & Pérez González <i>in</i> Kury, 2003	Recent
no fossil record	
KIMULIDAE Pérez González, Kury & Alonso-Zarazaga <i>in</i> Pérez González & Kury, 2007	Neogene – Recent
<i>Kimula</i> Goodnight & Goodnight, 1942	Neogene – Recent
<i>Kimula</i> sp. <i>in</i> Cokendolpher & Poinar (1992)	Ne Dominican amber
PODOCTIDAE Roewer, 1912	Recent
no fossil record	
SAMOIDEAE Sørensen, 1886	Neogene – Recent
<i>Hummelinckiolus</i> Šilhavý, 1979	Neogene – Recent
33. <i>Hummelinckiolus silhavyi</i> Cokendolpher & Poinar, 1998	Ne Dominican amber
<i>Pellobunus</i> Banks, 1905	Neogene – Recent
34. <i>Pellobunus proavus</i> Cokendolpher, 1987	Ne Dominican amber
STYGNOMMATIDAE Roewer, 1923	Recent
no fossil record	
ASSAMIOIDEA Sørensen, 1884	Recent
ASSAMIIDAE Sørensen, 1884	Recent
no fossil record	
EPEDANIDAE Sørensen, 1886	Recent
no fossil record	
PETROBUNIDAE Sharma & Giribet, 2011	Recent
no fossil record	
PYRAMIDOPIIDAE Sharma, Prieto & Giribet, 2011	Recent
no fossil record	
STYGNOPSIDAE Sørensen, 1932	Recent
no fossil record	
TITHAEIDAE Sharma & Giribet, 2011	Recent
no fossil record	

GONYLEPTOIDEA Sundevall, 1833	Recent
AGORISTENIDAE Šilhavý, 1973	Recent
no fossil record	
COSMETIDAE C. L. Koch, 1839a	Recent
no fossil record	
CRANAIIDAE Roewer, 1913	Recent
no fossil record	
GONYLEPTIDAE Sundevall, 1833	Recent
no fossil record	
MANAOSBIIDAE Roewer, 1943	Recent
no fossil record	
STYGNIDAE Simon, 1879b	Recent
no fossil record	
PHALANGODOIDEA Simon, 1879a	Recent
ONCOPODIDAE Thorell, 1876c	Recent
no fossil record	
PHALANGODIDAE Simon, 1879a	Recent
no fossil record	
ZALMOXOIDEA Sørensen, 1886	Recent
FISSIPHALLIIDAE Martens, 1988	Recent
no fossil record	
GUASINIIDAE González-Sponga, 1997	Recent
no fossil record	
ICALEPTIDAE Kury & Pérez González, 2002	Recent
no fossil record	
ZALMOXIDAE Sørensen, 1886	Recent
no fossil record	
OPILIONES <i>incertae sedis</i>	
unnamed specimen <i>in</i> Jell & Duncan (1986)	K Koonwarra

NOMINA DUBIA

1. *Cheiomachus coriaceus* Menge, 1854 Pa Baltic amber
2. *Phalangium succineum* Presl, 1822 Pa Baltic amber

MISIDENTIFICATIONS

1. *Hasseltides primigenius* Weyenbergh, 1869 [crinoid] J Solnhofen
2. *Rhabdotarachnoides simoni* Haupt, 1957 [plant fragment] P Rotliegend

6,491 Recent species according to Kury (2011)

PHALANGIOTARBIDA

31 currently valid species of fossil phalangiotarbid

- † **PHALANGIOTARBIDA Haase, 1890** Devonian – Permian
 = † ARCHITARBIDA Petrunkevitch, 1945a
- † **DEVONOTARBIDAE Poschmann & Dunlop, 2012** Devonian
- † ***Devonotarbus* Poschmann, Anderson & Dunlop, 2005** Devonian
1. *Devonotarbus hombachensis* Poschmann, Anderson & Dunlop, 2005* D Germany
- † **ANTHRACOTARBIDAE Kjellesvig-Waering, 1969** Carboniferous
- † ***Anthracotarbus* Kjellesvig-Waering, 1969** Carboniferous
2. *Anthracotarbus hintoni* Kjellesvig-Waering, 1969* C Oklahoma
- † **ARCHITARBIDAE Karsch, 1882** Carboniferous
 = † PHALANGIOTARBIDAE Haase, 1890
- † ***Architarbus* Scudder, 1868** Carboniferous
3. *Architarbus hoffmanni* Guthörl, 1934 C Saar basin
- i. = *Opiliotarbus kliveri* Waterlot, 1935 C Saar basin
- ii. = *Goniotarbus sarana* Guthörl, 1965 C Saar basin
4. *Architarbus minor* Petrunkevitch, 1913 C Mazon Creek
5. *Architarbus rotundatus* Scudder, 1868* C Mazon Creek
- † ***Bornatarbus* Rößler & Schneider, 1997** Carboniferous
6. *Bornatarbus mayasii* (Haupt in Nindel, 1955)* C Germany / UK
- † ***Discotarbus* Petrunkevitch, 1913** Carboniferous
7. *Discotarbus deplanatus* Petrunkevitch, 1913* C Mazon Creek
- † ***Geratarbus* Scudder, 1890b** Carboniferous
8. *Geratarbus lacoeyi* Scudder, 1890b* C Mazon Creek
9. *Geratarbus bohemicus* Petrunkevitch, 1953 C Nýřany
- † ***Goniotarbus* Petrunkevitch, 1949** Carboniferous
10. *Goniotarbus angulatus* (Pocock, 1911) C Coseley
11. *Goniotarbus tuberculatus* (Pocock, 1911)* C Coseley
- i. = *Goniotarbus tuberculatus* Petrunkevitch, 1949 C Coseley
- † ***Hadrachne* Melander, 1903** Carboniferous
12. *Hadrachne horribilis* Melander, 1903* C Mazon Creek
- † ***Leptotarbus* Petrunkevitch, 1945a** Carboniferous
13. *Leptotarbus torpedo* (Pocock, 1911)* C Coseley
- † ***Mesotarbus* Petrunkevitch, 1949** Carboniferous
14. *Mesotarbus angustus* (Pocock, 1911) C Coseley

15. <i>Mesotarbus eggintoni</i> (Pocock, 1911)	C Coseley
16. <i>Mesotarbus hindi</i> (Pocock, 1911)	C Coseley
17. <i>Mesotarbus intermedius</i> Petrunkevitch, 1949*	C Coseley
18. <i>Mesotarbus peteri</i> Dunlop & Horrocks, 1997	C Westhoughton
† Metatarbus Petrunkevitch, 1913	Carboniferous
19. <i>Metatarbus triangularis</i> Petrunkevitch, 1913*	C Mazon Creek
† Ootarbus Petrunkevitch, 1945a	Carboniferous
20. <i>Ootarbus pulcher</i> Petrunkevitch, 1945a*	C Mazon Creek
21. <i>Ootarbus ovatus</i> Petrunkevitch, 1945a	C Mazon Creek
† Orthotarbus Petrunkevitch, 1945a	Carboniferous
22. <i>Orthotarbus longipes</i> Simon, 1971	C Halleschen Mulde
23. <i>Orthotarbus minutus</i> (Petrunkevitch, 1913)*	C Mazon Creek
24. <i>Orthotarbus robustus</i> Petrunkevitch, 1945a	C Mazon Creek
25. <i>Orthotarbus nyranensis</i> Petrunkevitch, 1953	C Nýřany
† Paratarbus Petrunkevitch, 1945a	Carboniferous
26. <i>Paratarbus carbonarius</i> Petrunkevitch, 1945a*	C Mazon Creek
† Phalangiotarbus Haase, 1890	Carboniferous
27. <i>Phalangiotarbus subovalis</i> (Woodward, 1872b)*	C Burnley
† Pycnotarbus Darber, 1990	Carboniferous
28. <i>Pycnotarbus verrucosus</i> Darber, 1990*	C Oelsnitz
† Triangulotarbus Patrick, 1989	Carboniferous
29. <i>Triangulotarbus terrehautensis</i> Patrick, 1989*	C Indiana
† HETEROTARBIDAE Petrunkevitch, 1913	Carboniferous
† Heterotarbus Petrunkevitch, 1913	Carboniferous
30. <i>Heterotarbus ovatus</i> Petrunkevitch, 1913*	C Mazon Creek
† OPILIOTARBIDAE Petrunkevitch, 1945a	Carb. – Permian
† Opiliotarbus Pocock, 1910	Carb. – Permian
31. <i>Opiliotarbus elongatus</i> (Scudder, 1890b)*	C – P USA / Germany

NOMINA DUBIA

1. <i>Eotarbus litoralis</i> Kuřta, 1888	C Rakovník
2. <i>Nemastomoides depressus</i> Petrunkevitch, 1913	C Mazon Creek

no Recent species

PSEUDOSCORPIONES

44 currently valid species of fossil pseudoscorpion

PSEUDOSCORPIONES De Geer, 1778	Devonian – Recent
= CHERNETES Simon, 1879a	
† DRACOCHELIDAE Schawaller, Shear & Bonamo, 1991 (plesion family)	Devonian
† <i>Dracochela</i> Schawaller, Shear & Bonamo, 1991	Devonian
1. <i>Dracochela deprehendor</i> Schawaller, Shear & Bonamo, 1991*	D Gilboa
CHELONETHI Thorell, 1882	Cretaceous – Recent
EPIOCHIERATA Harvey, 1992	Cretaceous – Recent
CHTHONOIDEA Daday, 1888	Palaeogene – Recent
CHTHONIIDAE Daday, 1888	Palaeogene – Recent
<i>Chthonius</i> C. L. Koch, 1843a	Palaeogene – Recent
2. <i>Chthonius (Chthonius) mengei</i> Beier, 1937	Pa Baltic amber
3. <i>Chthonius (Chthonius) pristinus</i> Schawaller, 1978	Pa Baltic amber
<i>Pseudochthonius</i> Balzan, 1892	Neogene – Recent
4. <i>Pseudochthonius squamosus</i> Schawaller, 1980a	Ne Dominican amber
Tyrannchthonius Chamberlin, 1929	Quaternary – Recent
<i>Tyrannchthonius</i> sp. in Judson (2010)	Qt Madagascan copal
LECHYTIDAE Chamberlin, 1929	Neogene – Recent
<i>Lechyti</i> Balzan, 1892	Neogene – Recent
5. <i>Lechyti tertiaria</i> Schawaller, 1980a	Ne Dominican amber
TRIDENCHTHONIIDAE Balzan, 1892	Palaeogene – Recent
= DITHIDAE Chamberlin, 1929	
† <i>Chelignathus</i> Menge, 1854	Palaeogene
6. <i>Chelignathus kochii</i> Menge, 1854*	Pa Baltic amber
FEAELLOIDEA Ellingsen, 1906	Palaeogene – Recent
FEAELLIDAE Ellingsen, 1906	Recent
no fossil record	
PSEUDOGARYPIDAE Chamberlin, 1923a	Palaeogene – Recent
<i>Pseudogarypus</i> Ellingsen, 1909	Palaeogene – Recent
7. <i>Pseudogarypus extensus</i> Beier, 1937	Pa Baltic amber

8. <i>Pseudogarypus hemprichii</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
9. <i>Pseudogarypus minor</i> Beier, 1947a	Pa Baltic amber
10. <i>Pseudogarypus pangaea</i> Henderickx in Henderickx <i>et al.</i> , 2006.....	Pa Baltic amber
11. <i>Pseudogarypus synchrotron</i> Henderickx in Henderickx <i>et al.</i> , 2012	Pa Baltic amber
IOCHIERATA Harvey, 1992	Cretaceous – Recent
HEMICTENATA Balzan, 1892	Cretaceous – Recent
NEOBISIOIDEA Chamberlin, 1930	Cretaceous – Recent
BOCHICIDAE Chamberlin, 1930	Recent
= VACHONIIDAE Chamberlin, 1947	
no fossil record	
GYMNOBISIIDAE Beier, 1947b	Recent
no fossil record	
HYIDAE Chamberlin, 1930	Recent
no fossil record	
IDEORONCIDAE Chamberlin, 1930	Recent
no fossil record	
NEOBISIIDAE Chamberlin, 1930	Cretaceous – Recent
= OBISIIDAE Sundevall, 1833	
† <i>Electrobisium</i> Cockerell, 1917	Cretaceous
12. <i>Electrobisium acutum</i> Cockerell, 1917a*	K Myanmar amber
<i>Microcreagris</i> Balzan, 1892	Palaeogene – Recent
13. <i>Microcreagris koellnerorum</i> Schawaller, 1978	Pa Baltic amber
<i>Neobisium</i> Chamberlin, 1930	Palaeogene – Recent
14. <i>Neobisium (Neobisium) extinctum</i> Beier, 1955	Pa Baltic amber
15. <i>Neobisium henderickxi</i> Judson, 2003	Pa Baltic amber
<i>Roncus</i> L. Koch, 1873	Palaeogene – Recent
16. <i>Roncus succineus</i> Beier, 1955	Pa Baltic amber
PARAHYIDAE Harvey, 1992	Recent
no fossil record	
SYARINIDAE Chamberlin, 1930	Recent
no fossil record	
PANCTENATA Balzan, 1892	Cretaceous – Recent
GARYPOIDEA Simon, 1879a	Cretaceous – Recent
GARYPIDAE Simon, 1879a	Recent
= SYNSPHRONIDAE Beier, 1932a	

no fossil record

GARYPINIDAE Daday, 1888	Cretaceous – Recent
<i>Amblyolpium</i> Simon, 1898b	Cretaceous – Recent
17. <i>Amblyolpium burmiticum</i> (Cockerell, 1920)	K Myanmar amber
Garypinus Daday, 1888	Palaeogene – Recent
18. <i>Garypinus electri</i> Beier, 1937	Pa Baltic amber
GEOGARYPIDAE Chamberlin, 1930	Palaeogene – Recent
Geogarypus Chamberlin, 1930	Palaeogene – Recent
19. <i>Geogarypus gorskii</i> Henderickx, 2005	Pa Baltic amber
20. <i>Geogarypus macrodactylus</i> Beier, 1937	Pa Baltic amber
21. <i>Geogarypus major</i> Beier, 1937	Pa Baltic amber
LARCIDAE Harvey, 1992	Recent
no fossil record	
MENTHIDAE Chamberlin, 1930	Recent
no fossil record	
OLPIIDAE Banks, 1895	Palaeogene – Recent
no fossil record	
STERNOPHOROIDEA Chamberlin, 1923b	Neogene – Recent
STERNOPHORIDAE Chamberlin, 1923b	Neogene – Recent
<i>Idiogaryops</i> Hoff, 1963	Neogene – Recent
22. <i>Idiogaryops pumilus</i> (Hoff, 1963) [Recent]	Ne–R Dominican amber
CHEIRIDIOIDEA Hansen, 1894	Palaeogene – Recent
CHEIRIDIIDAE Hansen, 1894	Palaeogene – Recent
<i>Cheiridium</i> Menge, 1855	Palaeogene – Recent
23. <i>Cheiridium hartmanni</i> (Menge, 1854)	Pa Baltic amber
<i>Cryptocheiridium</i> Chamberlin, 1931a	Neogene – Recent
24. <i>Cryptocheiridium (Cryptocheiridium) antiquum</i> Schawaller, 1981	Ne Dominican amber
PSEUDOCHIRIDIIDAE Chamberlin, 1923b	Neogene – Recent
<i>Pseudochiridium</i> With, 1906	Neogene – Recent
25. <i>Pseudochiridium lindae</i> Judson, 2007	Ne Dominican amber
CHELIFEROIDEA Risso, 1826	Cretaceous – Recent
ATEMNIDAE Kishida, 1929	Palaeogene – Recent
Atemninae indet. <i>in</i> Judson (2010)	Qt Dominican amber
<i>Paratemnoides</i> Harvey, 1991	Quaternary – Recent

26. <i>Paratemnoides nidificator</i> (Balzan, 1888) [Recent]	Qt–R Colombian copal
† Progonatemnus Beier, 1955	Palaeogene
27. <i>Progonatemnus succineus</i> Beier, 1955*	Pa Baltic amber
CHELIFERIDAE Risso, 1826	Cretaceous – Recent
Cheliferidae? indet. <i>in</i> Judson (2009)	K Archingeay amber
† Dichela Menge, 1854	Palaeogene
= † <i>Oligochelifer</i> Beier, 1937	
28. <i>Dichela berendtii</i> Menge, 1954*	Pa Baltic amber
29. <i>Dichela gracilis</i> (Beier, 1937)	Pa Baltic amber
30. <i>Dichela granulatus</i> (Beier, 1937)	Pa Baltic amber
31. <i>Dichela serratidentatus</i> (Beier, 1937)	Pa Baltic amber
† Electrochelifer Beier, 1937	Palaeogene
32. <i>Electrochelifer bachofeni</i> Beier, 1947a	Pa Baltic amber
33. <i>Electrochelifer balticus</i> Beier, 1955	Pa Baltic amber
34. <i>Electrochelifer mengei</i> Beier, 1937*	Pa Baltic amber
35. <i>Electrochelifer rapulitarsatus</i> Beier, 1947a	Pa Baltic amber
† Heurtaultia Judson, 2009 [tentative referral to family]	Cretaceous
36. <i>Heurtaultia rossiorum</i> Judson, 2009	K Archingeay amber
† Pycnochelifer Beier, 1937	Palaeogene
37. <i>Pycnochelifer kleemanni</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
i. = <i>Obisium rathkii</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† Trachychelifer Hong, 1983b	Palaeogene
38. <i>Trachychelifer liaoningense</i> Hong, 1983b*	Pa Chinese amber
CHERNETIDAE Menge, 1855	Cretaceous – Recent
Chernetidae gen. et sp. indet. <i>in</i> Schawaller (1991)	K Canadian amber
Chernetidae gen. et sp. indet. <i>in</i> Schawaller (1982b)	Ne Chiapas amber
† Oligochernes Beier, 1937	Palaeogene
39. <i>Oligochernes bachofeni</i> Beier, 1937	Pa Baltic amber
40. <i>Oligochernes wigandi</i> (Menge, 1854)	Pa Baltic amber
Pachychernes Beier, 1932b	Neogene – Recent
41. <i>Pachychernes effossus</i> Schawaller, 1980b	Ne Dominican amber
42. <i>Pachychernes</i> aff. <i>subrobustus</i> (Balzan, 1892) [Recent]	Qt–R Colombian copal
WITHIIDAE Chamberlin, 1931b	Palaeogene – Recent
† Beierowithius Mahnert, 1979	Palaeogene
43. <i>Beierowithius sieboldtii</i> (Menge, 1854)*	Pa Baltic amber
Withius Kew, 1911	Quaternary – Recent
44. <i>Chelifer eucarpus</i> Dalman, 1826	Qt East African opal

NOMINA DUBIA

1. *Chelifer ehrenbergii* C. L. Koch & Berendt, 1854Pa Baltic amber

NOMINA NUDA

1. *Chelifer fossilis* Weyenbergh, 1874J Solnhofen

3,385 Recent species according to Harvey (2009)

SOLIFUGAE

5 currently valid species of camel spider

- *Schneidarachne* appears to show some solifuge-like features and was tentatively assigned to the stem-lineage of this order; for convenience it is listed here alongside the camel spiders
- a family name Protosolpugidae has been proposed for *Protosolpuga*, but was not recognised in most of the subsequent literature – cf. Selden & Shear's (1996) revision

stem-lineage?

- † ***Schneidarachne* Dunlop & Rössler, 2003** **Carboniferous**
 1. *Schneidarachne saganii* Dunlop & Rössler, 2003* C Kamienna Góra

SOLIFUGAE Sundevall, 1833 **Carbon. – Recent**

- † ***Protosolpuga* Petrunkevitch, 1913** **Carboniferous**
 2. *Protosolpuga carbonaria* Petrunkevitch, 1913* C Mazon Creek

AMMOTRECHIDAE Roewer, 1934 **Neogene – Recent**

- † ***Happlodontus* Poinar & Santiago-Blay, 1989** **Neogene**
 3. *Happlodontus proterus* Poinar & Santiago-Blay, 1989* Ne Dominican amber

CEROMIDAE Roewer, 1933 **Cretaceous – Recent**

- † ***Cratosolpuga* Selden in Selden & Shear, 1996** **Cretaceous**
 4. *Cratosolpuga wunderlichi* Selden in Selden & Shear, 1996* K Crato Formation

DAESIIDAE Kraepelin, 1899 **Palaeogene – Recent**

- † ***Palaeoblossia* Dunlop, Wunderlich & Poinar, 2004** **Palaeogene**
 5. *Palaeoblossia groehni* Dunlop, Wunderlich & Poinar, 2004* Pa Baltic amber

EREMOBATIDAE Kraepelin, 1901 **Recent**

no fossil record

GALEODIDAE Sundevall, 1833 **Recent**

no fossil record

GYLIPPIDAE Roewer, 1933 **Recent**

no fossil record

HEXISOPODIDAE Pocock, 1897 **Recent**

no fossil record

KARSCHIIDAE Kraepelin, 1899 **Recent**

no fossil record

MELANOBLOSSIDAE Roewer, 1933 **Recent**

no fossil record

MUMMUCIIDAE Roewer, 1934 **Recent**

no fossil record

RHAGODIDAE Pocock, 1897 **Recent**

no fossil record

SOLPUGIDAE Leach, 1815 **Recent**

no fossil record

1,075 Recent species according to Harvey (2003)

PALPIGRADI

1 currently valid species of fossil palpigrade

PALPIGRADI Thorell, 1888 **Neogene – Recent**

= MICROTHELYPHONIDA Grassi & Calandruccio, 1885

family uncertain

† *Paleokoenenia* Rowland & Sissom, 1980 **Neogene**

1. *Paleokoenenia mordax* Rowland & Sissom, 1980* Ne Onyx Marble

EUKOENENIIDAE Petrunkevitch, 1955a **Recent**

no fossil record

PROKOENENIIDAE Condé, 1996 **Recent**

no fossil record

MISIDENTIFICATIONS

1. *Sternarthron zitteli* Haase, 1890 [insect] J Solnhofen

2. *Sternarthron zitteli* var. *minor* (Oppenheim, 1887) [insect] J Solnhofen

78 Recent species according to Harvey (2003)

ACARI: PARASITIFORMES

15 currently valid species of fossil parasitiform mite

- higher systematics and sequence of taxa follows the third edition of *A Manual of Acarology* (Krantz & Walter, eds, 2009), except that their orders are listed here as suborders, and suborders as infraorders to achieve some degree of consistency with other arachnid higher taxa throughout this list

PARASITIFORMES Reuter, 1909	Cretaceous – Recent
= ANACTINOTRICHIDA author, date?	
OPILIOACARIDA Zachvatkin, 1952 (suborder)	Palaeogene – Recent
= NOTOSTIGMATA author, date?	
OPILIOACAROIDEA Vitzthum, 1931	Palaeogene – Recent
OPILIOACARIDAE Vitzthum, 1931	Palaeogene – Recent
= NEOACARIDAE Chamberlin & Mulaik, 1942	
<i>Opilioacarus</i> With, 1902	?Palaeogene – Recent
1. ? <i>Opilioacarus aenigmus</i> Dunlop, Sempf & Wunderlich, 2010	Pa Baltic amber
<i>Paracarus</i> Chamberlin & Mulaik, 1942	Palaeogene – Recent
2. <i>Paracarus pristinus</i> Dunlop, Wunderlich & Poinar, 2004	Pa Baltic amber
HOLOTHYRIDA Thorell, 1882 (suborder)	Recent
= TETRASTIGMATA author, date?	
HOLOTYHROIDEA Thorell, 1882	Recent
ALLOTHYRIDAE van der Hammen, 1972	Recent
no fossil record	
HOLOTHYRIDAE Thorell, 1882	Recent
no fossil record	
NEOTHYRIDAE Lehtinen, 1981	Recent
no fossil record	
IXODIDA Leach, 1815 (suborder)	Cretaceous – Recent
= METASTIGMATA author, date?	
IXODOIDEA Banks, 1907	Cretaceous – Recent
ARGASIDAE Murray, 1877	Cretaceous – Recent
<i>Carios</i> Latreille, 1796	Cretaceous – Recent
3. <i>Carios jerseyi</i> Klompen & Grimaldi, 2001	K New Jersey amber
<i>Ornithodoros</i> C. L. Koch, 1844	Neogene – Recent

4. <i>Ornithodoros antiquus</i> Poinar, 1995	Ne Dominican amber
IXODIDAE Banks, 1907	Cretaceous – Recent
<i>Amblyomma</i> C. L. Koch, 1844	Neogene – Recent
5. <i>Amblyomma</i> near <i>argentinae</i> Neumann, 1905 [Recent] (as <i>testudinis</i>) in Lane & Poinar (1986).....	Ne–R Dominican amber
6. <i>Amblyomma</i> near <i>dissimile</i> C. L. Koch, 1844 [Recent] in Kierens <i>et al.</i> (1986)	Ne–R Dominican amber
† <i>Compluriscutata</i> Poinar & Buckley, 2008	Cretaceous
7. <i>Compluriscutata vetulum</i> Poinar & Buckley, 2008*	K Myanmar amber
† <i>Cornupalpatum</i> Poinar & Brown, 2003	Cretaceous
8. <i>Cornupalpatum burmanicum</i> Poinar & Brown, 2003*	K Myanmar amber
<i>Dermacentor</i> C. L. Koch, 1844	Neogene – Recent
9. <i>Dermacentor</i> nr. <i>reticulatus</i> (Fabricius, 1794) [Recent] (in Kulczyński in Schille 1916).....	Ne–R in a Rhino's ear
<i>Hyalomma</i> C. L. Koch, 1844	Palaeogene – Recent
<i>Hyalomma</i> spp.	Pa Baltic amber
<i>Ixodes</i> Latreille, 1795	Palaeogene – Recent
10. <i>Ixodes sigelos</i> Keirans, Clifford & Corwin, 1976 [Recent]	Qt Argentina
11. <i>Ixodes succineus</i> Weidner, 1964	Pa Baltic amber
NUTALLIELLIDAE Schulze, 1935	Recent
no fossil record	
MESOSTIGMATA G. Canestrini, 1891 (suborder)	Palaeogene – Recent
= GAMASIDA Leach, 1815	
SEJIDA Kramer, 1885 (infraorder)	Palaeogene – Recent
= LIROASPINA author, date?	
= TRICHOPYGIDIINA author, date?	
SEJOIDEA Berlese, 1885	Palaeogene – Recent
ICHTHYOSTOMATOGASTERIDAE Sellnick, 1953	Recent
no fossil record	
SEJIDAE Berlese, 1885	Palaeogene – Recent
= LIROASPIDIDAE Trägårdh, 1946	
<i>Sejus</i> C. L. Koch, 1836 [NB: <i>Seius</i> in an invalid emendation].....	Palaeogene – Recent
12. <i>Sejus bdelloides</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
UROPODELLIDAE Camin, 1955	Recent
no fossil record	
TRIGYNASPIDA Camin & Gorirossi, 1955 (infraorder)	Recent

CERCOMEGISTINA Camin & Gorirossi, 1955 (cohort)	Recent
CERCOMEGISTOIDEA Trägårdh, 1937	Recent
ASTERNOSEIIDAE Vale, 1955	Recent
no fossil record	
CERCOMEGISTIDAE Trägårdh, 1937	Recent
no fossil record	
DAVACARIDAE Kethley, 1979	Recent
no fossil record	
PYROSEJIDAE Lindquist & Moraza, 1993	Recent
no fossil record	
SALTISEIIDAE Walter, 2000	Recent
no fossil record	
SEIODIDAE Kethley, 1979	Recent
no fossil record	
ANTENNOPHORINA Berlese, 1882 (cohort)	Recent
ANTENNOPHOROIDEA Berlese, 1892	Recent
ANTENNOPHORIDAE Berlese, 1892	Recent
no fossil record	
CELAENOPSOIDEA Berlese, 1892	Recent
CELAENOPSIDAE Berlese, 1892	Recent
no fossil record	
COSTACARIDAE Hunter, 1993	Recent
no fossil record	
DIPLOGYNIIDAE Trägårdh, 1941	Recent
no fossil record	
EUZERCONIDAE Trägårdh, 1938	Recent
no fossil record	
MEGACELAENOPSIDAE Funck, 1975	Recent
no fossil record	
MEINERTULIDAE Trägårdh, 1950	Recent
no fossil record	

NEOTENOGYNIIDAE Kethley, 1974	Recent
no fossil record	
SCHIZOGYNIIDAE Trägårdh, 1950	Recent
no fossil record	
TRIPLOGYNIIDAE Funck, 1977	Recent
no fossil record	
PARAMEGISTOIDEA Trägårdh, 1946	Recent
PARAMEGISTIDAE Trägårdh, 1946	Recent
no fossil record	
FEDRIZZIOIDEA Trägårdh, 1937	Recent
FEDRIZZIIDAE Trägårdh, 1937	Recent
no fossil record	
KLINCKOWSTROEMIIDAE Camin & Gorirossi, 1955	Recent
no fossil record	
PROMEGISTIDAE Kethley, 1979	Recent
no fossil record	
MEGISTHANOIDEA Berlese, 1914	Recent
HOPLOMEGISTIDAE Camin & Gorirossi, 1955	Recent
no fossil record	
MEGISTHANIDAE Berlese, 1914	Recent
no fossil record	
PARANTENNULOIDEA Willmann, 1940	Recent
PARANTENNULIDAE Willmann, 1940	Recent
no fossil record	
PHILODANIDAE Kethley, 1977b	Recent
no fossil record	
AENICTEQUOIDEA Kethley, 1979	Recent
AENICTEQUIDAE Kethley, 1979	Recent
no fossil record	
EUPHYSALOZERCONIDAE Kim, 2008	Recent

no fossil record

MESSORACARIDAE Kethley, 1977 **Recent**

no fossil record

PHYSALOZERCONIDAE Kethley, 1977 **Recent**

no fossil record

PTOCHACARIDAE Kethley, 1979 **Recent**

no fossil record

MONOGYNASPIDA Camin & Gorioffi, 1955 (infrorder) **Palaeogene – Recent**

MICROGYNIINA Trägårdh, 1942 (cohort) **Palaeogene – Recent**

MICROGYNOIDEA Trägårdh, 1942 **Palaeogene – Recent**

Microgynoidea sp. *in* Dunlop *et al.* (2013) Pa Baltic amber

MICROGYNIIDAE Trägårdh, 1942 **Recent**

 = **MICROSEJIDAE Trägårdh, 1942**

no fossil record

NOTHOGYNIDAE Walter & Kranz, 1999 **Recent**

no fossil record

HEATHERELLINA author, date? (cohort) **Recent**

HEATHERELLOIDEA Walter, 1997 **Recent**

HEATHERELLIDAE Walter, 1997 **Recent**

no fossil record

UROPODOIDEA Kramer, 1881 (cohort) **Palaeogene – Recent**

UROPODIAE Kramer, 1881 (subcohort) **Palaeogene – Recent**

PROTODINYCHOIDEA Evans, 1957 **Recent**

PROTODINYCHIDAE Evans, 1957 **Recent**

no fossil record

THINOZERCONOIDEA Halbert, 1915 **Recent**

THINOZERCONIDAE Halbert, 1915 **Recent**

no fossil record

POLYASPIDOIDEA Berlese, 1913 **Recent**

DITHINOZERCONIDAE Ainscough, 1979 **Recent**

no fossil record

POLYASPIDIDAE Berlese, 1913 **Recent**

no fossil record

TRACHYTIDAE Trägårdh, 1938 **Recent**

no fossil record

UROPODOIDEA Kramer, 1881 **Palaeogene – Recent**

BALOGHJKASZABIIDAE Hirschmann, 1979 **Recent**

no fossil record

BRASILUROPODIDAE Hirschmann, 1979 **Recent**

no fossil record

CILLIBIDAE Trägårdh, 1944 **Recent**

no fossil record

CLAUSIADINYCHIDAE Hirschmann, 1979 **Recent**

no fossil record

CIRCOCYLLIBAMIDAE Sellnick, 1926 **Recent**

no fossil record

CYLLIBULIDAE Hirschmann, 1979 **Recent**

no fossil record

DERAIOPHORIDAE Trägårdh, 1952 **Recent**

no fossil record

DINYCHIDAE Berlese, 1916 **Recent**

no fossil record

DISCOURELLIDAE Baker & Wharton, 1952 **Recent**

no fossil record

EUTRACHYTIDAE Trägårdh, 1944 **Recent**

no fossil record

HUTUFEIDERIIDAE Hirschmann, 1979 **Recent**

no fossil record

KASZABJBALOGHIIDAE Hirschmann, 1979 **Recent**

no fossil record

MACRODINYCHIDAE Hirschmann, 1979 **Recent**

no fossil record

METAGYNURIDAE Balogh, 1943	Recent
no fossil record	
NENTERIIDAE Hirschmann, 1979	Recent
no fossil record	
OPLITIDAE Johnston, 1968	Recent
no fossil record	
PHYMATODISCIDAE Hirschmann, 1979	Recent
no fossil record	
PRODINYCHIDAE Berlese, 1917	Recent
no fossil record	
ROTUNDABALOGHIIDAE Hirschmann, 1979	Recent
no fossil record	
TERASEJASPIDAE Hirschmann, 1979	Recent
no fossil record	
TREMATURIDAE Berlese, 1917	?Palaeogene – Recent
= TREMATURELLIDAE Trägårdh, 1944	
?Trematuridae <i>in</i> Lyubarsky & Perkovsky (2012)	Pa Rovno amber
Trichouropoda Berlese, 1916	?Palaeogene – Recent
?Trichouropoda sp. [as <i>Oodinychus</i> sp.] <i>in</i> Ramsay (1960)	Qt New Zealand
TRICHOCYLLIBIDAE Hirschmann, 1979	Recent
no fossil record	
TRICHOUROPODELLIDAE Hirschmann, 1979	Recent
no fossil record	
TRIGONUPODIDAE Hirschmann <i>in</i> Wisniewski, 1979	Recent
no fossil record	
UROACTINIIDAE Hirschmann & Zirngiebl-Nicol, 1964	Recent
no fossil record	
URODIASPIDIDAE Trägårdh, 1944	Recent
no fossil record	

URODINYCHIDAE Berlese, 1917	Palaeogene – Recent
<i>Uroobovella</i> Berlese, 1903	?Palaeogene – Recent
? <i>Uroobovella</i> sp. in Dunlop <i>et al.</i> (2013)	Pa Baltic amber
UROPODIDAE Kramer, 1881	Recent
no fossil record	
TRACHYUROPODOIDEA Berlese, 1917	Recent
TRACHYUROPODIDAE Berlese, 1917	Recent
no fossil record	
DIARTHROPHALLIAE Trägårdh, 1946 (subcohort)	Recent
DIARTHROPHALLOIDEA Trägårdh, 1946	Recent
DIARTHROPHALLIDAE Trägårdh, 1946	Recent
no fossil record	
HETEROZERCONINA author, date? (cohort)	Recent
HETEROZERCONOIDEA Berlese, 1892	Recent
DISCOZERCONIDAE Berlese, 1910	Recent
no fossil record	
HETEROZERCONIDAE Berlese, 1892	Recent
no fossil record	
GAMASINA Kramer, 1881 (cohort)	Palaeogene – Recent
Gamasina indet in Perkovsky <i>et al.</i> (2007)	Pa Rovno amber
EPICRIIAE Vitzthum, 1938 (subcohort)	Neogene – Recent
EPICRIOIDEA Berlese, 1885	Recent
EPICRIIDAE Berlese, 1885	Recent
no fossil record	
ZERCONOIDEA Berlese, 1892	Neogene – Recent
COPROZERCONIDAE Moraza & Lindquist, 1999	Recent
no fossil record	
ZERCONIDAE Berlese, 1892	Neogene – Recent
† <i>Paleozercon</i> Błaszak, Cokendolpher & Polyak, 1995	Neogene
13. <i>Paleozercon cavernicolus</i> Błaszak, Cokendolpher & Polyak, 1995	Ne New Mexico
ARCTACARIAE Johnston, 1982 (subcohort)	Recent
ARCTACAROIDEA Evans, 1955	Recent
ARCTACARIDAE Evans, 1955	Recent

no fossil record

PARASITIAE Reuter, 1909 (subcohort)	Palaeogene – Recent
PARASITOIDEA Oudemans, 1901	Palaeogene – Recent
PARASITIDAE Oudemans, 1901	Palaeogene – Recent
<i>Aclerogamasus</i> Athias, 1971	Palaeogene – Recent
14. <i>Aclerogamasus stenocornis</i> Witaliński, 2000	Pa Baltic amber

DERMANYSSIAE Evans & Till, 1997 (subcohort)	Neogene – Recent
VEIGAIIOIDEA Oudemans, 1939	Recent
VEIGAIIDAE Oudemans, 1939	Recent
= GAMASOLAEALAPTIDAE Oudemans, 1939	

no fossil record

RHODACAROIDEA Oudemans, 1902	Palaeogene – Recent
DIGAMASELLIDAE Evans, 1954 ...[or 57?]	Palaeogene – Recent
Digamasellidae sp. <i>in</i> Perkovsky <i>et al.</i> (2007).....	Pa Rovno amber
<i>Dendrolaelaps</i> Halbert, 1915	Neogene – Recent
15. <i>Dendrolaelaps fossilis</i> Hirschman, 1971	Ne Chiapas amber

EURYPARASITIDAE d’Antony, 1987	Recent
---	---------------

no fossil record

GAMASIPHIDAE author, date?	Recent
---	---------------

no fossil record

LAELAPTONYSSIDAE Womersley, 1956	Recent
---	---------------

no fossil record

OLOGAMASIDAE Ryke, 1962	Recent
--------------------------------------	---------------

no fossil record

PANTENIPHIDIDAE d’Antony, 1987	Recent
---	---------------

no fossil record

RHODACARIDAE Oudemans, 1902	Recent
--	---------------

no fossil record

TERANYSSIDAE Halliday, 2006	Recent
--	---------------

no fossil record

EVIPHIDOIDEA Berlese, 1913	Quaternary–Recent
---	--------------------------

EVIPHIDIDAE Berlese, 1913	Recent
no fossil record	
MACROCHELIDAE Vitzthum, 1930	Quaternary–Recent
Macrocheles Latreille, 1829	Quaternary–Recent
<i>Macrocheles</i> sp. in Ramsay (1960)	Qt New Zealand
MEGALOLAELAPIDAE author, date?	Recent
no fossil record	
PACHYLAELAPIDAE Berlese, 1913	Recent
= NEOPARASITIDAE Oudemans, 1939	
= BULBOGAMASIDAE Gu, Wang & Duan, 1991	
no fossil record	
PARHOLASPIDIDAE Evans, 1956	Recent
no fossil record	
ASCOIDEA Oudemans, 1905	Palaeogene – Recent
AMEROSEIIDAE Evans in Hughs, 1961	Recent
no fossil record	
ASCIDAE Voigts & Oudemans, 1905	?Palaeogene – Recent
?Ascidae sp. in Dunlop <i>et al.</i> (2013)	Pa Baltic amber
HALOLAELAPIDAE Karg, 1965	Recent
no fossil record	
MELICHARIDAE Hirschmann, 1962	Recent
no fossil record	
PODOCINIDAE Berlese, 1913	Quaternary – Recent
Podocinidae sp. in Aoki (1974)	Qt Mizunami copal
PHYTOSEIOIDEA Berlese, 1916	Recent
BLATTISCOIIDAE Garman, 1948	Recent
no fossil record	
OTOPHEIDOMENIDAE Treat, 1955	Recent
no fossil record	
PHYTOSEIIDAE Berlese, 1916	Recent
no fossil record	

DERMANYSSOIDEA Kolenati, 1859	Recent
DASYPONYSSIDAE Fonseca, 1940	Recent
no fossil record	
DERMANYSSIDAE Kolenati, 1859	Recent
no fossil record	
ENTONYSSIDAE Ewing, 1922	Recent
no fossil record	
HAEMOGAMASIDAE Oudemans, 1939	Recent
no fossil record	
HALARACHNIDAE Oudemans, 1906	Recent
no fossil record	
HIRSTIONYSSIDAE Evans & Till, 1966	Recent
no fossil record	
HYSTRICHONYSSIDAE Keegan, Yunker & Baker, 1960	Recent
no fossil record	
IPIHIOPSIDIDAE Kramer, 1886	Recent
no fossil record	
IXODORHYNCHIDAE Ewing, 1923	Recent
no fossil record	
LAELAPIDAE Berlese, 1892	Recent
no fossil record	
LARVAMIMIDAE Elzinga, 1993	Recent
no fossil record	
LEPTOLAELAPIDAE Karg, 1978	Recent
no fossil record	
MACRONYSSIDAE Oudemans, 1936	Recent
no fossil record	
MANITHERIONYSSIDAE Radovsky & Yunker, 1971	Recent
no fossil record	

OMENTOLAEALAPTIDAE Fain, 1961	Recent
no fossil record	
PNEUMOPHIONYSSIDAE Fonseca, 1940	Recent
no fossil record	
RAILLIETIIDAE Vitzthum, 1942	Recent
no fossil record	
RHINONYSSIDAE Trouessart, 1895	Recent
no fossil record	
SPELAEORHYNCHIDAE Oudemans, 1902	Recent
no fossil record	
SPINTURNICIDAE Oudemans, 1902	Recent
no fossil record	
TRICHOASPIDIDAE Gu, Wang & Li, 1991	Recent
no fossil record	
VARROIDAE Delfinado & Baker, 1974	Recent
no fossil record	

nomum dubium

1. *Ixodes tertiarius* Scudder, 1885 Pa Wyoming

c. 12,500 Recent species

ACARIFORMES

294 currently valid species of fossil acariform mite

- higher systematics and sequence of taxa follows the third edition of *A Manual of Acarology* (Krantz & Walter, eds, 2009), except that their orders are listed here as suborders, and suborders as infraorders to achieve some degree of consistency with other arachnid higher taxa throughout this list
- a putative Ordovician mite assigned to the derived Brachypylina group of the oribatids remains controversial and is not formally listed below

ACARIFORMES Zachvatkin, 1952 Devonian – Recent

= ACTINOTRICHIDA author, date?

TROMBIDIFORMES Reuter, 1909 (suborder) Devonian – Recent

SPHAEROLICHIDA OConnor, 1984 (infraorder) Recent

LORDALYCOIDEA Grandjean, 1939 Recent

LORDALYCHIDAE Grandjean, 1939 Recent

= HYBALICIDAE Theron, 1974

no fossil record

SPHAEROLICHOIDEA Berlese, 1913 Recent

SPHAEROLICHIDAE Berlese, 1913 Recent

no fossil record

PROSTIGMATA Kramer, 1877 (infraorder) Devonian – Recent

LABIDOSTOMMATIDES Lindquist, Krantz & Walter, 2009 (s.cohort) .. Palaeogene – Recent

LABIDOSTOMMATOIDEA Oudemans, 1906 Palaeogene – Recent

LABIDOSTOMMATIDAE Oudemans, 1906 Palaeogene – Recent

= NICOLETIELLIDAE Canestrini, 1891

Labidostomatidae sp. *in* Sidorchuk & Bertrand (2013) Pa Rovno amber

Labidostomatidae sp. *in* Sidorchuk & Bertrand (2013) Pa Bitterfeld amber

***Labidostomma* Kramer, 1879** Palaeogene – Recent

1. *Labidostomma (Nicoletiella) paleoluteum* Dunlop & Bertrand, 2011 Pa Baltic amber

2. *Labidostomma (Pseudocornutella) electri* Sidorchuk & Bertrand, 2013 .. Pa Baltic amber

***Sellnickiella* Feider & Vasiliu, 1969** Palaeogene – Recent

3. *Sellnickiella balticae* Sidorchuk & Bertrand, 2013 Pa Baltic amber

EUPODIDES Krantz, 1978 (supercohort) Devonian – Recent

BDELLOIDEA Dugès, 1834 Cretaceous – Recent

BDELLIDAE Dugès, 1834 Cretaceous – Recent

Bdellidae sp. <i>in Aoki</i> (1974)	Qt Mizunami copal
<i>Bdella</i> Latreille, 1795	Cretaceous – Recent
4. <i>Bdella bicincta</i> Menge <i>in C. L. Koch & Berendt</i> , 1854	Pa Baltic amber
5. <i>Bdella bombycina</i> Menge <i>in C. L. Koch & Berendt</i> , 1854	Pa Baltic amber
6. <i>Bdella obconica</i> Menge <i>in C. L. Koch & Berendt</i> , 1854	Pa Baltic amber
7. <i>Bdella vetusta</i> Ewing, 1937	K Manitobian amber
<i>Bdellodes</i> Oudemans, 1937	Palaeogene – Recent
8. <i>Bdellodes lata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
CUNAXIDAE Thor, 1902	Recent
no fossil record	
HALACAROIDEA Murray, 1877	Recent
HALACARIDAE Murray, 1877	Recent
no fossil record	
PEZIDAE Harvey, 1990	Recent
no fossil record	
EUPODOIDEA C. L. Koch, 1842	Palaeogene – Recent
COCCEUPODIDAE Jesionowska, 2010	Recent
no fossil record	
DENDOCHAETIDAE Oliver, 2008	Recent
no fossil record	
EUPODIDAE C. L. Koch, 1842	Recent
no fossil record	
ERIORHYNCHIDAE Qin & Halliday, 1997	Recent
no fossil record	
PENTAPALPIDAE Oliver & Theron, 2000	Recent
no fossil record	
PENTHALEIDAE Oudemans, 1931	Recent
no fossil record	
PENTHALODIDAE Thor, 1933	Palaeogene – Recent
<i>Penthalodes</i> Murray, 1877	Palaeogene – Recent
9. <i>Penthalodes tristiculus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber

PROTERORHAGIIDAE Lindquist & Palacios-Vargas, 1991	Recent
no fossil record	
RHAGIDIIDAE Oudemans, 1922	Paleogene – Recent
Rhagidiidae indet. <i>in</i> Judson & Wunderlich (2003)	Pa Baltic amber
<i>Poecilophysis</i> O. P.-Cambridge, 1876	Paleogene – Recent
? <i>Poecilophysis</i> sp. <i>in</i> Judson & Wunderlich (2003)	Pa Baltic amber
† <i>Zachardia</i> Judson & Wunderlich, 2003	Paleogene
10. <i>Zachardia flexipes</i> Judson & Wunderlich, 2003	Pa Baltic amber
STRANDTMANNIIDAE Zacharda, 1979	Recent
no fossil record	
TYDEOIDEA Kramer, 1877	Devonian – Recent
EREYNETIDAE Oudemans, 1931	Recent
= MICROEREUNETIDAE Bottazzi, 1950	
no fossil record	
IOLINIDAE Pritchard, 1956	Recent
no fossil record	
TRIOPHTYDEIDAE Andrè, 1980	Recent
= MEYERELLIDAE André, 1979	
no fossil record	
TYDEIDAE Kramer, 1877	Devonian – Recent
† <i>Palaeotydeus</i> Dubinin, 1962	Devonian – Recent
11. <i>Palaeotydeus devonicus</i> Dubinin, 1962	D Rhynie chert
† <i>Parapotacarus</i> Dubinin, 1962	Devonian – Recent
12. <i>Paraprotacarus hirsti</i> Dubinin, 1962	D Rhynie chert
ERIOPHYOIDEA Nalepa, 1898	Triassic – Recent
= TETRAPODILI author, date?	
† <i>Ampezzo</i> Linquist & Grimaldi <i>in</i> Schmidt <i>et al.</i>, 1012,	Triassic – Recent
13. <i>Ampezzo triassica</i> Lindquist & Grimaldi <i>in</i> Schmidt <i>et al.</i> , 2012*	Tr Italian amber
† <i>Triasacarus</i> Linquist & Grimaldi <i>in</i> Schmidt <i>et al.</i>, 1012,	Triassic – Recent
14. <i>Triasacarus fedelei</i> Lindquist & Grimaldi <i>in</i> Schmidt <i>et al.</i> , 2012*	Tr Italian amber
DIPTILOMIOPIDAE Keifer, 1944	Recent
no fossil record	
ERIOPHYIDAE Nalepa, 1898	?Palaeogene – Recent

- Aculops* Keifer, 1966** ? Palaeogene – Recent
 15. *Aculops keiferi* Southcott & Lange, 1971 ?Pa Australia
- PHYTOPTIDAE Murray, 1877** Neogene – Recent
 = NALEPELLIDAE Roivainen, 1953
 no fossil record
- ANYSTIDES van der Hammen, 1972 (supercohort)** Cretaceous – Recent
ANYSTINA van der Hammen, 1972 (cohort) Cretaceous – Recent
CAECULOIDEA Berlese, 1883 Paleogene – Recent
CAECULIDAE Berlese, 1883 Paleogene – Recent
***Procaeculus* Jacot, 1936** Paleogene – Recent
 16. *Procaeculus dominicensis* Coineau & Poinar, 2001 Ne Dominican amber
 17. *Procaeculus eridosae* Coineau & Magowski, 1994 Pa Baltic amber
- ADAMYSTOIDEA Cunliffe, 1957** Recent
ADAMYSTIDAE Cunliffe, 1957 Recent
 = SAXIDROMIDAE Coineau, 1974
 no fossil record
- ANYSTOIDEA Oudemans, 1902** Cretaceous – Recent
ANYSTIDAE Oudemans, 1902 Cretaceous – Recent
Anystidae sp. *in* Aoki (1974) Qt Mizunami copal
***Anystis* von Heyden, 1826** Cretaceous – Recent
 18. *Anystis malleator* (Menge *in* C. L. Koch & Berendt, 1854) Pa Baltic amber
 19. *Anystis subnuda* (Menge *in* C. L. Koch & Berendt, 1854) Pa Baltic amber
 20. *Anystis venustula* (C. L. Koch & Berendt, 1854) Pa Baltic amber
 † ***Mesoanystis* Zacharda, 1985** Cretaceous
 21. *Mesoanystis taimirensis* Zacharda, 1985* K Siberian amber
 † ***Palaeoerythracarus* Zacharda, 1985** Palaeogene
 22. *Palaeoerythracarus sachalinensis* Zacharda, 1985* Pa Sachalin amber
- PSEUDOCHEYLIDAE Oudemans, 1909** Recent
 = STIGMOCHEYLIDAE Kethley, 1990
 no fossil record
- TENERIFFIIDAE Thor, 1911b** Paleogene – Recent
Teneriffiidae sp. *indet in* Sayre *et al.* (1992) Pa Baltic amber
- PARATYDEOIDEA Baker, 1949** Recent
PARATYDEIDAE Baker, 1949 Recent
 no fossil record

STIGMOCHEYLIDAE Kethley, 1990	Recent
no fossil record	
POMERANTZIOIDEA Baker, 1949	Recent
POMERANTZIIDAE Baker, 1949	Recent
no fossil record	
PARASITENGONA Oudemans, 1909 (cohort)	Cretaceous – Recent
ERYTHRAIAE author, date? (subcohort)	Cretaceous – Recent
CALYPTOSTOMATOIDEA Oudemans, 1923	Recent
CALYPTOSTOMATIDAE Oudemans, 1923	Recent
no fossil record	
ERYTHRAEOIDEA Grandjean, 1947a	Cretaceous – Recent
larval Erythraeoidea <i>in</i> Zacharda & Krivoluckij (1985)	K Siberian amber
† Pararainbowia Dunlop, 2007	Cretaceous
23. <i>Pararainbowia martilli</i> Dunlop, 2007*	K Crato Formation
ERYTHRAEIDAE Robineau-Desvoidy, 1828	Paleogene – Recent
= LEPTIDAE Billberg, 1820	
= BALUSTIIDAE Grandjean, 1947	
Erythraeidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
† Arytaena Menge, 1854 in C. L. Koch & Berendt, 1854	Paleogene
24. <i>Arytaena troguloides</i> Menge <i>in</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
Balaustium von Heyden, 1826	Paleogene – Recent
25. <i>Balaustium illustris</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
Erythraeus Latrielle, 1806	Paleogene – Recent
26. <i>Erythraeus bifrons</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber
27. <i>Erythraeus foveolatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
28. <i>Erythraeus hirsutus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
29. <i>Erythraeus lagopus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
30. <i>Erythraeus longipes</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
31. <i>Erythraeus proavus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
32. <i>Erythraeus procerus</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber
33. <i>Erythraeus raripilus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
34. <i>Erythraeus rostratus</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber
35. <i>Erythraeus saccatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
Leptus Latrielle, 1796	Paleogene – Recent
36. <i>Leptus incertus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
† PROTERYTHRAEIDAE Vercammen-Grandjean, 1973	Cretaceous
† Proterythraeus Vercammen-Grandjean, 1973	Cretaceous

37. *Proterythraeus southcotti* Vercammen-Grandjean, 1973* K Manitoba amber
- SMARIDIDAE Vitzthum, 1929** **Paleogene – Recent**
 Smarididae *in* Kulicka (1990) Pa Baltic amber
- TROMBIDIAE author, date? (subcohort)** **Creteaceous – Recent**
trombidiid mites?
38. *Megameropsis aquensis* Gourret, 1887 Pa Aix-en-Provence
 39. *Pseudopachygnathus maculatus* Gourret, 1887 Pa Aix-en-Provence
- AMPHOTROMBIOIDEA Zhang, 1998** **Recent**
AMPHOTROMBIIDAE, Zhang, 1998 **Recent**
 no fossil record
- ALLOTANAUPODOIDAE Zhang & Fan, 2007** **Recent**
ALLOTANAUPODIDAE Zhang & Fan, 2007 **Recent**
 no fossil record
- TANAUPODOIDEA Thor, 1935** **Creteaceous – Recent**
TANAUPODIDAE Thor, 1935 **Creteaceous – Recent**
 = ?AMPHOTROMBIIDAE Zhang, 1998
 = TANAUPODASTRIDAE Feider, 1959
- † ***Atanaupodus* Judson & Mağol, 2009** **Cretaceous**
 40. *Atanaupodus bakeri* Judson & Mağol, 2009 K Archingeay amber
- CHYZERIOIDEA Womersley, 1954** **Recent**
CHYZERIIDAE Womersley, 1954 **Recent**
 no fossil record
- TROMBIDIOIDEA Leach, 1815** **Paleogene – Recent**
ACHAEMENOTHROMBIIDAE Saboori, Wohltmann & Hakimitabar, 2010 **Recent**
 no fossil record
- EUTROMBIDIIDAE Thor, 1935** **Recent**
 no fossil record
- MICROTROMBIDIIDAE Thor, 1935** **Recent**
 no fossil record
- NEOTHROMBIIDAE Feider, 1955** **Recent**
 no fossil record

- TROMBIDIIDAE Leach, 1815** **Paleogene – Recent**
= PARATHROMBIIDAE Feider, 1959
- Allothrombium Berlese, 1903** **Paleogene – Recent**
41. *Allothrombium clavipes* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- Trombidium Fabricius, 1775** **Paleogene – Recent**
42. *Trombidium crassipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
43. *Trombidium granulatum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
44. *Trombidium heterotrichum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
45. *Trombidium scrobiculatum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- NB: the next two families may be synonyms
- WALCHIIDAE Ewing, 1946** **Recent**
no fossil record
- TROMBICULOIDEA Ewing, 1929** **Recent**
- AUDYANIDAE Southcott, 1987** **Recent**
no fossil record
- JOHNSTONIANIDAE Thor, 1935** **Recent**
= NOTOTHROMBIIDAE Feider, 1959
no fossil record
- NEOTROMBIDIIDAE Feider, 1959** **Recent**
no fossil record
- LEEUWENHOEKIIDAE Womersley, 1944** **Recent**
no fossil record
- TROMBELLIDAE Leach, 1815** **Recent**
no fossil record
- TROMBICULIDAE Ewing, 1929** **Recent**
= VATACARIDAE Southcott, 1957
no fossil record
- YUREBILLOIDEA Southcott, 1966** **Recent**
- YUREBILLIDAE Southcott, 1996** **Recent**
no fossil record
- HYDRACARNIDIAE van der Hoeven, 1849 (subcohort)** **Neogene – Recent**
= HYDRACHNIDIA author, date?
= HYDRACHNELLAE author, date?

Undetermined water mites

Hygrobatoidea, Arrenuroidea or Lebertiodes *in Poinar* (1985) Ne Dominican amber

HYDRYPHANTOIDEA Piersig, 1896 **Recent**

CTENOTHYADIDAE Lundblad, 1936 **Recent**

no fossil record

EUPATRELLIDAE Viets, 1935 **Recent**

no fossil record

HYDRODROMIDAE Viets, 1936 **Recent**

= DIPLODONTIDAE Lundblad, 1927

no fossil record

HYDRYPHANTIDAE Piersig, 1896 **Recent**

= PROTZIIDAE Viets, 1926

no fossil record

MALGASACARIDAE Tuzovskij, Gerecke & Goldschmidt, 2007 **Recent**

no fossil record

RHYNCHOHYDRACARIDAE Lundblad, 1936 **Recent**

= CHATHROSPERCHONIDAE Lundblad, 1936

no fossil record

TERATOTHYADIDAE Viets, 1929 **Recent**

no fossil record

THERMACARIDAE Sokolow, 1927 **Recent**

no fossil record

ZELANDOTHYADIDAE Cook, 1983 **Recent**

no fossil record

EYLAOIDEA Leach, 1815 **Recent**

APHEVIDERULICIDAE Gerecke, Smith & Cook, 1999 **Recent**

no fossil record

EYLAIIDAE Leach, 1815 **Recent**

no fossil record

LIMNOCHARIDAE Grube, 1859 **Recent**

no fossil record

PIERSIGIIDAE Oudemans, 1902	Recent
no fossil record	
HYDROVOLZIOIDEA Thor, 1905	Recent
ACHERONTACARIDAE Cook, 1967	Recent
no fossil record	
HYDROVOLZIIDAE Thor, 1905	Recent
= POLYXOHALACARIDAE Motas, 1972	
no fossil record	
HYDRACHNOIDEA Leach, 1815	Recent
HYDRACHNIDAE Leach, 1815	Recent
no fossil record	
LEBERTOIDEA Thor, 1900	Recent
ACUCAPITIDAE Wiles, 1996	Recent
no fossil record	
ANISITSIELLIDAE Koenicke, 1910	Recent
= MAMERSOPSIDAE Viets, 1914	
no fossil record	
BANDAKIOPSISIDAE Panesar, 2004	Recent
no fossil record	
LEBERTIIDAE Thor, 1900	Recent
no fossil record	
NILOTONIIDAE Viets, 1929	Recent
no fossil record	
OXIDAE Viets, 1926	Recent
no fossil record	
RUTRIPALPIDAE Solokow, 1834	Recent
no fossil record	
SPERCHONTIDAE Thor, 1900	Recent
no fossil record	
STYGOTONIIDAE Cook, 1992	Recent

no fossil record

TEUTONIDAE Koenike, 1910 **Recent**

no fossil record

TORRENTICOLIDAE Piersig, 1902 **Recent**

= ATRACTIDEIDAE Thor, 1902

no fossil record

HYGROBATOIDEA C. L. Koch, 1842 **Recent**

ASTACOCROTONIDAE Thor, 1927 **Recent**

no fossil record

ATURIDAE Thor, 1900 **Recent**

= BRADYPODIDAE Thor, 1900 [preoccupied]

= AXONOPSIDAE Viets, 1929

= LJANIIDAE Thor, 1929

no fossil record

FELTRIIDAE Viets, 1926 **Recent**

no fossil record

FERRADASIIDAE Cook, 1980 **Recent**

no fossil record

FRONTIPODOPSIDAE Viets, 1931 **Recent**

no fossil record

HYGROBATIDAE C. L. Koch, 1842b **Recent**

no fossil record

LETHAXONIDAE Cook, Smith & Harvey, 2000 **Recent**

no fossil record

LIMNESIIDAE Thor, 1900 **Recent**

= NEOTORRENTICOLIDAE Lundblad, 1936

= EPALLAGOPODIDAE Viets, 1953

no fossil record

OMARTACARIDAE Cook, 1963 **Recent**

no fossil record

PIONIDAE Thor, 1900 **Recent**

= CURVIPEDIDAE Thor, 1900

- = ACERCIDAE Thor, 1909
- = FORELIIDAE Thor, 1923
- = NAUTARACHNIDAE Walter, 1925
- = HYDROCHOREUTIDAE Viets, 1942

no fossil record

PONTARACHNIDAE Koenicke, 1910 **Recent**

no fossil record

UNIONICOLIDAE Oudemans, 1909 **Recent**

- = ATRACIDAE Thor, 1900
- = NEUMANIIDAE Thor, 1923

no fossil record

WETTINIDAE Cook, 1956 **Recent**

no fossil record

ARRENUROIDEA Thor, 1900 **Neogene – Recent**

Family uncertain

† *Protoarrenurus* Cook in Palmer, 1957 **Neogene – Recent**

46. *Protoarrenurus convergens* Cook in Palmer, 1957* Ne Mojave Desert

ACALYPTONOTIDAE Walter, 1911 **Recent**

no fossil record

AMOENACARIDAE Smith & Cook, 1997 **Recent**

no fossil record

ARENOHYDRACARIDAE Cook, 1974 **Recent**

no fossil record

ARRENURIDAE Thor, 1900 **Recent**

no fossil record

ATHIENEMANNIIDAE Viets, 1922 **Recent**

- = CHELOMIDEOPSIDAE Lundblad, 1962

no fossil record

BOGATIIDAE Motas & Tanasachi, 1938 **Recent**

no fossil record

CHAPPUISIDIDAE Motas & Tanasachi, 1946 **Recent**

no fossil record

GRETACARIDAE Viets, 1978	Recent
no fossil record	
HARPAGOPALPIDAE Viets, 1924	Recent
no fossil record	
HUNGAROHYDRACACARIDAE Motas & Tanasachi, 1959	Recent
no fossil record	
KANTACARIDAE Imamura, 1959	Recent
no fossil record	
KRENDOWSKIIDAE Viets, 1926	Recent
no fossil record	
LAVERSIIDAE Cook, 1955	Recent
no fossil record	
MIDEIDAE Thor, 1911a	Recent
no fossil record	
MIDEOPSIDAE Koenicke, 1910	Recent
no fossil record	
MOMONIIDAE Viets, 1926	Recent
= STYGOMOMONIDAE Szalay, 1943	
no fossil record	
NEOACARIDAE Motas & Tanasachi, 1947	Recent
no fossil record	
NIPPONACARIDAE Imamura, 1959	Recent
no fossil record	
NUDOMIDEOPSIDAE Smith, 1990	Recent
no fossil record	
UCHIDASTYGACARIDAE Imamura, 1956	Recent
no fossil record	
STYGOTHROMBIAE Thor, 1935 (subcohort)	Recent
STYGOTHROMBOIDEA Thor, 1935	Recent
STYGOTHROMBIIDAE Thor, 1935	Recent

ELEUTHERENGONIDES Oudemans, 1909 (supercohort)	Cretaceous – Recent
RAPHIGNATHINA Kethley, 1982 (cohort)	Cretaceous – Recent
MYOBIOIDEA Mégnin, 1877	Recent
MYOBIIDAE Mégnin, 1877	Recent
no fossil record	
PTERYGOSOMATOIDEA Oudemans, 1910	Recent
PTERYGOSOMATIDAE Oudemans, 1910	Recent
no fossil record	
RAPHIGNATHOIDEA Kramer, 1877	Paleogene – Recent
BARBUTIIDAE Robaux, 1975	Recent
no fossil record	
CALIGONELLIDAE Grandjean, 1944	Recent
no fossil record	
CAMEROBIIDAE Southcott, 1957	Paleogene – Recent
<i>Neophyllobius</i> Berlese, 1886	Paleogene – Recent
47. <i>Neophyllobius succineus</i> Bolland & Magowski, 1990.....	Pa Baltic amber
CRYPTOGNATHIDAE Oudemans, 1902	Paleogene – Recent
no fossil record	
DASYTHYREIDAE Walter & Gerson, 1998	Recent
no fossil record	
EUPALOPSELLIDAE Willmann, 1952	Recent
no fossil record	
HOMOCALIGIDAE Wood, 1969	Recent
no fossil record	
MECOGNATHIDAE Gerson & Walter, 1998	Recent
no fossil record	
RAPHIGNATHIDAE Kramer, 1877	Recent
no fossil record	
STIGMAEIDAE Oudemans, 1931	Paleogene – Recent
<i>Mediolata</i> Canestrini, 1890	Paleogene – Recent
48. <i>Mediolata eocenica</i> Kuznetsov, Khaustov & Perkovsky, 2010.....	Pa Rovno amber

- XENOCALIGONELLIDAE Gonzalez, 1978** **Recent**
no fossil record
- TETRANYCHOIDEA Donnadieu, 1876** **Palaeogene – Recent**
ALLOCHAETOPHORIDAE Reck, 1959 **Recent**
no fossil record
- LINOTETRANIDAE Baker & Pritchard, 1953** **Recent**
no fossil record
- TENUIPALPIDAE Berlese, 1913** **Recent**
no fossil record
- TETRANYCHIDAE Donnadieu, 1876** **Palaeogene – Recent**
= BRYOBIIDAE Berlese, date?
***Metatetranychus* Oudemans, 1931** **Palaeogene – Recent**
49. *Metatetranychus gibbus* (C. L. Koch & Berendt, 1854) Pa Baltic amber
***Schizotetranychus* Trägårdh, 1915** **Palaeogene – Recent**
50. *Schizotetranychus brevipes* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- TUCKERELLIDAE Baker & Pritchard, 1953** **Recent**
no fossil record
- CHEYLETOIDEA Leach, 1815** **Cretaceous – Recent**
CHEyleTIDAE Leach, 1815 **Cretaceous – Recent**
***Cheyletus* Latreille, 1796** **Cretaceous – Recent**
51. *Cheyletus burmiticus* Cockerell, 1917b K Myanmar amber
52. *Cheyletus portentosus* C. L. Koch & Berendt, 1854 Pa Baltic amber
- DEMODECIDAE Nicolet, 1855** **Recent**
no fossil record
- HARPIRHYNCHIDAE Dubinin, 1957** **Recent**
no fossil record
- OPHIOPTIDAE Southcott, 1956** **Recent**
no fossil record
- PSORERGATIDAE Dubinin in Bregatova et al., 1955** **Recent**
no fossil record

- SYRINGOPHILIDAE** Laviopierre, 1953 **Recent**
no fossil record
- HETEROSTIGMATINA** Berlese, 1899 (cohort) **Cretaceous – Recent**
- TARSOCHYLOIDEA** Atyeo & Baker, 1964 **Recent**
- TARSOCHYLIDAE** Atyeo & Baker, 1964 **Recent**
no fossil record
- HETEROCHEYLOIDEA** Trägårdh, 1950 **Recent**
- HETEROCHEYLIDAE** Trägårdh, 1950 **Recent**
no fossil record
- DOLICHOCYBOIDEA** Mahunka, 1970 **Recent**
- CROTALOMORPHIDAE** Lindquist & Kranz, 2002 **Recent**
no fossil record
- DOLICHOCYBIDAE** Mahunka, 1970 **Recent**
no fossil record
- TROCHOMETRIDIOIDEA** Mahunka, 1970 **Recent**
- ATHYREACARIDAE** Lindquist Kaliszewski & Rack, 1990 **Recent**
= BEMBIDIACARIDAE Khuastov, 2000
no fossil record
- TROCHOMETRIDIIDAE** Mahunka, 1970 **Recent**
no fossil record
- SCUTACAROIDEA** Oudemans, 1916 **Recent**
- MICRODISPIDAE** Cross, 1965 **Recent**
no fossil record
- SCUTACARIDAE** Oudemans, 1916 **Recent**
no fossil record
- PYGMEPHOROIDEA** Cross, 1965 **Palaeogene – Recent**
Pygmephoroida sp. *in* Magowski (1995) Pa Baltic amber
- NEOPYGMEPHORIDAE** Cross, 1965 **Recent**
no fossil record
- PYGMEPHORIDAE** Cross, 1965 **Recent**
no fossil record

SITEROPTIDAE Mahunka, 1970	Recent
no fossil record	
PYEMOTOIDEA Oudemans, 1937	Cretaceous – Recent
ACAROPHENACIDAE Cross, 1965	Cretaceous – Recent
† <i>Protophenax</i> Magowski, 1994	Cretaceous
53. <i>Protophenax kotejii</i> Magowski, 1994*	K Russian amber
CARABOACARIDAE Mahunka, 1970	Recent
no fossil record	
PYEMOTIDAE Oudemans, 1937	Recent
= TROCHOMETRIDAE Mahunka, 1970	
<i>Pyemotes</i> Amerling, 1862	Palaeogene – Recent
54. <i>Pyemotes primus</i> Khaustov & Perkovsky, 2010	Pa Rovno amber
RESINACARIDAE Mahunka, 1975	Cretaceous –Recent
<i>Protoresinacaris</i> Khaustov & Poinar, 2010	Cretaceous
55. <i>Protoresinacars brevipedis</i> Khaustov & Poinar, 2010*	K Myanmar amber
TARSONEMOIDEA Canestrini & Fanzago, 1877	Quaternary – Recent
PODAPOLIPIDAE Ewing, 1922	Recent
no fossil record	
TARSONEMIDAE Canestrini & Fanzago, 1877	Quaternary – Recent
Tarsonemidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
Cohort <i>incertae sedis</i>	
CLOACAROIDEA Camin, Moss, Oliver & Singer, 1967	Recent
CLOACARIDAE Camin, Moss, Oliver & Singer, 1967	Recent
no fossil record	
EPIMYODICIDAE Fain, Lukoschus & Rosmalen, 1982	Recent
no fossil record	
SARCOPTIFORMES author, date? (suborder)	Devonian – Recent
ENDEOSTIGMATA author, date? (infraorder)	Devonian – Recent
= PACHYGNATHINA author, date?	
ALYCINA author, date? (cohort)	

ALYCOIDEA Canestrini & Fanzago, 1877	Devonian – Recent
ALYCIDAE Canestrini & Fanzago, 1877	Devonian – Recent
= PACHYGNATHIDAE Kramer, 1877	
= BIMICHAELIIDAE Womersley, 1944	
† Protacarus Hirst, 1923	Devonian
56. <i>Protacarus crani</i> Hirst, 1923*	D Rhyne chert
GRANDJEANICIDAE Kethley, 1977a	Recent
no fossil record	
MICROPSAMMIDAE Coineau & Theorn, 1983	Recent
no fossil record	
NANORCHESTIDAE Grandjean, 1937	Devonian – Recent
† Protospeleorchestes Dubinin, 1962	Devonian – Recent
57. <i>Protospeleorchestes pseudoprotacarus</i> Dubinin, 1962*	D Rhyne chert
NEMATALYCINA author, date? (cohort)	Recent
NEMATALYCOIDEA Strenke, 1954	Recent
NEMATALYCIDAE Strenke, 1954	Recent
no fossil record	
PROTONEMATALYCIDAE Kethley, 1989 [superfamily correct?]	Recent
no fossil record	
TERPNACARINA author, date? (cohort)	Recent
OEHSERCHESTOIDEA Kethley, 1977a	Recent
OEHSERCHESTIDAE Kethley, 1977a	Recent
no fossil record	
TERPNACAROIDEA Grandjean, 1939	Recent
TERPNACARIDAE Grandjean, 1939	Recent
no fossil record	
ALICORHAGIINA author, date? (cohort)	Devonian – Recent
ALICORHAGIOIDEA Grandjean, 1939	Devonian – Recent
ALICORHAGIIDAE Grandjean, 1939	Devonian – Recent
† Archaeacarus Kethley & Norton <i>in</i> Kethley <i>et al.</i> , 1989	Devonian
58. <i>Archaeacarus dubinini</i> Kethley & Norton <i>in</i> Kethley <i>et al.</i> , 1989*	D Gilboa
† Pseudoprotacarus Dubinin, 1962	Devonian
59. <i>Pseudoprotacarus scoticus</i> Dubinin, 1962*	D Rhyne chert

ORIBATIDA Dugès, 1834 (infraorder) Devonian – Recent

= CRYPTOSTIGMATA author, date?

NB: see remarks on the Ordovician fossil above

PALAEOSOMATA Grandjean, 1969 (supercohort) Devonian–Recent

family uncertain

† *Marcvippeda* Pérez-DA, 1988 Palaeogene

60. *Marcvippeda magallanes* Pérez-DA, 1988* [*Acari incerate sedis?*] Pa Patagonia, Chile

ACARONYCHOIDEA Grandjean, 1932 Recent

ACARONYCHIDAE Grandjean, 1932b Recent

no fossil record

ARCHAEONOTHRIDAE Grandjean, 1932 Recent

no fossil record

CTENACAROIDEA Grandjean, 1954c Devonian – Recent

ADELPHACARIDAE Grandjean, 1954c Carbon. – Recent

† *Monoaphelacarus* Subías & Arillo, 2002 Carboniferous

61. *Monoaphelacarus carboniferus* Subías & Arillo, 2002* C County Antrim

APHELACARIDAE Grandjean, 1954c Recent

no fossil record

CTENACARIDAE Grandjean, 1954b Devonian – Recent

† *Ctenacaronychus* Subías & Arillo, 2002 Devonian

62. *Ctenacaronychus nortoni* Subías & Arillo, 2002* D New York

† *Palaeoctenacarus* Subías & Arillo, 2002 Carboniferous

63. *Palaeoctenacarus simmsoi* Subías & Arillo, 2002* C County Antrim

PALAEACAROIDEA Grandjean, 1932b Recent

PALAEACARIDAE Grandjean, 1932b Recent

no fossil record

ENARTHRONOTA Grandjean, 1947b (supercohort) Devonian – Recent

superfamily uncertain

† **DEVONACARIDAE** Norton *in* Norton *et al.*, 1988 Devonian – Recent

† *Devonacarus* Norton *in* Norton *et al.*, 1988 Devonian – Recent

64. *Devonacarus sellnicki* Norton *in* Norton *et al.*, 1988* D Gilboa

† **PROTOCHTHONIIDAE** Norton *in* Norton *et al.*, 1988 Devonian – Recent

- † *Protochthonius* Norton in Norton et al., 1988 Devonian – Recent
65. *Protochthonius gilboa* Norton in Norton et al., 1988* D Gilboa
- BRACHYCHTHONIOIDEA Thor, 1934** Recent
BRACHYCHTHONIIDAE Thor, 1934 Recent
no fossil record
- ATOPOCHTHONIOIDEA Grandjean, 1948** Recent
ATOPOCHTHONIIDAE Grandjean, 1948 Recent
no fossil record
- PHYLLOCHTHONIIDAE Travé, 1967** Recent
no fossil record
- PTEROCHTHONIIDAE Grandjean, 1950** Recent
no fossil record
- HYPOCHTHONIOIDEA Berlese, 1910** Carbon. – Recent
ENIOCHTHONIIDAE Grandjean, 1947b Recent
no fossil record
- HYPOCHTHONIIDAE Berlese, 1910** Carbon. – Recent
Hypochthonius C. L. Koch, 1835 Quaternary – Recent
66. *Hypochthonius rufulus* C. L. Koch, 1835 [Recent] Qt Finland
- † *Palaeohypochthonius* Subías & Arillo, 2002 Carboniferous
67. *Palaeohypochthonius jerami* Subías & Arillo, 2002* C County Antrim
- LOHMANNIIDAE Berlese, 1916** Recent
= XENOLOHMANNIIDAE Balogh & Mahunka, 1969
no fossil record
- MESOPLOPHORIDAE Ewing, 1917** Recent
= ARCHOPLOPHORIDAE Grandjean, 1965
no fossil record
- PROTOPLOPHOROIDEA Ewing, 1917** Carbon. – Recent
COSMOCHTHONIIDAE Grandjean, 1947b Carbon. – Recent
† *Carbochthonius* Subías & Arillo, 2002 Carboniferous
68. *Carbochthonius antrimensis* Subías & Arillo, 2002* C County Antrim
- HAPLOCHTHONIIDAE van der Hammen, 1959** Recent
no fossil record

PEDICULOCHELIDAE Lavoipierre, 1946	Recent
no fossil record	
PROTHOPOPHORIDAE Ewing, 1917	Carbon. – Recent
= APOPOPHORIDAE Niedbala, 1984	
† <i>Archaeoplophora</i> Subías & Arillo, 2002	Carboniferous
69. <i>Archaeoplophora bella</i> Subías & Arillo, 2002*	C County Antrim
SPHAEROCHTHONIIDAE Grandjean, 1947 <i>b</i>	Recent
no fossil record	
HETEROCHTHONOIDEA Grandjean, 1954 <i>b</i>	Recent
ARBORICHTHONIIDAE Balogh & Balogh, 1992	Recent
no fossil record	
HETEROCHTHONIIDAE Grandjean, 1954 <i>b</i>	Recent
no fossil record	
TRICHTOCHTHONIIDAE Lee, 1982	Recent
no fossil record	
PARHYPOSOMATA Grandjean, 1969 (supercohort)	Carbon. – Recent
PARHYPOCHTHONIOIDEA Grandjean, 1932 <i>b</i>	Carbon. – Recent
ELLIPTOCHTHONIIDAE Norton, 1975	Recent
no fossil record	
GEHYPOCHTHONIIDAE Strenzke, 1963	Carbon. – Recent
† <i>Gehypochthonimimus</i> Subías & Arillo, 2002	Carboniferous
70. <i>Gehypochthonimimus hibernicus</i> Subías & Arillo, 2002*	C County Antrim
PARHYPOCHTHONIIDAE Grandjean, 1932 <i>b</i>	Recent
no fossil record	
MIXONOMATA Grandjean, 1969(supercohort)	Paleogene – Recent
NEHYPOCHTHONOIDEA Norton & Metz, 1980	Recent
NEHYPOCHTHONIIDAE Norton & Metz, 1980	Recent
no fossil record	
EULOHMANNOIDEA Grandjean, 1931	Recent
EULOHMANNIIDAE Grandjean, 1931	Recent
no fossil record	

PERLOHMANNIOIDEA Grandjean, 1954b	Recent
PERLOHMANNIIDAE Grandjean, 1954b	Recent
no fossil record	
EPILOHMANNIOIDEA Oudemans, 1923	Recent
EPILOHMANNIIDAE Oudemans, 1923	Recent
= LESSIRIIDAE Oudemans, 1916	
no fossil record	
COLLOHMANNIOIDEA Grandjean, 1958a	Paleogene – Recent
COLLOHMANNIIDAE Grandjean, 1958a	Paleogene – Recent
Collohmanna Sellnick, 1922	Paleogene – Recent
71. <i>Collohmanna schusteri</i> Norton, 2006	Pa Baltic amber
† Embolacarus Sellnick, 1919	Palaeogene – Recent
72. <i>Embolacarus pergratus</i> Sellnick, 1919*	Pa Baltic amber
EUPYCTIMA Grandjean, 1967	Palaeogene – Recent
NB: Eupyctima is listed here as a mixonomatid clade, but is not recognised in all classifications, or else is removed from this group and given equal rank	
EUPHTHIRACAROIDEA Jacot, 1930	Palaeogene – Recent
EUPHTHIRACARIDAE Jacot, 1930	Palaeogene – Recent
Microtritia Märkel, 1964	Quaternary – Recent
73. <i>Microtritia minima</i> (Berlese, 1904) [Recent]	Qt Germany
Rhysotritia Märkel & Meyer, 1959	Quaternary – Recent
74. <i>Rhysotritia ardua</i> (C. L. Koch, 1841) [Recent]	Qt Germany
75. <i>Rhysotritia duplicata</i> (Grandjean, 1953) [Recent]	Qt Germany
ORIBOTRITIIDAE Grandjean, 1954b	Palaeogene – Recent
= SABAHRITIIDAE Mahunka, 1987	
Oribotritia Jacot, 1924	Palaeogene – Recent
76. <i>Oribotritia pyropus</i> (Sellnick, 1919)	Pa Baltic amber
77. <i>Oribotritia translucida</i> Sellnick, 1931	Pa Baltic amber
SYNICHOTRITIIDAE Walker, 1965	Recent
no fossil record	
PHTHIRACAROIDEA Perty, 1841	Palaeogene – Recent
PHTHIRACARIDAE Perty, 1841	Palaeogene – Recent
= STEGANACARIDAE Niedbala, 1986	
Hoplophthiacarus Jacot, 1933	Quaternary – Recent
78. <i>Hoplophthiacarus pavidus</i> (Berlese, 1913) [Recent]	Qt Karelia, Russia

Phthiacarus Perty, 1841	Palaeogene – Recent
79. <i>Phthiacarus borealis</i> Trägårdh, date? [Recent]	Qt Karelia, Russia
80. <i>Phthiacarus multipunctus</i> (Sellnick, 1919)	Pa Baltic amber
Steganacarus Ewing, 1917	Quaternary – Recent
81. <i>Steganacarus applicatus</i> (Sellnick, 1920) [Recent]	Qt Denmark
82. <i>Steganacarus carinatus</i> (C. L. Koch, 1841) [Recent]	Qt Finland
83. <i>Steganacarus striculus</i> (C. L. Koch, 1835) [Recent]	Qt Europe
<i>Steganacarus</i> sp.	Qt Finland
DESMONOMATA Woodley, 1873 (supercohort)	Jurassic – Recent
NOTHRINA van der Hammen, 1982 (cohort)	Jurassic – Recent
= HOLOSOMATA author, date?	
CROTONIOIDEA Thorell, 1876	Jurassic – Recent
CAMISIIDAE Oudemans, 1900	Cretaceous – Recent
Camisia von Heyden, 1826	Paleogene – Recent
84. <i>Camisia foveolata</i> Hammer, 1955 [Recent]	Qt western Norway
85. <i>Camisia horrida</i> [Recent] <i>fossilis</i> Sellnick, 1919	Pa Baltic amber
i. = <i>Nothrus kuehli</i> Karsch, 1884	Pa Baltic amber
NB: unclear why the older name is the synonym	
86. <i>Camisia invenusta</i> (Michael, 1888) [Recent]	Qt western Norway
87. <i>Camisia lapponica</i> Trägårdh, 1910 [Recent]	Qt Karelia, Russia
† Eocamisia Bulanova-Zachvatkina, 1974	Cretaceous
88. <i>Eocamisia sukatshevae</i> Bulanova-Zachvatkina, 1974*	K Siberian amber
Platynothrus Berlese, 1913	Quaternary – Recent
89. <i>Platynothrus peltifer</i> (C. L. Koch, 1839) [Recent]	Qt Greenland
90. <i>Platynothrus punctatus</i> (L. Koch, 1879) [Recent]	Qt northern Europe
CROTONIIDAE Thorell, 1876	Neogene – Recent
= HOLONOTHRIDAE Wallwork, 1963	
Crotonia Thorell, 1876	Neogene – Recent
91. <i>Crotonia ramus</i> (Womersley, 1957)	Ne Australian retinite
HERMANNIIDAE Sellnick, 1928	Palaeogene – Recent
= GALAPAGACARIDAE P. Balogh, 1985	
Hermannia Nicolet, 1855	Palaeogene – Recent
92. <i>Hermannia gibba</i> (C. L. Koch, 1839) [Recent]	Qt Finland
93. <i>Hermannia reticulata</i> Thorell, 1871 [Recent]	Qt Subarctic – Arctic
94. <i>Hermannia scabra</i> (L. Koch, 1879) [Recent]	Qt Greenland
95. <i>Hermannia sellnicki</i> Norton, 2006	Pa Baltic amber
MALACONOTHRIDAE Berlese, 1916	Quaternary – Recent

<i>Malacoethrus</i> Berlese, 1904	Quaternary – Recent
96. <i>Malacoethrus monodactylus</i> (Michael, 1888) [Recent]	Qt Europe
<i>Trimalacoethrus</i> Berlese, 1916	Quaternary – Recent
97. <i>Trimalacoethrus maior</i> (Berlese, 1910) [Recent]	Qt northern Europe
NANHERMANNIIDAE Sellnick, 1928	Quaternary – Recent
<i>Nanhermannia</i> Berlese, 1913	Quaternary – Recent
98. <i>Nanhermannia coronata</i> Berlese, 1913 [Recent]	Qt Karelia, Russia
99. <i>Nanhermannia elegantula</i> Berlese, 1913 [Recent]	Qt Germany
NOTHRIDAE Berlese, 1896	Paleogene – Recent
<i>Nothrus</i> C. L. Koch, 1836	Paleogene – Recent
100. <i>Nothrus illautus</i> Sellnick, 1919	Pa Baltic amber
101. <i>Nothrus punctulum</i> Karsch, 1884	Pa Baltic amber
102. <i>Nothrus silvestris</i> Nicolet, 1855 [Recent]	Qt Europe
TRHYPOCHTHONIIDAE Willmann, 1931	Jurassic – Recent
= ALLONOTHRIDAE Lee, 1985	
= MUCRONOTHRIDAE Kunst, 1972	
= PARALLONOTHRIDAE Badejo, Woas & Beck, 2002	
= TRHYPOCHTHONIELLIDAE Knülle, 1957	
<i>Allonothrus</i> van der Hammen, 1953	Neogene – Recent
<i>Allonothrus</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
† <i>Juracarus</i> Krivolutsky in Krivolutsky & Krasilov, 1977	Jurassic – Recent
103. <i>Juracarus serratus</i> Krivolutsky in Krivolutsky & Krasilov, 1977	J Russian far east
<i>Mucronothrus</i> Trägårdh, 1931	Quaternary – Recent
104. <i>Mucronothrus nasalis</i> (Willmann, 1929) [Recent]	Qt Karelia, Russia
† <i>Palaeochthonius</i> Krivolutsky in Krivolutsky & Krasilov, 1977	Jurassic – Recent
105. <i>Palaeochthonius krasilovi</i> Krivolutsky in Kriv. & Krasilov, 1977	J Russian far east
<i>Trhypochthonius</i> Berlese, 1904	Palaeogene – Recent
106. <i>Trhypochthonius badiformis</i> Sellnick, 1931	Pa Baltic amber
107. <i>Trhypochthonius cladonicola</i> (Willmann, 1919) [Recent]	Qt Germany
108. <i>Trhypochthonius corniculatus</i> Sellnick, 1931	Pa Baltic amber
109. <i>Trhypochthonius tectorum</i> (Berlese, 1896) [Recent]	Qt Karelia, Russia
BRACHYPYLINA Hull, 1918 (cohort)	Jurassic – Recent
= CIRCUMDEHISCENTIAE Grandjean, 1954b	
= PORONOTA Grandjean, 1954b [in part; taxon used for seven brachypyline superfamilies]	
superfamily uncertain	
ARIBATIDAE Aoki, Takaku & Ito, 1994	Recent
no fossil record	

HERMANNIELLOIDEA Grandjean, 1934	Paleogene – Recent
HERMANNIELLIDAE Grandjean, 1934	Paleogene – Recent
<i>Hermanniella</i> Berlese, 1908	Paleogene – Recent
110. <i>Hermanniella concamerata</i> Sellnick, 1931	Pa Baltic amber
111. <i>Hermanniella tuberculata</i> Sellnick, 1919	Pa Baltic amber
<i>Sacculobates</i> Grandjean, 1962	Neogene – Recent
<i>Sacculobates</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
PLASMOBATIDAE Grandjean, 1961a	Recent
no fossil record	
NEOLIODOIDEA Sellnick, 1928	Palaeogene – Recent
= LIODOIDEA Grandjean, 1954b	
NEOLIODIDAE Sellnick, 1928	Palaeogene – Recent
= LIODIDAE Grandjean, 1954b	
<i>Neoliodes</i> Berlese, 1888	Palaeogene – Recent
= <i>Liodes</i> von Heyden, 1826 [preoccupied]	
112. <i>Neoliodes brevitarsus</i> (Woolley, 1971)	Ne Chiapas amber
113. <i>Neoliodes dominicus</i> Heethoff, Helfen & Norton, 2009	Ne Dominican amber
114. <i>Neoliodes quadriscutatus</i> Sellnick, 1919	Pa Baltic amber
<i>Neoliodes</i> sp. in Norton & Poinar (1993) [as <i>Liodes</i>]	Ne Dominican amber
<i>Platyliodes</i> Berlese, 1917	Palaeogene – Recent
115. <i>Platyliodes ensigerus</i> (Sellnick, 1919)	Pa Baltic amber
<i>Teleliodes</i> author, date?	Neogene – Recent
<i>Teleliodes</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
PLATEREMAEOIDEA Trägårdh, 1926	Cretaceous – Recent
= GYMNODAMAEOIDEA Grandjean, 1954a	
ALEURODAMAEIDAE Paschoal & Johnston, 1985	Recent
no fossil record	
GYMNODAMAEIDAE Grandjean, 1954a	Paleogene – Recent
<i>Gymnodamaeus</i> Kulczynski, 1902	Paleogene – Recent
116. <i>Gymnodamaeus sepotisus</i> Sellnick, 1919	Pa Baltic amber
IDIODAMAEIDAE Paschoal, 1987	Recent
no fossil record	
LICNOBELBIDAE Grandjean, 1965a	Recent
no fossil record	

LICNODAMAEIDAE Grandjean, 1954b	Recent
= NACUNANSELLIDAE author, date	
no fossil record	
LYRIFISSIELLIDAE Paschoal, 1987	Recent
no fossil record	
PEDROCORTESELLIDAE Paschoal, 1987	Recent
no fossil record	
PHEROLIODIDAE Paschoal, 1987	Recent
= HAMMERIELLIDAE Paschoal, 1987	
= NOOLIODIDAE Paschoal, 1987	
no fossil record	
PLATEREMAEIDAE Trägårdh, 1926	Cretaceous – Recent
<i>Rasnitsynella</i> Krivoluckij, 1976	Cretaceous
117. <i>Rasnitsynella punctulata</i> Krivoluckij, 1976	K Taymir amber
DAMAEOIDEA Berlese, 1896	Paleogene – Recent
DAMAEIDAE Berlese, 1896	Paleogene – Recent
Damaeidae sp. <i>in Aoki</i> (1974)	Qt Mizunami copal
<i>Belba</i> von Heyden, 1826	Quaternary – Recent
118. <i>Belba compta</i> (Kulczynski, 1902) [Recent]	Qt western Norway
119. <i>Belba cornyops</i> (Hermann, 1804)* [Recent]	Qt Finland
† <i>Belbites</i> Pampaloni, 1902	Neogene
120. <i>Belbites disodilis</i> Pampaloni, 1902*	Ne? Sicily
<i>Damaeobelba</i> Sellnick, 1928	Quaternary – Recent
121. <i>Damaeobelba minutissima</i> (Sellnick, 1920) [Recent]	Qt Germany
<i>Damaeus</i> C. L. Koch, 1835	Paleogene – Recent
122. <i>Damaeus auritus</i> C. L. Koch, 1835* [Recent]	Qt Finland
123. <i>Damaeus genadensis</i> Sellnick, 1931	Pa Baltic amber
<i>Spatiodamaeus</i> Bulanova-Zachvatkina, 1967	Quaternary – Recent
124. <i>Spatiodamaeus verticillipes</i> (Nicolet, 1855)* [Recent]	Qt Finland
CEPHEOIDEA Berlese, 1896	Cretaceous – Recent
= EUTEGOIDEA Balogh, 1965	
ANDEREMAEIDAE Balogh, 1972	Recent
no fossil record	
CEPHEIDAE Berlese, 1896	Cretaceous – Recent
= COMPATOZETIDAE Luxton, 1988	

Cepheus C. L. Koch, 1835	Paleogene – Recent
125. <i>Cepheus cepheiformis</i> (Nicolet, 1855) [Recent]	Qt Finland
126. <i>Cepheus dentatus</i> (Michael, 1888) [Recent]	Qt Finland
127. <i>Cepheus implicatus</i> (Sellnick, 1919)	Pa Baltic amber
128. <i>Cepheus latus</i> C. L. Koch, 1835* [Recent]	Qt Finland
Eupterotegaeus Berlese, 1916	Cretaceous – Recent
129. <i>Eupterotegaeus bitranslamellatus</i> Arillo & Subías, 2002	K Álava amber
Ommatocepheus Berlese, 1913	Cretaceous – Recent
130. <i>Ommatocepheus nortoni</i> Arillo, Subías & Shtanchaeva, 2008	K Álava amber
CEROCEPHEIDAE Mahunka, 1986	Recent
no fossil record	
EUTEGAEIDAE Balogh, 1965	Recent
= PTEROZETIDAE Luxton, 1988	
no fossil record	
MICROTEGEIDAE Balogh, 1972	Recent
no fossil record	
NODOCEPHEIDAE Piffi, 1972	Recent
no fossil record	
NOSYBEIDAE Mahunka, 1994	Recent
no fossil record	
PTEROBATIDAE Balogh & Balogh, 1992	Recent
no fossil record	
POLYPTEROZETOIDEA Grandjean, 1959	Recent
PODOPTEROTEGAEIDAE Piffi, 1972	Recent
no fossil record	
POLYPTEROZETIDAE Grandjean, 1959	Recent
no fossil record	
TUMEROZETIDAE Hammer, 1966	Recent
no fossil record	
MICROZETOIDEA Grandjean, 1936a	Recent
MICROZETIDAE Grandjean, 1936a	Recent

no fossil record

AMEROIDEA Bulanova-Zachvatkina, 1957 **Palaeogene – Recent**

= AMEROBELBOIDEA Grandjean, 1954*b*

= CALEREMEIOIDEA Grandjean, 1965*c*

AMERIDAE Bulanova-Zachvatkina, 1957 **Recent**

no fossil record

AMEROBELBIDAE Grandjean, 1961*b* **Recent**

no fossil record

BASILOBELBIDAE Balogh, 1961 **Recent**

no fossil record

CALEREMAEIDAE Grandjean, 1965*c* **Palaeogene – Recent**

***Caleremaeus* Berlese, 1910** **Palaeogene – Recent**

131. *Caleremaeus gleso* Sellnick, 1931 Pa Baltic amber

CTENOBELBIDAE Grandjean, 1965*b* **Recent**

no fossil record

DAMAEOLIDAE Grandjean, 1965*b* **Recent**

no fossil record

EREMOBELBIDAE Balogh, 1961 **Recent**

no fossil record

EREMULIDAE Grandjean, 1965*b* **Recent**

no fossil record

HETEROBELBIDAE Balogh, 1961 **Recent**

no fossil record

HUNGAROBELBIDAE Miko & Travé, 1996 **Recent**

no fossil record

STAUROBATIDAE Grandjean, 1966 **Recent**

no fossil record

ZETORCHESTOIDEA Michael, 1898 **Cretaceous – Recent**

= EREMAEOIDEA Oudemans, 1900

= NIPHOCEPHOIDEA Travé, 1959 [a separate superfamily in some studies]

† ARCHAEORCHESTIDAE Arillo & Subías, 2000	Cretaceous
† Platigeocranus Sellnick, 1919	Palaeogene
132. <i>Platigeocranus sulcatus</i> (Karsch, 1884)*	Pa Baltic amber
† Strieremaeus Sellnick, 1919	Cretaceous – Recent
= † <i>Archaeorchestes</i> Arillo & Subías, 2000	
133. <i>Strieremaeus illibatus</i> Sellnick, 1919	Pa Baltic amber
134. <i>Strieremaeus minguezae</i> (Arillo & Subías, 2000)	K Álava amber
EREMAEIDAE Oudemans, 1900	Paleogene – Recent
Eremaeus C. L. Koch, 1836	Paleogene – Recent
135. <i>Eremaeus hepaticus</i> C. L. Koch, 1835* [Recent]	Qt Germany
136. <i>Eremaeus oblongus</i> [Recent] <i>fossilis</i> Sellnick, 1919	Pa Baltic amber
Eueremaeus Mihelcic, 1963	Quaternary – Recent
137. <i>Eueremaeus silvestris</i> (Forsslund, 1956) [Recent]	Qt Finland
† Gradidorsum Sellnick, 1919	Palaeogene – Recent
138. <i>Gradidorsum asper</i> Sellnick, 1919*	Pa Baltic amber
MEGEREMAEIDAE Woolley & Higgins, 1968	Recent
no fossil record	
NIPHOCEPHEIDAE Travé, 1959	Recent
no fossil record	
ZETORCHESTIDAE Michael, 1898	Palaeogene – Recent
Zetorchestidae spp. <i>in</i> Sidorchuk & Norton (2011)	Pa Rovno amber
GUSTAVIOIDEA Oudemans, 1900	Jurassic – Recent
= LIACAROIDEA Sellnick, 1928	
ASTEGISTIDAE Balogh, 1961	Jurassic – Recent
Astegistes Hull, 1916	Quaternary – Recent
139. <i>Astegistes pilosus</i> (C. L. Koch, 1840) [Recent]	Qt Karelia, Russia
Cultroribula Berlese, 1908	Jurassic – Recent
140. <i>Cultroribula jurassica</i> Krivolutsky <i>in</i> Krivolutsky & Krasilov, 1977	J Russian far east
141. <i>Cultroribula lauta</i> Sellnick, 1931	Pa Baltic amber
142. <i>Cultroribula superba</i> Sellnick, 1931	Pa Baltic amber
GUSTAVIIDAE Oudemans, 1900	Quaternary – Recent
Gustavia Kramer, 1879	Quaternary – Recent
143. <i>Gustavia microcephala</i> (Nicolet, 1855) [Recent]	Qt Finland
KODIAKELLIDAE Hammer, 1967	Recent
no fossil record	

LIACARIDAE Sellnick, 1928	Quaternary – Recent
= XENILLIDAE Woolley & Higgins, 1966	
Adoristes Hull, 1916	Quaternary – Recent
144. <i>Adoristes ovatus</i> (C. L. Koch, 1839)* [Recent]	Qt northern Europe
Liacarus Michael, 1898	Quaternary – Recent
145. <i>Liacarus coracinus</i> (C. L. Koch, 1841) [Recent]	Qt Finland
Xenillus Robineau-Desvoidy, 1839	Paleogene – Recent
146. <i>Xenillus tegeocraniformis</i> (Sellnick, 1919)	Pa Baltic amber
MULTORIBULIDAE Balogh, 1972	Recent
no fossil record	
PELOPPIIDAE Balogh, 1943	Paleogene – Recent
Ceratoppia Berlese, 1908	Paleogene – Recent
147. <i>Ceratoppia bipilis fossilis</i> Sellnick, 1919	Pa Baltic amber
i. = <i>Oribates politus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
148. <i>Ceratoppia quadridentata</i> (Haller, 1882) [Recent]	Qt Finland
TENUIALIDAE Jacot, 1929	Quaternary – Recent
Hafenrefferia Oudemans, 1906	Quaternary – Recent
149. <i>Hafenrefferia gilvipes</i> (C. L. Koch, 1839)* [Recent]	Qt Finland
CARABODOIDEA C. L. Koch, 1843b	Palaeogene – Recent
= OCTOCEPHOIDEA Balogh, 1961	
CARABOCEPHEIDAE Mahunka, 1986	Recent
no fossil record	
CARABODIDAE C. L. Koch, 1843b	Palaeogene – Recent
Carabodes C. L. Koch, 1835	Palaeogene – Recent
150. <i>Carabodes areolatus</i> Berlese, 1916 [Recent]	Qt Karelia, Russia
151. <i>Carabodes coriaceus</i> C. L. Koch, 1835* [Recent]	Qt Finland
152. <i>Carabodes coriaceus</i> [Recent] <i>fossilis</i> Sellnick, 1931	Pa Baltic amber
153. <i>Carabodes dissonus</i> Sellnick, 1931	Pa Baltic amber
154. <i>Carabodes gerberi</i> Sellnick, 1931	Pa Baltic amber
155. <i>Carabodes laybrinthicus</i> (Michael, 1879) [Recent]	Qt Europe
156. <i>Carabodes labyrinthicus</i> [Recent] <i>fossilis</i> Sellnick, 1931	Pa Baltic amber
157. <i>Carabodes marginatus</i> (Michael, 1884) [Recent]	Qt Finland
158. <i>Carabodes minusculus</i> Berlese, 1923 [Recent]	Qt Germany
159. <i>Carabodes ornatus</i> Storkan, 1925 [Recent]	Qt Finland
160. <i>Carabodes subarcticus</i> Trägårdh, 1902 [Recent]	Qt Finland
161. <i>Carabodes willmanni</i> Bernini, 1975 [Recent]	Qt western Norway

? <i>Carabodes</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
† Carabodites Pampaloni, 1902	Neogene?
162. <i>Carabodites pavesii</i> Pampaloni, 1902*	Ne? Sicily
Odontocepheus Berlese, 1913	Quaternary – Recent
163. <i>Odontocepheus elongatus</i> (Michael, 1879)* [Recent]	Qt Finland
DAMPFIELLIDAE Balogh, 1961	Recent
no fossil record	
HEXOPPIIDAE Balogh, 1983	Recent
no fossil record	
LUXTONIIDAE Mahunka, 2001	Recent
no fossil record	
NIPPOBODIDAE Aoki, 1959	Recent
no fossil record	
OTOCEPHEIDAE Balogh, 1961	Paleogene – Recent
Dolicheremaeus Jacot, 1938	Neogene – Recent
<i>Dolicheremaeus</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
Otocepheus Berlese, 1905	Paleogene – Recent
164. <i>Otocepheus niger</i> Sellnick, 1931	Pa Baltic amber
165. <i>Otocepheus praesignis</i> Sellnick, 1931	Pa Baltic amber
TOKUNOCEPHEIDAE Aoki, 1966a	Recent
no fossil record	
OPPIOIDEA Grandjean, 1951	Palaeogene – Recent
= EREMELLOIDEA Balogh, 1961 [in part]	
= TRIZETOIDEA Ewing, 1917 [in part]	
AUTOGNETIDAE Grandjean, 1960b	Quaternary – Recent
Conchogneta Grandjean, 1963	Quaternary – Recent
166. <i>Conchogneta traegardhi</i> (Forsslund, 1947) [Recent]	Qt Finland
ARCEREMAEIDAE Balogh, 1972	Recent
no fossil record	
BORHIDIIDAE Balogh, 1983	Recent
no fossil record	
CHAVINIIDAE Balogh, 1983	Recent

no fossil record

ENANTIOPPIIDAE Balogh, 1983 **Recent**

no fossil record

EPIMERELLIDAE Ayyildiz & Luxton, 1989 **Recent**

no fossil record

GRANULOPPIIDAE Balogh, 1983 **Recent**

no fossil record

MACHADOBELBIDAE Balogh, 1972 **Recent**

no fossil record

MACHUELLIDAE Balogh, 1893 **Recent**

no fossil record

NOSYBELBIDAE Mahunka, 1994 **Recent**

no fossil record

OPPIIDAE Grandjean, 1951 **Palaeogene – Recent**

***Dissorhina* Hull, 1916** **Quaternary – Recent**

167. *Dissorhina ornata* (Oudemans, 1900)* **[Recent]** Qt Germany

***Oppia* C. L. Koch, 1836** **Palaeogene – Recent**

168. *Oppia angustum* (Sellnick, 1931) Pa Baltic amber

169. *Oppia cervicornu* (Sellnick, 1919) Pa Baltic amber

170. *Oppites hurdi* Woolley, 1971 Ne Chiapas amber

171. *Oppia longilamellata* **[Recent]** *fossilis* (Sellnick, 1931) Pa Baltic amber

172. *Oppia medium* (Sellnick, 1931) Pa Baltic amber

173. *Oppia mexicana* (Woolley, 1971) Ne Chiapas amber

174. *Oppia setigera* (Woolley, 1971) Ne Chiapas amber

175. *Oppia sucinum* (Sellnick, 1931) Pa Baltic amber

? *Oppia* sp. *in* Norton & Poinar (1993) Ne Dominican amber

***Oppiella* Jacot, 1937** **Quaternary – Recent**

176. *Oppiella nova* (Oudemans, 1902)* **[Recent]** Qt northern Europe

177. *Oppiella ornata* (Oudemans, 1900) **[Recent]** Qt western Norway

178. *Oppiella splendens* (C. L. Koch, 1841) **[Recent]** Qt western Norway

179. *Oppiella subpectinata* (Oudemans, 1900) **[Recent]** Qt northern Europe

180. *Oppiella translamellata* (Willmann, 1923) **[Recent]** Qt northern Europe

† ***Oppites* Pampaloni, 1902** **Neogene**

181. *Oppites melilli* Pampaloni, 1902* Ne? Sicily

***Ramusella* Hammer, 1962** **Quaternary – Recent**

182. *Ramusella clavipectinata* (Michael, 1885) **[Recent]** Qt Germany

OXYAMERIDAE Aoki, 1965	Recent
no fossil record	
PAPILLONOTIDAE Balogh, 1983	Recent
no fossil record	
PLATYAMERIDAE Balogh & Balogh, 1983	Recent
no fossil record	
QUADROPPIIDAE Balogh, 1983	Recent
no fossil record	
RHYNCHORIBATIDAE Balogh, 1961	Recent
no fossil record	
SPINOZETIDAE Balogh, 1972	Recent
no fossil record	
STERNOPPIIDAE Balogh & Mahunka, 1969	Recent
no fossil record	
SUCTOBELBIDAE Jacot, 1938	Palaeogene – Recent
<i>Suctobelbella</i> Jacot, 1937	Palaeogene – Recent
183. <i>Suctobelbella falcata</i> (Forsslund, 1941) [Recent]	Qt Germany
184. <i>Suctobelbella latirostris</i> (Strenzke, 1950) [Recent]	Qt Germany
185. <i>Suctobelbella longirostris</i> (Forsslund, 1941) [Recent]	Qt western Norway
186. <i>Suctobelbella sarekensis</i> (Forsslund, 1941) [Recent]	Qt Europe
187. <i>Suctobelbella similis</i> (Forsslund, 1941) [Recent]	Qt Germany
188. <i>Suctobelbella subcornigera</i> (Forsslund, 1941) [Recent]	Qt Germany
189. <i>Suctobelbella subtrigona</i> (Oudemans, 1916) [Recent]	Qt Europe
190. <i>Suctobelbella subtrigona</i> [Recent] <i>fossilis</i> (Sellnick, 1931)	Pa Baltic amber
TERATOPPIIDAE Balogh, 1983	Recent
no fossil record	
TETRACONDYLIDAE Aoki, 1961	Recent
no fossil record	
THYRISOMIDAE Grandjean, 1954b	Quaternary – Recent
<i>Banksinoma</i> Oudemans, 1930	Quaternary – Recent
191. <i>Banksinoma lanceolata</i> (Michael, 1885)* [Recent]	Qt Europe

TRIZETIDAE Ewing, 1917	Recent
no fossil record	
TUPAREZETIDAE Balogh, 1972	Recent
no fossil record	
TECTOCEPHEOIDEA Grandjean, 1954b	Paleogene – Recent
TECTOCEPHEIDAE Oudemans, 1900	Paleogene – Recent
<i>Tectocepheus</i> Berlese, 1895	Paleogene – Recent
192. <i>Tectocepheus minor</i> Berlese, 1903 [Recent]	Qt western Norway
193. <i>Tectocepheus similis</i> Sellnick, 1931	Pa Baltic amber
194. <i>Tectocepheus velatus</i> (Michael, 1880)* [Recent]	Qt northern Europe
HYDROZETOIDEA Grandjean, 1954b	Jurassic – Recent
HYDROZETIDAE Grandjean, 1954b	Jurassic – Recent
<i>Hydrozetes</i> Berlese, 1902	Jurassic – Recent
195. <i>Hydrozetes confervae</i> (Schrank, 1791) [Recent]	Qt western Norway
196. <i>Hydrozetes lacustris</i> (Michael, 1882)* [Recent]	Qt northern Europe
197. <i>Hydrozetes oryktosis</i> Woolley, 1969	Qt Michigan
<i>Hydrozetes</i> sp. in Sivhead & Wallwork (1978)	J Sweden
LIMNOZETIDAE Thor, 1937	Quaternary – Recent
<i>Limnozetes</i> Hull, 1916	Quaternary – Recent
198. <i>Limnozetes ciliatus</i> (Schrank, 1803)* [Recent]	Qt northern Europe
199. <i>Limnozetes rugosus</i> (Sellnick, 1923) [Recent]	Qt northern Europe
AMERONOTHROIDEA Willmann, 1931	Quaternary – Recent
AMERONOTHRIDAE Willmann, 1931	Quaternary – Recent
<i>Ameronothrus</i> Berlese, 1896	Quaternary – Recent
200. <i>Ameronothrus lineatus</i> (Thorell, 1871)* [Recent]	Qt Europe / Greenland
201. <i>Ameronothrus maculatus</i> (Michael, 1882) [Recent]	Qt western Norway
FORTUYNIIDAE van der Hammen, 1963	Recent
no fossil record	
SELENORIBATIDAE Schuster, 1963	Recent
no fossil record	
TEGEOCRANELLIDAE Balogh, 1987	Recent
no fossil record	
CYMBAEREMAEOIDEA Sellnick, 1928	Jurassic – Recent

CYBAEREMAEIDAE Sellnick, 1928	Jurassic – Recent
= AMETROPROCTIDAE Subías, 2004	
= SCAPHEREMAEIDAE Subías, 2004	
<i>Ametroproctus</i> Higgins & Woolley, 1968	Cretaceous – Recent
202. <i>Ametroproctus valeriae</i> Arillo, Subías & Shtanchaeva, 2009	K San Just amber
<i>Cymbaeremaeus</i> Berlese, 1896	Paleogene – Recent
203. <i>Cymbaeremaeus cymba</i> (Nicolet, 1855)* [Recent]	Qt northern Europe
† <i>Jureremus</i> Krivolutsky in Krivolutsky & Krasilov, 1977	Jurassic
204. <i>Jureremus foveolatus</i> Krivolutsky in Krivolutsky & Krasilov, 1977*	J Russian far east
205. <i>Jureremus phippsi</i> Selden, Baker & Phipps, 2008	J Yorkshire, UK
<i>Scapheremaeus</i> Berlese, 1910	Paleogene – Recent
206. <i>Scapheremaeus undosus</i> Sellnick, 1919	Pa Baltic amber
† <i>Tectocymba</i> Sellnick, 1919	Paleogene – Recent
207. <i>Tectocymba rara</i> Sellnick, 1919*	Pa Baltic amber
EREMAEOZETOIDEA Piffli, 1972	Paleogene – Recent
= IDIOZETOIDEA Aoki, 1976	
EREMAEOZETIDAE Piffli, 1972	Paleogene – Recent
<i>Eremaeozetes</i> Berlese, 1913	Paleogene – Recent
= † <i>Scutoribates</i> Sellnick, 1919	
<i>Eremaeozetes</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
IDIOZETIDAE Aoki, 1976	Recent
no fossil record	
LICNEREMAEOIDEA Grandjean, 1931	Palaeogene – Recent
= CHARASSOBATOIDEA Grandjean, 1958b	
ADHAESOZETIDAE Hammer, 1973	Recent
no fossil record	
CHARASSOBATIDAE Grandjean, 1958b	Recent
no fossil record	
DENDEROEREMAEIDAE Behan-Pelletier, Eamer & Clavton, 2005	Recent
no fossil record	
EREMELLIDAE Balogh, 1961	Recent
no fossil record	
LAMELLAREIDAE Balogh, 1972	Recent
no fossil record	

LICNEREMAEIDAE Grandjean, 1931	Palaeogene – Recent
<i>Licneremaeus</i> Paoli, 1908	Palaeogene – Recent
208. <i>Licneremaeus fritschi</i> Sellnick, 1931	Pa Baltic amber
209. <i>Licneremaeus licnophorus</i> (Michael, 1882) [Recent]	Qt Germany
MICREREMIDAE Grandjean, 1954b	Jurassic – Recent
<i>Micreremus</i> Grandjean, 1954b[not Berlese 1908?].....	Paleogene – Recent
210. <i>Micreremus brevipes</i> (Michael, 1888)* [Recent]	Qt northern Europe
211. <i>Micreremus reticulatus</i> Sellnick, 1931	Pa Baltic amber
212. <i>Micreremus scrobiculatus</i> Sellnick, 1931	Pa Baltic amber
PASSALOZETIDAE Grandjean, 1954b	Quaternary – Recent
<i>Passalozetes</i> Grandjean, 1932a	Quaternary – Recent
213. <i>Passalozetes africanus</i> Grandjean, 1932a [Recent]	Qt Finland
SCUTOVERTICIDAE Grandjean, 1954b	Neogene – Recent
<i>Arthrovertex</i> Balogh, 1970	Neogene – Recent
214. <i>Arthrovertex hurdi</i> (Woolley, 1971).....	Ne Chiapas amber
<i>Arthrovertex</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
<i>Scutovertex</i> Michael, 1879	Quaternary – Recent
215. <i>Scutovertex minutus</i> (C. L. Koch, 1835) [Recent]	Qt Germany
PHENOPELOPOIDEA Petrunkevitch, 1955a	Palaeogene – Recent
PHENOPELOPIDAE Petrunkevitch, 1955a	Palaeogene – Recent
= PELOPIDAE author, date?	
<i>Eupelops</i> Ewing, 1917	Palaeogene – Recent
216. <i>Eupelops acromios</i> (Hermann, 1804) [Recent]	Qt Finland
217. <i>Eupelops curtipilus</i> (Berlese, 1916) [Recent]	Qt Germany
218. <i>Eupelops occultus</i> (C. L. Koch, 1835) [Recent]	Qt Kerelia, Russia
219. <i>Eupelops plicatus</i> (C. L. Koch, 1835) [Recent]	Qt northern Europe
220. <i>Eupelops punctulatus</i> (Sellnick, 1931)	Pa Baltic amber
221. <i>Eupelops uraceus</i> (C. L. Koch, 1839)* [Recent]	Qt Kerelia, Russia
<i>Eupelops</i> sp. in Karppinen & Koponen (1974)	Qt Finland
<i>Peloptulus</i> Berlese, 1908	Quaternary – Recent
222. <i>Peloptulus phaenotus</i> (C. L. Koch, 1844)* [Recent]	Qt Germany
UNDULORIBATIDAE Kunst, 1971	Palaeogene – Recent
<i>Scutoribates</i> Sellnick, 1918	Palaeogene – Recent
223. <i>Scutoribates perornatus</i> Sellnick, 1918	Pa Baltic amber
<i>Unduloribates</i> Balogh, 1943	?Palaeogene – Recent
224. <i>Unduloribates parvus</i> (Sellnick, 1931)	Pa Baltic amber

[generic affinities need clarification]

ACHIPTERIOIDEA Thor, 1929	?Jurassic – Recent
ACHIPTERIIDAE Thor, 1929	?Jurassic – Recent
<i>Achipteria</i> Berlese, 1885	?Jurassic – Recent
225. <i>Achipteria coleoptera</i> (Linnaeus, 1757) [Recent]	Qt Finland / Greenland
226. ? <i>Achipteria obscura</i> Krivolutsky in Krivolutsky & Krasilov, 1977	J Russian far east
[An <i>incertae sedis</i> taxon?]	
<i>Parachipteria</i> van der Hammen, 1952	Quaternary – Recent
227. <i>Parachipteria punctata</i> (Nicolet, 1855) [Recent]	Qt northern Europe
228. <i>Parachipteria willmanni</i> van der Hammen, 1952 [Recent]	Qt Germany
EPACTOZETIDAE Grandjean, 1936b	Recent
no fossil record	
TEGORIBATIDAE Grandjean, 1954b	Quaternary – Recent
<i>Tegoribates</i> Ewing, 1917	Quaternary – Recent
229. <i>Tegoribates latirostris</i> (C. L. Koch, 1844) [Recent]	Qt Finland
ORIBATELLOIDEA Jacot, 1925	Palaeogene – Recent
ORIBATELLIDAE Jacot, 1925	Palaeogene – Recent
<i>Oribatella</i> Banks, 1895	Palaeogene – Recent
230. <i>Oribatella berleseii</i> (Michael, 1898) [Recent]	Qt Finland
231. <i>Oribatella calcarata</i> (C. L. Koch, 1835) [Recent]	Qt Kerelia, Russia
232. <i>Oribatella mirabilis</i> Sellnick, 1931	Pa Baltic amber
ORIPODOIDEA Jacot, 1925	Palaeogene – Recent
CALOPPIIDAE Balogh, 1960	Recent
= ?CRASSORIBATULIDAE author, date?	
no fossil record	
CAMPBELLOBATIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
CHAUNOPROCTIDAE Balogh, 1961	Recent
no fossil record	
DRYMOBATIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
HAPLOZETIDAE Grandjean, 1936c	Palaeogene – Recent
= PROTORIBATIDAE J. Balogh & P. Balogh, 1984	

= XLOBATIDAE J. Balogh & P. Balogh, 1984

Protoribates Berlese, 1908	Palaeogene – Recent
233. <i>Protoribates longipilis</i> Sellnick, 1931	Pa Baltic amber
LAMELLAREIDAE Balogh, 1972	Recent
no fossil record	
MAUDHEIMIIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
MOCHLOZETIDAE Grandjean, 1960a	Neogene – Recent
Mochlozetidae sp. <i>in</i> Norton & Poinar (1993)	Ne Dominican amber
Mochloribatula Mahunka, 1978	Neogene – Recent
234. <i>Mochloribatula smithi</i> (Woolley, 1971)	Ne Chiapas amber
Mochlozetes Grandjean, 1930	Neogene – Recent
<i>Mochlozetes</i> sp. <i>in</i> Norton & Poinar (1993)	Ne Dominican amber
NASOBATIDAE Balogh, 1972	Recent
no fossil record	
NEOTRICHOSZETIDAE Balogh, 1965	Recent
no fossil record	
NESOSZETIDAE J. Balogh & P. Balogh, 1984	Recent
no fossil record	
ORIBATULIDAE Thor, 1929	Palaeogene – Recent
Oribatulidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
Lucoppia Berlese, 1908	Palaeogene – Recent
235. <i>Lucoppia simplex</i> Sellnick, 1919	Pa Baltic amber
Oribatula Berlese, 1895	Quaternary – Recent
236. <i>Oribatula tibialis</i> (Nicolet, 1855)* [Recent]	Qt Europe
Phauloppia Berlese, 1908	Palaeogene – Recent
237. <i>Phauloppia lucorum</i> (C. L. Koch, 1841) [Recent]	Qt northern Europe
238. <i>Phauloppia pellucida</i> (Sellnick, 1931)	Pa Baltic amber
† Sachalinella Rjabinin <i>in</i> Krivolutzkii & Rjabinin, 1976	Palaeogene – Recent
May be a homonym of a bivalve genus	
239. <i>Sachalinella zherichini</i> Rjabinin <i>in</i> Krivolutzkii & Rjabinin, 1976*	Pa Sachalin amber
Zygoribatula Berlese, 1916	Quaternary – Recent
240. <i>Zygoribatula exilis</i> (Nicolet, 1855) [Recent]	Qt northern Europe
ORIPODIDAE Jacot, 1925	Palaeogene – Recent

= BIROBATIDAE J. Balogh & P. Balogh, 1984

Benoibates Balogh, 1958	Neogene – Recent
241. <i>Benoibates chiapasensis</i> (Woolley, 1971)	Ne Chiapas amber
Oripoda Banks, 1904	Palaeogene – Recent
242. <i>Oripoda baltica</i> Sellnick, 1931	Pa Baltic amber
<i>Oripoda</i> sp. in Norton & Poinar (1993)	Ne Dominican amber
Parapirnodus Balogh & Mahunka, 1968	Neogene – Recent
243. <i>Parapirnodus denaius</i> (Woolley, 1971)	Ne Chiapas amber
PARAKALUMMIDAE Grandjean, 1936b	Palaeogene – Recent
Neoribates Berlese, 1914	Palaeogene – Recent
244. <i>Neoribates borussicus</i> Sellnick, 1931	Pa Baltic amber
SCHELORIBATIDAE Grandjean, 1933	Palaeogene – Recent
Liebstadia Oudemans, 1906	Palaeogene – Recent
245. <i>Liebstadia similiformis</i> Sellnick, 1931	Pa Baltic amber
246. <i>Liebstadia similis</i> (Michael, 1888)* [Recent]	Qt Europe / Greenland
Scheloribates Berlese, 1908	Palaeogene – Recent
247. <i>Scheloribates apterus</i> Sellnick, 1931	Pa Baltic amber
248. <i>Scheloribates areatus</i> Sellnick, 1931	Pa Baltic amber
249. <i>Scheloribates durhami</i> (Woolley, 1971)	Ne Chiapas amber
250. <i>Scheloribates initialis</i> (Berlese, 1908) [Recent]	Qt Europe
251. <i>Scheloribates laevigatus</i> (C. L. Koch, 1835) [Recent]	Qt northern Europe
252. <i>Scheloribates latipes</i> (C. L. Koch, 1844) [Recent]	Qt Europe
253. <i>Scheloribates pallidulus</i> (C. L. Koch, 1841) [Recent]	Qt Germany
254. <i>Scheloribates setatus</i> Sellnick, 1931	Pa Baltic amber
SELLNICKIIDAE Balogh & Balogh, 1984	Recent
no fossil record	
STELECHOBATIDAE Grandjean, 1965b	Recent
no fossil record	
SYMBIORIBATIDAE Aoki, 1966b	Recent
no fossil record	
TUBULOZETIDAE Balogh, 1989	Quaternary – Recent
Grandjeanobates Ramsay, 1967	Quaternary – Recent
? <i>Grandjeanobates</i> sp.	Qt New Zealand
ZETOMOTRICHIDAE Grandjean, 1954b	Paleogene – Recent
Zetomotrichidae sp. in Sidorchuk & Norton (2011)	P Baltic amber

CERATOZETOIDEA Jacot, 1925	Paleogene – Recent
CERATOKALUMMIDAE Balogh, 1970	Recent
no fossil record	
CERATOZETIDAE Jacot, 1925	Paleogene – Recent
Ceratozetes Berlese, 1908	Quaternary – Recent
255. <i>Ceratozetes gracilis</i> (Michael, 1884)* [Recent]	Qt Finland
256. <i>Ceratozetes minimus</i> Sellnick, 1928 [Recent]	Qt Germany
257. <i>Ceratozetes parvulus</i> Sellnick, 1922 [Recent]	Qt Germany
Diapterobates Grandjean, 1936b	Quaternary – Recent
258. <i>Diapterobates notatus</i> (Thorell, 1871) [Recent]	Qt Europe / Greenland
Edwardzetes Berlese, 1914	Quaternary – Recent
259. <i>Edwardzetes edwardsi</i> (Nicolet, 1855)* [Recent]	Qt western Norway
Fuscozetes Sellnick, 1928	Quaternary – Recent
260. <i>Fuscozetes fuscipes</i> (C. L. Koch, 1844)* [Recent]	Qt western Norway
Melanozetes Hull, 1916	Paleogene – Recent
261. <i>Melanozetes foderatus</i> Sellnick, 1931	Pa Baltic amber
262. <i>Melanozetes mollicornus</i> [Recent] <i>fossilis</i> Sellnick, 1931	Pa Baltic amber
263. <i>Melanozetes meridianus</i> Sellnick, 1928 [Recent]	Qt Greenland
<i>Melanozetes</i> sp. in Karppinen et al. (1979)	Qt Karelia, Russia
Oromucia Thor, 1930	Quaternary – Recent
264. <i>Oromucia bicuspidata</i> Thor, 1930* [Recent]	Qt western Norway
265. <i>Oromucia lucens</i> (C. L. Koch, date?) [Recent]	Qt Greenland
Sphaerozetes Berlese, 1885	Paleogene – Recent
266. <i>Sphaerozetes convexulus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
267. <i>Sphaerozetes piriformis</i> (Nicolet, 1855) [Recent]	Qt Finland
268. <i>Sphaerozetes primus</i> Sellnick, 1931	Pa Baltic amber
Trichoribates Berlese, 1910	Quaternary – Recent
269. <i>Trichoribates biarea</i> Gjelstrup & Solhøy, 1994 [Recent]	Qt western Norway
270. <i>Trichoribates incisellus</i> (Kramer, 1897) [Recent]	Qt Europe
271. <i>Trichoribates monticola</i> (Trägårdh, 1902) [Recent]	Qt western Norway
272. <i>Trichoribates setiger</i> (Trägårdh, 1910) [Recent]	Qt western Norway
273. <i>Trichoribates trimaculatus</i> (C. L. Koch, 1835)* [Recent]	Qt northern Europe
CHAMOBATIDAE Thor, 1937	Paleogene – Recent
Chamobates Hull, 1916	Paleogene – Recent
274. <i>Chamobates borealis</i> (Trägårdh, 1902) [Recent]	Qt western Norway
275. <i>Chamobates cuspidatus</i> (Michael, 1884) [Recent]	Qt Finland
276. <i>Chamobates difficilis</i> Sellnick, 1931	Pa Baltic amber

EUZETIDAE Grandjean, 1954b	Quaternary – Recent
<i>Euzetes</i> Berlese, 1908	Quaternary – Recent
277. <i>Euzetes globulus</i> (Nicolet, 1855) [Recent]	Qt Finland
HUMEROBATIDAE Grandjean, 1970	Recent
no fossil record	
MYCOBATIDAE Grandjean, 1954b	Quaternary – Recent
<i>Mycobates</i> Hull, 1916	Quaternary – Recent
278. <i>Mycobates consimilis</i> Hammer, 1952 [Recent]	Qt Greenland
279. <i>Mycobates parmelliae</i> (Michael, 1884) [Recent]	Qt Karelia, Russia
280. <i>Mycobates sarekenis</i> (Trägårdh, 1910) [Recent]	Qt western Norway
<i>Punctoribates</i> Berlese, 1908	Quaternary – Recent
281. <i>Punctoribates punctum</i> (C. L. Koch, 1839) [Recent]	Qt Karelia, Russia
282. <i>Punctoribates sellnicki</i> Willmann, 1928 [Recent]	Qt Europe
<i>Punctoribates</i> sp. in Karppinen & Koponen (1973)	Qt Finland
ONYCHOBATIDAE Luxton, 1985	Recent
no fossil record	
RAMSAYELLIDAE Luxton, 1985	Recent
no fossil record	
ZETOMIMIDAE Shaldybina, 1966	Quaternary – Recent
<i>Zetomimus</i> author, date?	Quaternary – Recent
283. <i>Zetomimus furcatus</i> (Pearce & Warburton, 1906)* [Recent]	Qt Karelia, Russia
GALUMNOIDEA Jacot, 1925	Palaeogene – Recent
GALUMNELLIDAE Piffli, 1970	Quaternary – Recent
<i>Galumnella</i> Berlese, 1917	Quaternary – Recent
<i>Galumnella</i> sp. in Aoki (1974)	Qt Mizunami copal
GALUMNIDAE Jacot, 1925	Palaeogene – Recent
Galumnidae spp. in Norton & Poinar (1993)	Pa Baltic amber
<i>Acrogalumna</i> Grandjean, 1956b	Quaternary – Recent
284. <i>Acrogalumna longipluma</i> (Berlese, 1904)* [Recent]	Qt Karelia, Russia
<i>Galumna</i> von Heyden, 1826	Palaeogene – Recent
285. <i>Galumna clavata</i> Sellnick, 1931	Pa Baltic amber
286. <i>Galumna diversa</i> Sellnick, 1931	Pa Baltic amber
287. <i>Galumna lanceata</i> (Oudemans, 1900) [Recent]	Qt Karelia, Russia
288. <i>Galumna obvia</i> (Berlese, 1915) [Recent]	Qt Finland
<i>Galumna</i> sp. in Karppinen & Koponen (1974)	Qt Finland

<i>Pergalumna</i> Grandjean, 1936b	Quaternary – Recent
289. <i>Pergalumna dorsalis</i> (C. L. Koch, 1835) [Recent]	Qt Finland
290. <i>Pergalumna nervosa</i> (Berlese, 1914)* [Recent]	Qt northern Europe
<i>Pilogalumna</i> Grandjean, 1956b	Quaternary – Recent
291. <i>Pilogalumna tenuiclava</i> (Berlese, 1908) [Recent]	Qt Germany

ASTIGMATA G. Canestrini, 1891 (cohort) **Palaeogene – Recent**
 = ACARIDIDA author, date?

SCHIZOGLYPHOIDEA Mahunka, 1978 **Recent**

SCHIZOGLYPHIDAE Mahunka, 1978 **Recent**

no fossil record

HISTIOSTOMATOIDEA Berlese, 1897 **?Palaeogene – Recent**

GUANOLICHIDAE Fain, 1968 **Recent**

no fossil record

HISTIOSTOMATIDAE Berlese, 1897 **?Palaeogene – Recent**

Hististomatidae? [alternatively Acaridae] *in* Dunlop *et al.* (2012) Pa Baltic amber

CANESTRINIOIDEA Berlese, 1884 **Recent**

CANESTRINIIDAE Berlese, 1884 **Recent**

no fossil record

CHETOCHELACARIDAE Fain, 1987 **Recent**

no fossil record

HETEROOPTIDAE Fain, 1967b **Recent**

no fossil record

LEMANNIELLIDAE Wurst, 2001 **Recent**

no fossil record

Superfamily?

[NB: Sidorchuk & Klimov (2011) discussed the problems in placing this extinct family.]

† **GLAESACARIDAE Klimov & Sidorchuk *in* Sidorchuk & Klimov, 2011** **Palaeogene**

† ***Glaesacarus* Klimov & Sidorchuk *in* Sidorchuk & Klimov, 2011** **Palaeogene – Recent**

292. *Glaesacarus rhombeus* (C. L. Koch & Berendt, 1854)* Pa Baltic amber

HEMISCARPOCTOIDEA Oudemans, 1908 **Neogene – Recent**

ALGOPHAGIDAE Fain, 1974 **Recent**

no fossil record

CARPOGLYPHIDAE Oudemans, 1923	Recent
no fossil record	
CHAETODACTYLIDAE Zachvatkin, 1941	Recent
no fossil record	
HEMISARCOPTIDAE Oudemans, 1908	Recent
no fossil record	
HYADESIIDAE Halbert, 1915	Recent
no fossil record	
MELIPONOCOPTIDAE Fain & Rosa, 1983	Recent
no fossil record	
WINTERSCHMIDTIIDAE Oudemans, 1923	Neogene – Recent
† <i>Amphicalvolia</i> Türk, 1963	Neogene – Recent
293. <i>Amphicalvolia hurdi</i> Türk, 1963*	Ne Chiapas amber
GLYCOPHAGOIDEA Berlese, 1897	Recent
AEROGLYPHIDAE Zachvatkin, 1941	Recent
no fossil record	
CHORTOGLYPHIDAE Berlese, 1897	Recent
no fossil record	
ECHIMYOPODIDAE Fain, 1967a	Recent
no fossil record	
EUGLYCYPHAGIDAE Fain & Phillips, 1977	Recent
no fossil record	
GLYCYPHAGIDAE Berlese, 1897	Recent
no fossil record	
PEDETOPODIDAE Fain, 1969	Recent
no fossil record	
ROSENSTEINIIDAE Coorman, 1954	Recent
= LOPHONOTACARIDAE Fain, 1987	
= TROGLOTACARIDAE Fain, 1977	
no fossil record	

ACAROIDEA Latreille, 1802	Neogene – Recent
ACARIDAE Latreille, 1802	Recent
[query family placement?]	
† Tyroglyphites Pampaloni, 1902	Neogene – Recent
294. <i>Tyroglyphites miocenicus</i> Pampaloni, 1902*	Ne Sicily
GAUDIPELLIDAE Atyeo et al., 1974	Recent
= PARTAMONACOPTIDAE author, date?	
= PLATYGLYPHIDAE Kurosa, 1976	
no fossil record	
GLYCACARIDAE Griffiths, 1977	Recent
no fossil record	
LARDOGLYPHIDAE Oudemans, 1877	Recent
no fossil record	
SAPRACARIDAE Fain, 1988	Recent
no fossil record	
SCATOGLYPHIDAE Zachvatkin & Volgin, 1956	Recent
no fossil record	
SUIDASIIDAE Hughes, 1948	Recent
no fossil record	
TYROGLYPHIDAE Donnadieu, 1868	Quaternary – Recent
Tyroglyphidae sp. <i>in</i> Aoki (1974)	Qt Mizunami copal
HYPODERATOIDEA Murray, 1877	Recent
HYPODERATIDAE Murray, 1877	Recent
no fossil record	
PSOROPTIDIA Yunker, 1955 (unranked clade)	Neogene – Recent
PTEROLICHOIDEA Trouessart & Mégnin, 1884	Recent
= FREYANOIDEA Dubinin, 1953	
ASCOURACARIDAE Gaud & Atyeo, 1976	Recent
no fossil record	
CAUDIFERIDAE Gaud & Atyeo, 1978	Recent
no fossil record	
CHEYLABIDIDAE Gaud, 1983	Recent

no fossil record

CRYPTUROPTIDAE Gaud, Atyeo & Berla, 1972 **Recent**

no fossil record

EUSTATHIIDAE Oudemans, 1905 **Recent**

no fossil record

FALCULIFERIDAE Oudemans, 1905 **Recent**

no fossil record

FREYANIDAE Dubinin, 1953 **Recent**

no fossil record

GABUCINIIDAE Gaud & Atyeo, 1975 **Recent**

no fossil record

KIWILICHIDAE Dabert, 1994 **Recent**

no fossil record

KRAMERELLIDAE Gaud & Mouchet, 1961 **Recent**

no fossil record

OCHROLICHIDAE Gaud & Atyeo, 1978 **Recent**

no fossil record

OCONNORIIDAE Gaud, Atyeo & Klompen, 1989 **Recent**

no fossil record

PTEROLICHIDAE Trouessart & Mégnin, 1884 **Recent**

no fossil record

PTILOXENIDAE Gaud, 1982 **Recent**

no fossil record

RECTIJANUIDAE Gaud, 1961 **Recent**

no fossil record

SYRINGOBIIDAE Trouessart, 1897 **Recent**

no fossil record

THORACOSATHESIDAE Gaud & Mouchet, 1959 **Recent**

no fossil record

VEXILLARIIDAE Gaud & Mouchet, 1959	Recent
no fossil record	
ANALGOIDEA Trouessart & Mégnin, 1884	Recent
ALLOPTIDAE Gaud, 1957	Recent
no fossil record	
ANALGIDAE Trouessart & Mégnin, 1884	Recent
no fossil record	
APIONACARIDAE Gaud & Atyeo, 1977	Recent
no fossil record	
AVENZOARIIDAE Oudemans, 1905	Recent
no fossil record	
CYTODITIDAE Oudemans, 1908	Recent
no fossil record	
DERMATIONIDAE Fain, 1965	Recent
no fossil record	
DERMOGLYPHIDAE Mégnin & Trouessart, 1884	Recent
no fossil record	
EPIDERMOPTIDAE Trouessart, 1892	Recent
no fossil record	
GAUDOGLYPHIDAE Bruce & Johnston, 1976	Recent
no fossil record	
HETEROPSORIDAE Oudemans, 1908	Recent
no fossil record	
KNEMIDOKOPTIDAE Dubinin, 1953	Recent
no fossil record	
LAMINOSIOPTIDAE Vitzthum, 1931	Recent
no fossil record	
PROCTOPHYLLODIDAE Mégnin & Trouessart, 1884	Recent
no fossil record	

- PSORALGIDAE Oudemans, 1908** **Recent**
no fossil record
- PSOROPTOIDIDAE Gaud, 1983** **Recent**
no fossil record
- PTERONYSSIDAE Oudemans, 1941** **Recent**
no fossil record
- PTYSSALGIDAE Atyeo & Gaud, 1979** **Recent**
no fossil record
- PYROGLYPHIDAE Cunliffe, 1958** **Recent**
no fossil record
- TARSOCHYLIDAE Atyeo & Gaud, 1979** **Recent**
no fossil record
- THYSANOCERCIDAE Atyeo & Peterson, 1972** **Recent**
no fossil record
- TROUESSARTIIDAE Gaud, 1957** **Recent**
no fossil record
- TURBINOPTIDAE Fain, 1957** **Recent**
no fossil record
- XOLALGIDAE Dubinin, 1953** **Recent**
no fossil record
- SARCOPTOIDEA Murray, 1877** **Neogene–Recent**
= PSOROPTOIDEA Canestrini, 1892
- ACAROPTIDAE Womersley, 1953** **Recent**
no fossil record
- ATOPEMELIDAE Gunter, 1942** **Neogene–Recent**
?Apotomelidae sp. [originally as Listrophoridae in Poinar 1988] Ne Dominican amber
- AUDYCOPTIDAE Lavoipierre, 1964** **Recent**
no fossil record
- CHIRODISCIDAE Trouessart, 1892** **Recent**
no fossil record

CHIRORHYNCHOBIIDAE Fain, 1967 **Recent**

no fossil record

GALAGALIDAE Fain, 1963 **Recent**

no fossil record

GASTRONYSSIDAE Fain, 1956 **Recent**

no fossil record

LEMURNYSIIDAE Fain, 1957 **Recent**

no fossil record

LISTROPHORIDAE Mégnin & Trouessart, 1884 **Recent**

no fossil record

LOBALGIDAE Fain, 1965 **Recent**

no fossil record

MYCOPTIDAE Gunther, 1942 **Recent**

no fossil record

PSOROPTIDAE Canestrini, 1892 **Recent**

no fossil record

PNEUMOCOPTIDAE Fain, 1957 **Recent**

no fossil record

RHYNCOPTIDAE Lawrence, 1956 **Recent**

no fossil record

SARCOPTIDAE Murray, 1877 **Recent**

no fossil record

NOMINA DUBIA

1. *Acarus resinosus* Presl, 1822 Pa Baltic amber
2. *Strieremaeus cordiformatus* Sellnick, 1919 [as *species inquirenda*] Pa Baltic amber

NOMINA NUDA

1. *Erythraeus hirsutissimus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
2. *Gymnodamaeus kulczynskii* Petrunkevitch, 1955a Pa Baltic amber
3. *Trombidium fossile* Keferstein, 1834 Pa Aix-en-Provence?

MISIDENTIFICATIONS

1. *Limnochaeres antiquus* Heyden, 1862 [larval hemipteran insect] Pa Rott, Germany

NON NAMES IN ZOOLOGY

Taxa assigned to living mite genera based on the fossil responses of plant tissue (galls); see discussion in Dunlop & Braddy (2011)

1. *Eriophyes daphnogene* Ambrus & Hably, 1979 [fossil gall] Pa Hungary
2. *Eryophies [sic] vilarrubiae* Villalta, 1957 [fossil gall] Ne Spain
3. *Phytopus antiquus* van Heyden, 1860 [fossil gall] Ne Rott, Germany

c. 36,900 Recent species according to Hallan (2004)

RICINULEI

16 currently valid species of fossil ricinuleid

RICINULEI Thorell, 1876c	Carbon. – Recent
= RHINOASTRA Cook, 1899	
= PODOGONA Cook, 1899	
† PALAEORICINULEI Selden, 1992 (suborder)	Carboniferous – ?Cret.
NB: Wunderlich (2012e) treated the two suborders as superfamilies.	
Ricinulei indet. <i>in</i> Wunderlich (2012e)	K Myanmar amber
† CURCULIOIDIDAE Cockerell, 1916	Carboniferous
† <i>Amarixys</i> Selden, 1992	Carboniferous
1. <i>Amarixys gracilis</i> (Petrunkevitch, 1945a)	C Mazon Creek
2. <i>Amarixys stellaris</i> Selden, 1992	C Mazon Creek
3. <i>Amarixys sulcata</i> (Melander, 1903)*	C Mazon Creek
† <i>Curculioides</i> Buckland, 1837	Carboniferous
4. <i>Curculioides adompha</i> Brauckmann, 1987	C Hagen-Vorhalle
5. <i>Curculioides ansticii</i> Buckland, 1837*	C Coalbrookdale
6. <i>Curculioides eltringhami</i> Petrunkevitch, 1949	C Crawcrook
7. <i>Curculioides gigas</i> Selden, 1992	C Mazon Creek
8. <i>Curculioides granulatus</i> Petrunkevitch, 1949	C Ilkeston
9. <i>Curculioides mcluckiei</i> Selden, 1992	C Mazon Creek
10. <i>Curculioides pococki</i> Selden, 1992	C Coseley
11. <i>Curculioides scaber</i> (Scudder, 1890b)	C Mazon Creek
† POLIOCHERIDAE Scudder, 1884	Carboniferous – ?Cret.
† <i>Poliochera</i> Scudder, 1884	Carboniferous – ?Cret.
12. ? <i>Poliochera cretacea</i> Wunderlich, 2012e	K Myanmar amber
13. <i>Poliochera gibbsi</i> Selden, 1992	C Illinois
14. <i>Poliochera glabra</i> Petrunkevitch, 1913	C Mazon Creek
15. <i>Poliochera punctulata</i> Scudder, 1884*	C Mazon Creek
† <i>Terpsicroton</i> Selden, 1992	Carboniferous
16. <i>Terpsicroton alticeps</i> Selden, 1992*	C Coseley
NEORICINULEI Selden, 1992 (suborder)	Recent
RICINOIDIDAE Ewing, 1929	Recent
= CRYPTOSTEMMIDAE Westwood, 1874	

no fossil record

NOMINA DUBIA

1. *Poliochera* / *Curculioides pustulatus* Laurentiaux-Viera & Laurentiaux, 1963 C Kiaping

55 Recent species according to Harvey (2003)

ARACHNIDA and/or PANTETRAPULMONATA

incertae sedis

3 currently valid, unplaced fossil arachnid and/or tetrapulmonate species

- all three species below have been suggested as possible members of the so-called pantetrapulmonate arachnids; i.e. spiders and their closest relatives

†	<i>Ecchosis</i> Selden & Shear, 1991	Devonian
	1. <i>Ecchosis pulchribothrium</i> Selden & Shear in Selden <i>et al.</i> 1991*	D Gilboa
†	<i>Saccogulus</i> Dunlop, Fayers, Hass & Kerp, 2006	Devonian
	2. <i>Saccogulus seldeni</i> Dunlop, Fayers, Hass & Kerp, 2006*	D Rhynie chert
†	<i>Xenarachne</i> Dunlop & Poschmann, 1997	Devonian
	3. <i>Xenarachne wilwerathensis</i> Dunlop & Poschmann, 1997*	D Willwerath

no Recent species

TRIGONOTARBIDA

65 currently valid species of fossil trigonotarbid

- † **TRIGONOTARBIDA Petrunkevitch, 1949** **Silurian – Permian**
 = ANTHRACOMARTI Karsch, 1882
 = MERIDOGASTRA Thorell & Lindström, 1885
 = EURYMARTI Matthew, 1895
- plesion genus**
- † **Palaeotarbus Dunlop, 1999** **Silurian**
 = † *Eotarbus* Dunlop, 1996 [preoccupied]
 1. *Palaeotarbus jerami* (Dunlop, 1996)* S Ludford Lane
- † **PALAEOCHARINIDAE Hirst, 1923** **Devonian**
- † **Aculeatarbus Shear, Selden & Rolfe, 1987** **Devonian**
 2. *Aculeatarbus depressus* Shear, Selden & Rolfe, 1987* D Gilboa
- † **Gelasinotarbus Shear, Selden & Rolfe, 1987** **Devonian**
 3. *Gelasinotarbus bifidus* Shear, Selden & Rolfe, 1987 D Gilboa
 4. *Gelasinotarbus bonamoae* Shear, Selden & Rolfe, 1987* D Gilboa
 5. *Gelasinotarbus heptops* Shear, Selden & Rolfe, 1987 D Gilboa
 6. *Gelasinotarbus reticulatus* Shear, Selden & Rolfe, 1987 D Gilboa
- † **Gigantocharinus Shear, 2000** **Devonian**
 7. *Gigantocharinus szatmaryi* Shear, 2000* D Red Hill, USA
- † **Gilboarachne Shear, Selden & Rolfe, 1987** **Devonian**
 8. *Gilboarachne griersoni* Shear, Selden & Rolfe, 1987* D Gilboa
- † **Palaeocharinus Hirst, 1923** **Devonian**
 = † *Palaeocharinoides* Hirst, 1923
 9. *Palaeocharinus calmani* Hirst, 1923 D Rhyne cherts
 10. *Palaeocharinus hornei* (Hirst, 1923) D Rhyne cherts
 11. *Palaeocharinus kidstoni* Hirst, 1923 D Rhyne cherts
 12. *Palaeocharinus rhyniensis* Hirst, 1923* D Rhyne cherts
 13. *Palaeocharinus scourfieldi* Hirst, 1923 D Rhyne cherts
 14. *Palaeocharinus tuberculatus* Fayers, Dunlop & Trewin, 2005 D Rhyne cherts
- † **Spiniocharinus Poschmann & Dunlop, 2011** **Devonian**
 15. *Spiniocharinus steinmeyeri* Poschman & Dunlop, 2011* D Bürdenbach
- † **ARCHAEOMARTIDAE Poschmann & Dunlop, 2010** **Devonian**
- † **Archaeomartus Størmer, 1970** **Devonian**
 16. *Archaeomartus levis* Størmer, 1970* D Alken an der Mosel
 i. = *Archaeomartus tuberculatus* Størmer, 1970 D Alken an der Mosel

- † **ANTHRACOMARTIDAE Haase, 1890** **Carboniferous**
- = † PROMYGALIDAE Frič, 1904
- = † BRACHYPYGIDAE Pocock, 1911
- = † CORYPHOMARTIDAE Petrunkevitch, 1945
- = † PLEOMARTIDAE Petrunkevitch, 1945
- † ***Anthracomartus* Karsch, 1882** **Carboniferous**
- = † *Brachylycosa* Frič, 1904
- = † *Cleptomartus* Petrunkevitch, 1949
- = † *Coryphomartus* Petrunkevitch, 1945a
- = † *Cryptomartus* Petrunkevitch, 1945a
- = † *Oomartus* Petrunkevitch, 1953
- = † *Perneria* Frič, 1904
- = † *Pleomartus* Petrunkevitch, 1945a
- = † *Promygale* Frič, 1901
17. *Anthracomartus bohemica* (Frič, 1901) C Nýřany
18. *Anthracomartus carcinoides* (Frič, 1901) C Nýřany
- i. = *Promygale rotundata* Frič, 1901 C Nýřany
- ii. = *Perneria salticoides* Frič, 1904 C ?Nýřany
19. *Anthracomartus elegans* Frič, 1901 C Nýřany
20. *Anthracomartus hindi* Pocock, 1911 C Coseley
- i. = *Cleptomartus hangardi* Guthörl, 1965 C Saar, Germany
- ii. = *Cryptomartus meyeri* Guthörl, 1964 C Aachen
- iii. = *Cleptomartus planus* Petrunkevitch, 1949 C Coseley
- iv. = *Cryptomartus rebskei* Brauckmann, 1984 C Saarbrücken
21. *Anthracomartus granulatus* Frič, 1904 C Nowa Ruda
22. *Anthracomartus janae* (Opluštil, 1986) C Kladno
23. *Anthracomartus kustae* Petrunkevitch, 1953 C Rakovník
24. *Anthracomartus minor* Kušta, 1884 C Rakovník
- i. = *Anthracomartus socius* Kušta, 1888 C Rakovník
25. *Anthracomartus nyranensis* (Petrunkevitch, 1953) C Nýřany
26. *Anthracomartus palatinus* Ammon, 1901 C Brücken, Germany
27. *Anthracomartus preisti* Pocock, 1911 C Coseley
- i. = *Anthracomartus denuiti* Pruvost, 1922 C Charleroi
- ii. = *Cleptomartus plautus* Petrunkevitch, 1949 C Coseley
28. *Anthracomartus radvanicensis* (Opluštil, 1985) C Radvanice
29. *Anthracomartus triangularis* Petrunkevitch, 1913 C Joggins
30. *Anthracomartus trilobitus* Scudder, 1884 C Fayetteville
31. *Anthracomartus voelkelianus* Karsch, 1882* C Europe
- Anthracomartus* sp. in Wright & Selden (2011) C Kansas
- † ***Brachypyge* Woodward, 1878b** **Carboniferous**
32. *Brachypyge carbonis* Woodward, 1878b* C Mons

- † *Maiocercus* Pocock, 1911 **Carboniferous**
 33. *Maiocercus celticus* (Pocock, 1902)* C Coal Measures
 i. = *Maiocercus orbicularis* Gill, 1911 C Westhoughton
- † **ANTHRACOSIRONIDAE** Pocock, 1903a **Devonian – Carbon.**
- † *Anthracosiro* Pocock, 1903a **Carboniferous**
 34. *Anthracosiro fritschii* Pocock, 1903b C Coseley
 i. = *Anthracosiro elongatus* Waterlot, 1934 C Marlebach, France
 35. *Anthracosiro woodwardi* Pocock, 1903a* C Coal Measures
 i. = *Anthracosiro corsini* Pruvost, 1926 C Noeux, France
 ii. = *Anthracosiro latipes* Gill, 1909 C Ryton-on-Tyne, UK
- † *Arianrhoda* Dunlop & Selden, 2004 **Devonian**
 36. *Arianrhoda bennetti* Dunlop & Selden, 2004* D Tredomen
- † *Vratislavia* Frič, 1904 **Carboniferous**
 37. *Vratislavia silesica* (Roemer, 1878)* C Silesia
- † **TRIGONOTARBIDAE** Petrunkevitch, 1949 **Devonian – Carbon.**
- † *Trigonotarbus* Pocock, 1911 **Devonian – Carbon.**
 38. *Trigonotarbus arnoldi* Petrunkevitch, 1955b C Decazeville
 39. *Trigonotarbus johnsoni* Pocock, 1911* C Coseley
 40. *Trigonotarbus stoermeri* Schultka, 1991 D Rheinischen Schief.
- Family uncertain**
- † *Namurotarbus* Poschmann & Dunlop, 2010 **Carboniferous**
 41. *Namurotarbus roessleri* (Dunlop & Brauckmann, 2006)* C Hagen-Vorhalle
- † **LISSOMARTIDAE** Dunlop, 1995 **Carboniferous**
- † *Lissomartus* Petrunkevitch, 1949 **Carboniferous**
 42. *Lissomartus carbonarius* (Petrunkevitch, 1913) C Mazon Creek
 43. *Lissomartus schucherti* (Petrunkevitch, 1913)* C Mazon Creek
- † **APHANTOMARTIDAE** Petrunkevitch, 1945a **Devonian – Permian**
 = † **TRIGONOMARTIDAE** Petrunkevitch, 1949
- † *Alkenia* Størmer, 1970 **Devonian**
 44. *Alkenia mirabilis* Størmer, 1970* D Alken an der Mosel
- † *Aphantomartus* Pocock, 1911 **Carbon. – Permian**
 = † *Trigonomartus* Petrunkevitch, 1913
 = † *Phrynomartus* Petrunkevitch, 1945a
 45. *Aphantomartus areolatus* Pocock, 1911* C–P Coal Measures
 i. = *Aphantomartus pococki* Pruvost, 1912 C Anzin, France
 ii. = *Trigonomartus dorlodoti* Pruvost, 1930 C Rien, France
 iii. = *Eophrynus waechteri* Guthörl, 1938 C Saar

- iv. = ?*Trigonomartus pruvosti* van der Heide, 1951 C Limbourg
v. = ?*Brachylycosa manebachensis* Müller, 1957 C Rotliegenden
46. *Aphantomartus ifeldicus* (Scharf, 1924) P Rotliegend
47. *Aphantomartus pustulatus* (Scudder, 1884) C Coal Measures
i. = ?*Kreischeria villeti* Pruvost, 1912 C Pas de Calais
ii. = *Cleptomartus plötzensis* Simon, 1971 C Halleschen Mulde
- † **KREISCHERIIDAE Haase, 1890** **Carboniferous**
- † **Anzinia Petrunkevitch, 1953** **Carboniferous**
48. *Anzinia thevenini* (Pruvost, 1919)* C Anzin
- † **Gondwanarache Pinto & Hünicken, 1980** **Carboniferous**
49. *Gondwanarache argentinensis* Pinto & Hünicken, 1980* C Bajo de Véliz
- † **Hemikreischeria Frič, 1904** **Carboniferous**
50. *Hemikreischeria geinitzi* (Thevenin, 1902)* C France
- † **Kreischeria Geinitz, 1882** **Carboniferous**
51. *Kreischeria wiedei* Geinitz, 1882* C Zwickau
- † **Pseudokreischeria Petrunkevitch, 1953** **Carboniferous**
52. *Pseudokreischeria pococki* (Gill, 1924) C Crawcrook
i. = *Eophrynus varius* Petrunkevitch, 1949 C Crawcrook
- † **EOPHRYNIDAE Karsch, 1882** **Carboniferous**
= †HEMIPHRYNIDAE Frič, 1904
- † **Eophrynus Woodward, 1871b** **Carboniferous**
53. *Eophrynus prestvicii* (Buckland, 1837)* C Coalbrookdale
54. *Eophrynus udus* Brauckmann, Koch & Kemper, 1985 C Hagen-Vorhalle
- † **Nyranytarbus Harvey & Selden, 1995** **Carboniferous**
= †*Hemiphrynus* Frič, 1901 [preoccupied]
55. *Nyranytarbus hofmanni* (Frič, 1901) C Nýřany
56. *Nyranytarbus longipes* (Frič, 1901)* C Nýřany
- † **Petrovicia Frič, 1904** **Carboniferous**
57. *Petrovicia proditoria* Frič, 1904* C Petrovice
- † **Planomartus Petrunkevitch, 1953** **Carboniferous**
58. *Planomartus krejcii* (Kušta, 1883)* C Rakovník
i. = *Anthracomartus affinis* Kušta, 1885 C Rakovník
- † **Pleophrynus Petrunkevitch, 1945a** **Carboniferous**
59. *Pleophrynus verrucosus* (Pocock, 1911) C Coal Measures
i. = *Eophrynus warei* Dix & Pringle, 1930 C Glyncoch, UK
ii. = *Pleophrynus ensifer* Petrunkevitch, 1945a* C Mazon Creek
iii. = *Eophrynus jugatus* Ambrose & Romano, 1972 C Kilmersdon, UK
- † **Pocononia Petrunkevitch, 1953** **Carboniferous**
60. *Pocononia whitei* (Ewing, 1930)* C Pocono Shales
- † **Somaspidion Jux, 1982** **Carboniferous**

61. *Somaspidion hammapheron* Jux, 1982* C Dinslaken
† ***Stenotrogulus Frič, 1904*** **Carboniferous**
 = † *Cyclotrogulus Frič, 1904*
 = † *Pseudoeophrynus Příbyl, 1958*
62. *Stenotrogulus salmii* (Stur, 1877)* C Ostrava
 i. = *Cyclotrogulus sturii* Frič, 1904 [*non* Hasse, 1890] C Ostrava
 ii. = *Pseudoeophrynus ostraviensis* Příbyl, 1958 C Ostrava

TRIGONOTARBIDA *incertae sedis*

- † ***Anthracophrynus André, 1913*** **Carboniferous**
 63. *Anthracophrynus tuberculatus* André, 1913* C Dudweiler
- † ***Areomartus Petrunkevitch, 1913*** **Carboniferous**
 64. *Areomartus ovatus* Petrunkevitch, 1913* C West Virginia
- † **'*Eophrynus***
 65. '*Eophrynus*' *scharfi* Scharf, 1924 P Rotliegend

NOMINA DUBIA

1. *Anthracomartus buchi* (Goldenberg, 1873) C Saarbrücken
2. *Anthracomartus hageni* (Goldenberg, 1873) C Saarbrücken
3. *Elaverimartus pococki* Petrunkevitch, 1953 C Ellismuir
4. *Eurymartus latus* Matthew, 1895 C Fern Ledges
5. ?*Eurymartus spinulosus* Matthew, 1895 C Fern Ledges
6. *Trigonomartus woodruffi* (Scudder, 1893) C Rhode Island

no Recent species

URARANEIDA

2 currently valid species of uraraneid

- The uraraneids were previously interpreted as true spiders (Araneae), but are now thought to be a more basal lineage which produced silk but lacked spinnerets.

† **URARANEIDA Selden & Shear *in* Selden *et al.*, 2008** Devonian – Permian

† ***Attercopus* Selden & Shear *in* Selden *et al.* (1991)** Devonian

1. *Attercopus fimbriunguis* (Shear, Selden & Rolfe, 1987)* D Gilboa, New York

† **PERMARACHNIDAE Eskov & Selden, 2005** Permian

† ***Permarachne* Eskov & Selden, 2005** Permian

2. *Permarachne novokshonovi* Eskov & Selden, 2005* P Matveyevka

ARANEAE

1,185 currently valid species of fossil spider

ARANEAE Clerck, 1757	Carbon. – Recent
‘mesotheles’	Carbon. – Recent
† ARTHROLYCOSIDAE Frič, 1904	Carboniferous
† <i>Arthrolycosa</i> Harger, 1874	Carbon. – Permian
1. <i>Arthrolycosa antiqua</i> Harger, 1874*	C Mazon Creek
2. <i>Arthrolycosa danielsi</i> Petrunkevitch, 1913	C Mazon Creek
<i>Arthrolycosa</i> sp. in Eskov & Selden (2005)	P Kityak river
† <i>Eocteniza</i> Pocock, 1911	Carboniferous
3. <i>Eocteniza silvicola</i> Pocock, 1911*	C Coseley
† ARTHROMYGALIDAE Petrunkevitch, 1923	Carboniferous
† <i>Arthromygale</i> Petrunkevitch, 1923	Carboniferous
4. <i>Arthromygale fortis</i> (Frič, 1904)*	C Rakovník
i. = <i>Arthrolycosa beecheri</i> Frič, 1904	C Rakovník
† <i>Eolycosa</i> Kušta, 1885	Carboniferous
5. <i>Eolycosa lorenzi</i> Kušta, 1885*	C Rakovník
† <i>Geralycosa</i> Kušta, 1888	Carboniferous
6. <i>Geralycosa fritschi</i> Kušta, 1888*	C Rakovník
† <i>Kustaria</i> Petrunkevitch, 1953	Carboniferous
= † <i>Scudderia</i> Kušta, 1888 [preoccupied]	
7. <i>Kustaria carbonaria</i> (Kušta, 1888)*	C Rakovník
† <i>Palaranaea</i> Frič, 1873	Carboniferous
8. <i>Palaranaea borassifoliae</i> Frič, 1873*	C Czech Republic
† <i>Protocteniza</i> Petrunkevitch, 1949	Carboniferous
9. <i>Protocteniza britannica</i> Petrunkevitch, 1949*	C Coseley
† <i>Protolycosa</i> Roemer, 1866	Carboniferous
10. <i>Protolycosa anthracophila</i> Roemer, 1866*	C Silesia
11. <i>Protolycosa cebennensis</i> Laurentiaux-Viera & Laurentiaux, 1963	C Cévennes, France
† <i>Rakovnicia</i> Kušta, 1884a	Carboniferous
12. <i>Rakovnicia antiqua</i> Kušta, 1884a*	C Rakovník
† PYRITARANEIDAE Petrunkevitch, 1953	Carboniferous
† <i>Dinopilio</i> Frič, 1904	Carboniferous
13. <i>Dinopilio gigas</i> Frič, 1904*	C Rakovník

14. *Dinopilo parvus* Petrunkevitch, 1953 C Kent, UK
- † *Pyritaranea* Frič, 1901 **Carboniferous**
15. *Pyritaranea tubifera* Frič, 1901* C Nyřany
- MESOTHELAE Pocock, 1892** **Carbon. – Recent**
- plesion genus**
- † *Palaeothele* Selden, 2000 **Carboniferous**
- = † *Eothele* Selden, 1996 [preoccupied]
16. *Palaeothele montceauensis* (Selden, 1996)* C Montceau-les-Mines
- LIPHISTIIDAE Pocock, 1892** **Recent**
- = HEPTATHELIDAE Haupt, 1983
- no fossil record
- OPISTHOTHELAE Pocock, 1892** **Triassic – Recent**
- Opisthotelae incertae sedis**
- † *Eoatypus* McCook, 1888 **Palaeogene**
17. *Eoatypus woodwardii* McCook, 1888* Pa Isle of Wight
- MYGALOMORPHAE Pocock, 1892** **Triassic – Recent**
- Mygalomorpha indet. 1–3 in Wunderlich (2008d) K Myanmar amber
- ATYPOIDEA Thorell, 1870a** **Triassic – Recent**
- † *Friularachne* Dalla Vecchia & Selden, 2013 **Triassic**
18. *Friularachne rigoi* Dalla Vecchia & Selden, 2013* Tr Friurli, Italy
- ATYPIDAE Thorell, 1870a** **Cretaceous – Recent**
- = CALOMMATOIDAE Thorell, 1887
- † *Ambiortiphagus* Eskov & Zonstein, 1990 **Cretaceous**
19. *Ambiortiphagus ponomarenkoi* Eskov & Zonstein, 1990* K Central Mongolia
- † *Balticatypus* Wunderlich, 2011h **Palaeogene**
20. *Balticatypus beigeli* Wunderlich, 2011h Pa Baltic amber
21. *Balticatypus juvenis* Wunderlich, 2011h* Pa Baltic amber
22. *Balticatypus spinosus* Wunderlich, 2011h Pa Baltic amber
- ANTRODIAETIDAE Gertsch in Comstock, 1940** **Cretaceous – Recent**
- = BRACHYBOTHRIDAE Simon, 1892
- = ACCATYMIDAE Kishida, 1930
- † *Cretacattyma* Eskov & Zonstein, 1990 **Cretaceous**
23. *Cretacattyma raveni* Eskov & Zonstein, 1990* K Central Mongolia
- MECICOBOTHRIIDAE Holmberg, 1882** **Cretaceous – Recent**
- = HEXURIDAE Simon, 1889b
- † *Cretohexura* Eskov & Zonstein, 1990 **Cretaceous**

24. <i>Cretohexura coylei</i> Eskov & Zonstein, 1990*	K Transbaikalia
† Cretomegahexura Eskov & Zonstein, 1990	Cretaceous
25. <i>Cretomegahexura platnicki</i> Eskov & Zonstein, 1990*	K Central Mongolia
HEXATHELIDAE Simon, 1892 <i>b</i>	Triassic – Recent
† Rosamygale Selden & Gall, 1992	Triassic
26. <i>Rosamygale grauvogeli</i> Selden & Gall, 1992*	Tr Vosges, France
DIPLURIDAE Simon, 1889 <i>b</i>	Cretaceous – Recent
† Clostes Menge, 1869	Palaeogene
27. <i>Clostes priscus</i> Menge, 1869*	Pa Baltic / Bitt. amber
† Cretadiplura Selden <i>in</i> Selden <i>et al.</i> , 2006	Cretaceous
28. <i>Cretadiplura ceara</i> Selden <i>in</i> Selden <i>et al.</i> , 2006*	K Crato Formation
† Dinodiplura Selden <i>in</i> Selden <i>et al.</i> , 2006	Cretaceous
29. <i>Dinodiplura ambulacra</i> Selden <i>in</i> Selden <i>et al.</i> , 2006*	K Crato Formation
Ischnothele Ausserer, 1875	?Neogene – Recent
? <i>Ischnothele</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
Masteria L. Koch, 1873	Neogene – Recent
= † <i>Microsteria</i> Wunderlich, 1988	
30. <i>Masteria sexoculata</i> (Wunderlich, 1988)	Ne Dominican amber
? <i>Masteria</i> sp. <i>in</i> Schawaller (1982 <i>c</i> : as ? <i>Ischnothele</i>)	Ne Dominican amber
genus uncertain	
Dipluridae sp. 1–3 <i>in</i> Wunderlich (2004 <i>a</i>)	Pa Baltic amber
Dipluridae sp. <i>in</i> Wunderlich (2004 <i>a</i>)	Ne Dominican amber
Dipluridae indet. <i>in</i> Wunderlich (2012 <i>d</i>)	K Myanmar amber
CYRTAUCHENIIDAE Simon, 1892 <i>b</i>	Neogene – Recent
Bolostromus Ausserer, 1875	Neogene – Recent
31. <i>Bolostromus destructus</i> Wunderlich, 1988	Ne Dominican amber
CTENIZIDAE Thorell, 1887	Palaeogene – Recent
= HALONOPROCTIDAE Pocock, 1903	
† Baltocteniza Eskov & Zonstein, 2000	Palaeogene
32. <i>Baltocteniza kulicka</i> Eskov & Zonstein, 2000	Pa Baltic amber
† Electrocteniza Eskov & Zonstein, 2000	Palaeogene
33. <i>Electrocteniza sadilenkoi</i> Eskov & Zonstein, 2000	Pa Baltic amber
Ummidia Thorell, 1875	Palaeogene – Recent
34. <i>Ummidia damzeni</i> Wunderlich, 2000	Pa Baltic amber
35. <i>Ummidia malinowskii</i> Wunderlich, 2000	Pa Baltic amber
<i>Ummidia</i> sp. <i>in</i> Wunderlich (2004 <i>a</i>)	Pa Baltic amber
? <i>Ummidia</i> sp. <i>in</i> Wunderlich (2011 <i>h</i>)	Pa Baltic amber

EUCTENIZIDAE Raven, 1985	Recent
no fossil record	
IDIOPIDAE Simon, 1892b	Recent
no fossil record	
ACTINOPODIDAE Simon, 1892b	Recent
= ERIODONTIDAE C. L. Koch & Berendt, 1854	
[based on a generic synonym; listed in Bonnet as syn. of Clubionidae!]	
no fossil record	
MIGIDAE Simon, 1892b	Recent
no fossil record	
NEMESIIDAE Simon, 1892b	Cretaceous – Recent
= PYCNOTHELIDAE Chamberlin, 1917	
† <i>Cretamygale</i> Selden, 2002	Cretaceous
36. <i>Cretamygale chasei</i> Selden, 2002*	K Isle of Wight
† <i>Eodiplurina</i> Petrunkevitch, 1922	Palaeogene
[NB: Selden (2001) questioned this familial placement based on claw structure]	
37. <i>Eodiplurina cockerelli</i> Petrunkevitch, 1922*	Pa Florissant
MICROSTIGMATIDAE Roewer, 1942	Neogene – Recent
= MICROMYGALIDAE Wunderlich, 2004b	
† <i>Parvomygale</i> Wunderlich, 2004b	Neogene
38. <i>Parvomygale distincta</i> Wunderlich, 2004b*	Ne Dominican amber
BARYCHELIDAE Simon, 1889b	Neogene – Recent
<i>Psalistops</i> Simon, 1889b	Neogene – Recent
39. <i>Psalistops hispaniolensis</i> Wunderlich, 1988*	Ne Dominican amber
THERAPHOSIDAE Thorell, 1870a	Neogene – Recent
= AVICULARIIDAE Simon, 1874	
Theraphosidae gen. et sp. indet. in Dunlop <i>et al.</i> (2008)	Ne Chiapas amber
<i>Hemirraghus</i> Simon, 1903	Neogene – Recent
<i>Hemirraghus</i> sp. in García-Villafuerte (2008)	Ne Chiapas amber
† <i>Ischnocolinopsis</i> Wunderlich, 1988	Neogene
40. <i>Ischnocolinopsis acutus</i> Wunderlich, 1988*	Ne Dominican amber
PARATROPIDIDAE Simon, 1889a	Recent
no fossil record	
ARANEOMORPHAE Smith, 1902	Triassic – Recent

ARANEOMORPHAE indet.

- † *Argyrarachne* Selden *in* Selden *et al.*, 1999 **Triassic**
 41. *Argyrarachne solitus* Selden *in* Selden *et al.*, 1999* Tr Virginia
- † *Triassaraneus* Selden *in* Selden *et al.*, 1999 **Triassic**
 42. *Triassaraneus andersonorum* Selden *in* Selden *et al.*, 1999* Tr KwaZulu-Natal

HYPOCHILIDAE Marx, 1888 **Recent**

= ECTATOSTICTIDAE Lehtinen, 1967

no fossil record

AUSTROCHILOIDEA Zapfe, 1955 **Recent****AUSTROCHILIDAE Zapfe, 1955** **Recent**

= THAIDIDAE Lehtinen, 1967

= HICKMANIIDAE Lehtinen, 1967

no fossil record

GRADUNGULIDAE Forster, 1955 **Recent**

no fossil record

ARANEOCLADA Platnick, 1977 **Triassic – Recent****HAPLOGYNAE Simon, 1893** **Jurassic – Recent****FILISTATIDAE Ausserer, 1867** **Neogene – Recent***Misionella* Ramírez & Grismado, 1997 **Neogene – Recent**

- 43.
- Misionella didicostae*
- Penney, 2005a Ne Dominican amber

SICARIIDAE Keyserling, 1880a **Neogene – Recent**

= LOXOSCELIDAE Simon, 1893

Loxosceles Heineken & Lowe, 1832 **Neogene – Recent**

- 44.
- Loxosceles aculicaput*
- Wunderlich, 2004c Ne Dominican amber

- 45.
- Loxosceles defecta*
- Wunderlich, 1988 Ne Dominican amber

- 46.
- Loxosceles deformis*
- Wunderlich, 1988 Ne Dominican amber

- Loxosceles*
- sp.
- in*
- Wunderlich (1988) Ne Dominican amber

SCYTODIDAE Blackwall, 1864 **?Cretaceous – Recent**

- Scytodidae sp. 1–2
- in*
- Wunderlich (2004b) Pa Bitterfeld amber

Scytodes Latreille, 1804a **?Cretaceous – Recent**

47. ?
- Scytodes hani*
- Wunderlich, 2012d K Jordanian amber

- 48.
- Scytodes marginalis*
- Wunderlich, 2004as Qt Madagascan copal

- 49.
- Scytodes piliformis*
- Wunderlich, 1988 Ne Dominican amber

- 50.
- Scytodes planithorax*
- Wunderlich, 1988 Ne Dominican amber

- 51.
- Scytodes stridulans*
- Wunderlich, 1988 Ne Dominican amber

- 52.
- Scytodes weitschati*
- Wunderlich, 1993a Pa Baltic amber

<i>Scytodes</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
<i>Scytodes</i> sp. <i>in</i> Wunderlich (2011 <i>h</i>)	Pa Baltic amber
PERIEGOPIDAE Simon, 1893	Recent
no fossil record	
DRYMUSIDAE Simon, 1893	Recent
no fossil record	
† PRAETERLEPTONETIDAE Wunderlich 2008<i>d</i>	Cretaceous
<i>Praeterleptonetidae</i> indet. <i>in</i> Wunderlich (2008 <i>d</i>)	K Myanmar amber
† <i>Palaeohygropoda</i> Penney, 2004<i>c</i>	Cretaceous
53. <i>Palaeohygropoda myanmarensis</i> Penney, 2004 <i>c</i> *	K Myanmar amber
† <i>Praeterleptoneta</i> Wunderlich, 2008<i>d</i>	Cretaceous
54. <i>Praeterleptoneta spinipes</i> Wunderlich, 2008 <i>d</i> *	K Myanmar amber
55. <i>Praeterleptoneta tibialis</i> Wunderlich, 2011 <i>i</i>	K Myanmar amber
† PHOLCOCHYROCERIDAE Wunderlich 2012<i>d</i>	Cretaceous
† <i>Pholcochyrocer</i> Wunderlich, 2008<i>d</i>	Cretaceous
56. ? <i>Pholcochyrocer baculum</i> Wunderlich, 2012 <i>d</i>	K Myanmar amber
57. <i>Pholcochyrocer guttulaequeae</i> Wunderlich, 2008 <i>d</i> *	K Myanmar amber
58. <i>Pholcochyrocer pecten</i> Wunderlich, 2012 <i>d</i>	K Myanmar amber
LEPTONETIDAE Simon, 1890	Cretaceous – Recent
† <i>Eoleptoneta</i> Wunderlich, 1991	Palaeogene
59. <i>Eoleptoneta curvata</i> Wunderlich, 2004 <i>c</i>	Pa Bitterfeld amber
60. <i>Eoleptoneta duocalcar</i> Wunderlich, 2004 <i>c</i>	Pa Baltic amber
61. <i>Eoleptoneta kutscheri</i> Wunderlich, 1991*	Pa Bitterfeld amber
62. <i>Eoleptoneta multispinae</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
63. <i>Eoleptoneta pseudoarticulata</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
64. <i>Eoleptoneta similis</i> Wunderlich, 2004 <i>c</i>	Pa Baltic amber
† <i>Oligoleptoneta</i> Wunderlich 2004<i>c</i>	Palaeogene
65. <i>Oligoleptoneta altoculus</i> Wunderlich 2004 <i>c</i> *	Pa Baltic amber
66. <i>Oligoleptoneta cymbiospina</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
† <i>Palaeoleptoneta</i> Wunderlich 2012<i>d</i>	Cretaceous
67. <i>Paleoleptoneta calcar</i> Wunderlich, 2012 <i>d</i> *	K Myanmar amber
TELEMIDAE Fage, 1913	Palaeogene – Recent
<i>Telema</i> Simon, 1882	Palaeogene – Recent
68. ? <i>Telema moritzi</i> Wunderlich, 2004 <i>c</i>	Pa Baltic / Bitt. amber
OCHYROCERATIDAE Fage, 1912	Neogene – Recent

= † EOPSILODERCIDAE Wunderlich, 2008d

[NB: Wunderlich (2012d) recognised this as a junior synonym of Psilodercidae; Platnick does not recognise this family]

?Eopsilodercidae indet. 1–3 in Wunderlich (2008d)	K Myanmar amber
† Arachnolithulus Wunderlich, 1988	Neogene
69. <i>Arachnolithulus longipes</i> Wunderlich, 2004c	Ne Dominican amber
70. <i>Arachnolithulus pygmaeus</i> Wunderlich, 1988*	Ne Dominican amber
? <i>Arachnolithulus</i> sp. in Wunderlich (1988)	Ne Dominican amber
† Eopsiloderces Wunderlich, 2008d	Cretaceous
71. <i>Eopsiloderces loxosceloides</i> Wunderlich, 2008d	K Myanmar amber
† Furcembolus Wunderlich, 2008d	Cretaceous
72. <i>Furembolus andersoni</i> Wunderlich, 2008d	K Myanmar amber
Leclercera Deeleman-Reinhold, 1995	Cretaceous – Recent
73. <i>Leclercera longissipes</i> Wunderlich, 2012d	K Myanmar amber
74. <i>Leclercera spicula</i> Wunderlich, 2012d	K Myanmar amber
Psiloderces Simon, 1892	?Cretaceous – Recent
75. ? <i>Psiloderces filiformis</i> Wunderlich, 2012d	K Myanmar amber
PHOLCIDAE C. L. Koch, 1851	Palaeogene – Recent
Pholcidae sp. 1–2 in Wunderlich (2004b)	Pa Baltic amber
Pholcidae sp. in Wunderlich (2004au)	Pa Fu Shun amber
Coryssocnemis Simon, 1893	Neogene – Recent
76. ? <i>Coryssocnemis velteni</i> Wunderlich, 2004c	Ne Dominican amber
Leptopholcus Simon, 1893	Neogene
77. <i>Leptopholcus kiskeya</i> Huber & Wunderlich, 2006	Ne Dominican amber
Modisimus Simon, 1893	Neogene – Recent
78. <i>Modisimus calcar</i> Wunderlich, 1988	Ne Dominican amber
79. <i>Modisimus calcaroides</i> Wunderlich, 1988	Ne Dominican amber
80. <i>Modisimus crassifemoralis</i> Wunderlich, 1988	Ne Dominican amber
81. <i>Modisimus oculatus</i> Wunderlich, 1988	Ne Dominican amber
82. <i>Modisimus tuberosus</i> Wunderlich, 1988	Ne Dominican amber
<i>Modisimus</i> sp. in Wunderlich (1988)	Ne Dominican amber
† Paraspermophora Wunderlich, 2004c	Palaeogene
83. <i>Paraspermophora bitterfeldensis</i> Wunderlich, 2004c	Pa Bitterfeld amber
84. <i>Paraspermophora perplexa</i> Wunderlich, 2004c*	Pa Baltic amber
<i>Paraspermophora</i> sp. in Wunderlich (2004c, 2011h)	Pa Baltic / Bitt. amber
Pholcophora Banks, 1896	Neogene – Recent
85. <i>Pholcophora brevipes</i> Wunderlich, 1988	Ne Dominican amber
86. <i>Pholcophora gracilis</i> Wunderlich, 1988	Ne Dominican amber
87. <i>Pholcophora longicornis</i> Wunderlich, 1988	Ne Dominican amber
Quamtana Huber, 2003	Palaeogene – Recent

88. *Quamtana huberi* Penney, 2007a Pa Le Quesnoy amber
† **Serratochorus Wunderlich, 1988** **Neogene**
89. *Serratochorus pygmaeus* Wunderlich, 1988* Ne Dominican amber
- PLECTREURIDAE Simon, 1893** **Jurassic – Recent**
† **Eoplectreurys Selden & Huang, 2010** **Jurassic**
90. *Eoplectreurys gertschi* Selden & Huang, 2010 J Daohugou
† **Palaeoplectreurys Wunderlich, 2004c** **Palaeogene**
91. *Palaeoplectreurys baltica* Wunderlich, 2004c* Pa Baltic amber
Plectreurys Simon, 1893 **Neogene – Recent**
92. *Plectreurys pittfieldi* Penney, 2009 Ne Dominican amber
- DIGUETIDAE F. O. P.-Cambridge, 1899** **Recent**
no fossil record
- CAPONIIDAE Simon, 1890** **Neogene – Recent**
= COLOPHONIDAE O. P.-Cambridge, 1874 [based on a generic homonym]
Nops MacLeay, 1839 **Neogene – Recent**
93. *Nops lobatus* Wunderlich, 1988 Ne Dominican amber
i. = *Nops segmentatus* Wunderlich, 1988 Ne Dominican amber
Nops sp. *in* Wunderlich (1988) Ne Dominican amber
- TETRABLEMMIDAE O. P.-Cambridge, 1873** **Palaeogene – Recent**
= PHAEDOMOIDAE Thorell, 1890 [based on a generic homonym]
= PACULLIDAE Simon, 1894
Tetrablemmidae gen. indet. *in* Wunderlich (2012d) K Myanmar amber
† **Balticoblemma Wunderlich, 2004c** **Palaeogene**
94. *Balticoblemma unicorniculum* Wunderlich, 2004c* Pa Baltic amber
† **Eogamasomorpha Wunderlich, 2008d** **Cretaceous**
95. *Eogamasomorpha nubila* Wunderlich, 2008d* K Myanmar amber
† **Eoscaphiella Wunderlich, 2011i** **Cretaceous**
96. *Eoscaphiella ohlhoffi* Wunderlich, 2011i* K Myanmar amber
Monoblemma Gertsch, 1941 **Neogene**
97. ?*Monoblemma spinosum* Wunderlich, 1988* Ne Dominican amber
† **Saetosoma Wunderlich, 2012d** **Cretaceous**
98. *Saetosoma filiembolus* Wunderlich, 2012d* K Myanmar amber
- TROGLORAPTORIDAE Griswold, Audisio & Ledford, 2012** **Recent**
no fossil record
- DYSDEROIDEA Bristowe, 1938** **Cretaceous – Recent**
?Dysderoidea s. l. indet 1–2 *in* Wunderlich (2008d) K Myanmar amber

SEGESTRIIDAE Simon, 1893	Cretaceous – Recent
?Segestriidae indet <i>in</i> Wunderlich (2008d)	K Myanmar amber
Ariadna Audouin, 1826	Cretaceous – Recent
99. ? <i>Ariadna amissicoli</i> Wunderlich, 2008d	K Jordanian amber
100. <i>Ariadna copalis</i> Wunderlich, 2008a	Qt ?Madagascan copal
101. <i>Ariadna defuncta</i> Wunderlich 2004c	Pa Bitterfeld amber
102. <i>Ariadna hintzei</i> Wunderlich, 2004as	Qt Madagascan copal
103. <i>Ariadna ovalis</i> Wunderlich, 2008a	Pa Baltic amber
104. <i>Ariadna parva</i> Wunderlich, 2008a	Pa Baltic amber
105. <i>Ariadna paucispinosa</i> Wunderlich, 1988	Ne Dominican amber
106. <i>Ariadna resiniae</i> Hickman, 1957	Ne? Australian copal
? <i>Ariadna</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
† Lebansegestria Wunderlich 2008d	Cretaceous
107. <i>Lebansegestria azari</i> Wunderlich, 2008d*	K Lebanese amber
† Microsegestria Wunderlich & Milki, 2004	Cretaceous
108. <i>Microsegestria poinari</i> Wunderlich & Milki, 2004*	K Lebanese amber
† Palaeosegestria Penney, 2004a	Cretaceous
109. <i>Palaeosegestria lutzii</i> Penney, 2004a*	K New Jersey amber
Segestria Latreille, 1804a	Cretaceous – Recent
110. <i>Segestria cristata</i> Menge <i>in</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
111. <i>Segestria flexio</i> Wunderlich, 2004c	Pa Baltic amber
112. <i>Segestria mortalis</i> Wunderlich 2004c	Pa Baltic amber
113. <i>Segestria plicata</i> Petrunkevitch, 1950	Pa Baltic amber
114. <i>Segestria scudderi</i> Petrunkevitch, 1922	Pa Florissant
115. <i>Segestria secessa</i> Scudder, 1890a	Pa Florissant
116. <i>Segestria succinei</i> Berland, 1939	Pa Baltic amber
117. <i>Segestria tomentosa</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
i. = <i>Segestria plicata</i> Petrunkevitch, 1950 [provisional]	Pa Baltic amber
<i>Segestria</i> sp. <i>in</i> Penney (2002)	K New Jersey amber
<i>Segestria</i> sp. <i>in</i> Wunderlich (2004c)	Pa Baltic amber
† Vetsegestria Wunderlich, 2004c	Palaeogene
118. <i>Vetsegestria quinquespinosa</i> Wunderlich, 2004c*	Pa Bitterfeld amber
DYSDERIDAE C. L. Koch, 1837	Palaeogene – Recent
† Dasumiana Wunderlich, 2004c	Palaeogene
119. <i>Dasumiana emicans</i> Wunderlich, 2004c*	Pa Baltic amber
120. ? <i>Dasumiana subita</i> (Petrunkevitch, 1958)	Pa Baltic amber
121. <i>Dasumiana valga</i> Wunderlich, 2004c	Pa Baltic amber
Dysdera Latreille, 1804	Palaeogene – Recent
122. <i>Dysdera dilatata</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
Harpactea Bristowe, 1939	Palaeogene – Recent

123.	<i>Harpactea communis</i> Wunderlich, 2004c	Pa	Baltic amber
124.	<i>Harpactea extincta</i> Petrunkevitch, 1950	Pa	Baltic amber
125.	<i>Harpactea hombergi</i> (Scopoli, 1763) [Recent]	Qt	England
126.	<i>Harpactea longibulbus</i> Wunderlich, 2011h	Pa	Baltic amber
127.	<i>Harpactea tersa</i> (C. L. Koch & Berendt, 1854) ... [provisional transfer]	Pa	Baltic amber
	<i>Harpactea</i> sp. in Wunderlich (2011h)	Pa	Bitterfeld amber
Dysderidae?			
†	<i>Mistura</i> Petrunkevitch, 1971		Neogene
128.	<i>Mistura perplexa</i> Petrunkevitch, 1971*	Ne	Chiapas amber
OONOPIDAE Simon, 1890 Cretaceous – Recent			
	Oonopidae gen. et sp. in Penney (2002)	K	New Jersey amber
†	<i>Burmorchestina</i> Wunderlich, 2008a		Cretaceous
129.	<i>Burmorchestina pulcher</i> Wunderlich, 2008a*	K	Myanmar amber
†	<i>Canadaorchestina</i> Wunderlich, 2008a		Cretaceous
130.	<i>Canadaorchestina albertensis</i> (Penney, 2006a)*	K	Manitobian amber
†	<i>Fossilopaea</i> Wunderlich, 1988		Neogene
131.	<i>Fossilopaea sulci</i> Wunderlich, 1988*	Ne	Dominican amber
<i>Heteroonops</i> Dalmás, 1916 ?Neogene – Recent			
	<i>Heteroonops</i> sp. in Wunderlich (1988)	Ne	Dominican amber
<i>Opopaea</i> Simon, 1891 ?Neogene – Recent			
	? <i>Opopaea</i> sp. in Wunderlich (1988)	Ne	Dominican amber
<i>Orchestina</i> Simon, 1882 Cretaceous – Recent			
132.	<i>Orchestina (Baltorchestina) angulata</i> Wunderlich, 2012f [replacement name].....	Pa	Bitterfeld amber
	i. = <i>Orchestina (B.) rectangulata</i> Wunderlich, 2011h [preoccupied]		
133.	<i>Orchestina baltica</i> Petrunkevitch, 1942	Pa	Baltic amber
134.	<i>Orchestina (Baltorchestina) bitterfeldensis</i> Wunderlich, 2008a	Pa	Bitterfeld amber
135.	<i>Orchestina breviembolus</i> Wunderlich, 1981	Pa	Baltic amber
136.	<i>Orchestina (Baltorchestina) brevis</i> Wunderlich, 2008a	Pa	Baltic amber
137.	<i>Orchestina crassimbolus</i> Wunderlich, 1981	Pa	Baltic amber
138.	<i>Orchestina (Baltorchestina) crassipatellaris</i> Wunderlich, 1981	Pa	Baltic amber
139.	<i>Orchestina (Baltorchestina) crassitibialis</i> Wunderlich, 1981	Pa	Baltic amber
140.	<i>Orchestina (Baltorchestina) colchembolus</i> Wunderlich, 1981	Pa	Baltic amber
141.	<i>Orchestina colombiensis</i> Wunderlich, 2004at	Qt	Colombian copal
142.	<i>Orchestina dominicana</i> Wunderlich, 1981	Ne	Dominican amber
143.	<i>Orchestina forceps</i> Wunderlich, 1981	Pa	Baltic amber
144.	<i>Orchestina (Baltorchestina) forfex</i> Wunderlich, 2011h	Pa	Baltic amber
145.	<i>Orchestina (Baltorchestina) furca</i> Wunderlich, 1981	Pa	Baltic amber
146.	<i>Orchestina fushunensis</i> Wunderlich, 2004au	Pa	Fu Shun amber

147. <i>Orchestina gappi</i> Saupe et al., 2012	K Archingeay amber
148. <i>Orchestina gracilitibialis</i> Wunderlich, 2004c	Pa Baltic amber
149. <i>Orchestina (Baltorchestina) imperialis</i> Petrunkevitch, 1963	Pa Baltic/Bitter. amber
150. <i>Orchestina kenyana</i> Wunderlich, 1981	Qt East African copal
151. <i>Orchestina longimana</i> Wunderlich, 1981	Qt East African copal
152. <i>Orchestina madagascariensis</i> Wunderlich, 2004as	Qt Madagascan copal
153. <i>Orchestina mortua</i> Petrunkevitch, 1971	Ne Chiapas amber
154. <i>Orchestina (Baltorchestina) multisetae</i> Wunderlich, 2008a	Pa Baltic amber
155. <i>Orchestina (Gallorchestina) parisiensis</i> Penney, 2007b	Pa Le Quesnoy amber
156. <i>Orchestina (Baltorchestina) perfecta</i> Wunderlich, 2008a	Pa Baltic amber
157. <i>Orchestina pusilla</i> (Menge in C. L. Koch & Berendt, 1854)	Pa Baltic amber
158. <i>Orchestina rabagensis</i> Saupe et al., 2012	K El Soplao amber
159. <i>Orchestina (Baltorchestina) rectangulata</i> Wunderlich, 2008a	Pa Baltic amber
160. <i>Orchestina (Baltorchestina) sternalis</i> Wunderlich, 2008a	Pa Baltic amber
161. <i>Orchestina tibialis</i> Wunderlich, 1988	Ne Dominican amber
162. <i>Orchestina truncata</i> Wunderlich, 2004at	Qt Colombian copal
163. <i>Orchestina tuberosa</i> Wunderlich, 1981	Pa Baltic amber
<i>Orchestina</i> sp. in Nishikawa (1974)	Qt Mizunami copal
<i>Orchestina</i> sp. in Saupe et al. (2012)	K Álava amber
<i>Orchestina</i> sp. in Soriano et al. (2010)	K San Just amber
<i>Orchestina</i> sp. in Wunderlich (2011h)	Pa Bitterfeld amber
Stenoonops Simon, 1891	Palaeogene – Recent
164. <i>Stenoonops incertus</i> (Wunderlich, 1988)	Ne Dominican amber
165. ? <i>Stenoonops rugosus</i> Wunderlich, 2004c	Pa Bitterfeld amber
166. <i>Stenoonops seldeni</i> (Penney, 2000)	Ne Dominican amber
ORSOLOBIDAE Cooke, 1965	Recent
no fossil record	
† PLUMORSOLIDAE Wunderlich, 2008d	Cretaceous
?Plumorsolidae indet. in Wunderlich (2008d)	K Myanmar amber
?Plumorsolidae indet. in Wunderlich (2011i)	K Myanmar amber
† Plumorsolus Wunderlich, 2008d	Cretaceous
167. <i>Plumorsolus gondwanensis</i> Wunderlich, 2008d	K Lebanese amber
ENTELEGYNAE Simon, 1893	Triassic – Recent
PALPIMANOIDEA Thorell, 1870a	Jurassic – Recent
family uncertain	
† Sinaranea Selden, Huang & Ren, 2008	Jurassic
168. <i>Sinaranea metaxyostraca</i> Selden, Huang & Ren, 2008*	J Daohugou, China
ARCHAEIDAE C. L. Koch & Berendt, 1854	Jurassic – Recent

Archaea C. L. Koch & Berendt, 1854	Palaeogene – Recent
169. ? <i>Archaea bitterfeldensis</i> Wunderlich, 2004 <i>d</i>	Pa Bitterfeld amber
170. <i>Archaea compacta</i> Wunderlich, 2004 <i>d</i>	Pa Baltic amber
171. <i>Archaea paradoxa</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
i. = <i>Archaea laevigata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
ii. = <i>Archaea incompta</i> Menge in C. L. Koch & Berendt, 1854	Pa Baltic amber
172. <i>Archaea pogneti</i> Simon, 1884 <i>b</i>	Pa Baltic amber
† Baltarchaea Eskov, 1992	Palaeogene
173. <i>Baltarchaea conica</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
† Burmesarchaea Wunderlich, 2008<i>d</i>	Cretaceous
174. <i>Burmesarchaea grimaldii</i> (Penney, 2003 <i>a</i>)	K Myanmar amber
† Eoarchaea Forster & Platnick, 1984	Palaeogene
175. <i>Eoarchaea hyperoptica</i> (Menge in C. L. Koch & Berendt, 1854)*	Pa Baltic amber
176. <i>Eoarchaea vidua</i> Wunderlich, 2004 <i>d</i>	Pa Baltic amber
† Eomysmauchenius Wunderlich, 2008<i>d</i>	Cretaceous
177. <i>Eomysmauchenius septentrionalis</i> Wunderlich, 2008 <i>d</i> *	K Myanmar amber
Eriauchenius O. P.-Cambridge, 1881	Quaternary – Recent
178. <i>Eriauchenius gracilicollis</i> (Millot, 1948) [Recent]	Qt Copal
i. = <i>Archaea copalensis</i> Lourenço, 2000 <i>b</i>	Qt Copal
† Filiauchenius Wunderlich, 2008<i>d</i>	Cretaceous
179. <i>Filiauchenius paucidentatus</i> Wunderlich, 2008 <i>d</i> *	K Myanmar amber
† Jurarchaea Eskov, 1987	Jurassic
180. <i>Jurarchaea zherikhini</i> Eskov, 1987*	J Kazakhstan
† Lacunauchenius Wunderlich, 2008<i>d</i>	Cretaceous
181. <i>Launauchenius speciosus</i> Wunderlich, 2008 <i>d</i> *	K Myanmar amber
† Myrmecarchaea Wunderlich, 2004<i>d</i>	Palaeogene
182. <i>Myrmecarchaea petiolus</i> Wunderlich, 2004 <i>d</i> *	Pa Baltic amber
183. <i>Myrmecarchaea pediculus</i> Wunderlich, 2004 <i>d</i>	Pa Baltic amber
† Patarchaea Selden, Huang & Ren, 2008	Jurassic
184. <i>Patarchaea muralis</i> Selden, Huang & Ren, 2008*	J Daohugou, China
† Saxonarchaea Wunderlich, 2004<i>d</i>	Palaeogene
185. <i>Saxonarchaea dentata</i> Wunderlich, 2004 <i>d</i> *	Pa Bitterfeld amber
186. <i>Saxonarchaea diabolica</i> Wunderlich, 2004 <i>d</i>	Pa Bitterfeld amber
MECY SMAUCHENIIDAE Simon, 1895	Cretaceous – Recent
† Archaemecys Saupe & Selden, 2009	Cretaceous
187. <i>Archaemecys arcantiensis</i> Saupe & Selden, 2009	K Charente amber
PARARCHAEIDAE Forster & Platnick, 1984	Recent
no fossil record	

HOLARCHAEIDAE Forster & Platnick, 1984	Recent
no fossil record	
MICROPHOLCOMMATIDAE Hickman, 1944	Palaeogene – Recent
† <i>Cenotextricella</i> Penney in Penney <i>et al.</i>, 2007	Palaeogene
188. <i>Cenotextricella simoni</i> Penney in Penney <i>et al.</i> , 2007	Pa Le Quesnoy amber
HUTTONIIDAE Simon, 1893	Cretaceous – Recent
unnamed genus and species in Penney & Selden (2006)	K Manitoban amber
STENOCHILIDAE Thorell, 1873	Recent
no fossil record	
† MICROPALPIMANIDAE Wunderlich, 2008d	Cretaceous
† <i>Micropalpimanus</i> Wunderlich, 2008d	Cretaceous
<i>Micropalpimanus</i> sp. indet in Wunderlich (2012d)	K Myanmar amber
189. <i>Micropalpimanus poinari</i> Wunderlich, 2008d	K Myanmar amber
PALPIMANIDAE Thorell, 1870a	Neogene – Recent
= OTITHOPOIDAE Thorell, 1869 [younger name protected by useage]	
= CHERSIDAE Canestrini & Pavesi, 1870	
<i>Otiothops</i> MacLeay, 1839	Neogene – Recent
<i>Otiothops</i> sp. 1–2 in Wunderlich (1988)	Ne Dominican amber
† LAGONOMEGOPIDAE Eskov & Wunderlich, 1995	Cretaceous
† <i>Archaelagonops</i> Wunderlich, 2012d	Cretaceous
190. <i>Archaelagonops salticoides</i> Wunderlich, 2012d*	K Myanmar amber
† <i>Burlagonomegops</i> Penney, 2005b	Cretaceous
191. <i>Burlagonomegops alavensis</i> Penney, 2006b	K Álava amber
192. <i>Burlagonomegops eskovi</i> Penney, 2005b*	K Myanmar amber
† <i>Lagonoburmops</i> Wunderlich, 2012d	Cretaceous
193. <i>Lagonoburmops plumosus</i> Wunderlich, 2012d*	K Myanmar amber
† <i>Lagonomegops</i> Eskov & Wunderlich, 1995	Cretaceous
194. <i>Lagonomegops americanus</i> Penney, 2005b	K New Jersey amber
195. <i>Lagonomegops sukatchevae</i> Eskov & Wunderlich, 1995*	K Taimyr amber
† <i>Myanlagonops</i> Wunderlich, 2012d	Cretaceous
196. <i>Myanlagonops gracilipes</i> Wunderlich, 2012d*	K Myanmar amber
† <i>Zarquagonomegops</i> Kaddumi, 2007	Cretaceous
197. <i>Zarquagonomegops wunderlichi</i> Kaddumi, 2007*	K Jordanian amber
† GRANDOCULIDAE Penney, 2011	Cretaceous

NB: The validity of this family has been challenged (cf. Wunderlich 2012d; Pérez-de la Fuente *et al.* 2013).

- † **Grandoculus Penney, 2004b** **Cretaceous**
 198. *Grandoculus chemahawinensis* Penney, 2004b* K Manitobian amber
- † **SPATIATORIDAE Petrunkevitch, 1942** **Palaeogene**
 † **Spatiator Petrunkevitch, 1942** **Palaeogene**
 199. *Spatiator caulis* Wunderlich, 2008a Pa Baltic amber
 200. *Spatiator martensi* Wunderlich, 2006 Pa Baltic amber
 201. *Spatiator praeceps* Petrunkevitch, 1942* Pa Baltic amber
Spatiator sp. in Wunderlich (2011h) Pa Baltic amber
- MALKARIDAE Davies, 1980** **Recent**
 = STERNODIDAE Moran, 1986
 no fossil record
- MIMETIDAE Simon, 1881** **Palaeogene – Recent**
 = CTENOPHORIDAE Blackwall, 1870 [younger name protected by useage]
 Mimetidae gen. et sp. indet. in Penney *et al.* (2012a) Pa Indian amber
 Mimetini sp. 1–4 in Wunderlich (2004q) Pa Baltic amber
- Ero C. L. Koch, 1836** **Palaeogene – Recent**
 = † *Palaeoero* Wunderlich, 2004q
 = † *Succinero* Wunderlich, 2004q
 [Wunderlich revalidated both as putative subgenera]
 202. *Ero carboneana* Petrunkevitch, 1942 Pa Baltic amber
 203. *Ero aberrans* Petrunkevitch, 1958 Pa Baltic amber
 [Treated as a *nomen dubium* by Harms & Dunlop (2009)]
 204. *Ero (Succinero) clunis* Wunderlich, 2012c Pa Baltic amber
 205. *Ero (Succinero) gracilitibialis* Wunderlich, 2012c Pa Baltic amber
 206. *Ero (Paleoero) longitarsus* (Wunderlich, 2004q) Pa Baltic amber
 207. *Ero permunda* Petrunkevitch, 1942 Pa Baltic amber
 208. *Ero (Succinero) rovnoensis* (Wunderlich, 2004ar) Pa Rovno amber
 209. *Ero (Succinero) veta* Wunderlich, 2012c Pa Baltic amber
- Mimetus Hentz, 1832** **Palaeogene – Recent**
 ? *Mimetus* sp. in Wunderlich (1988) Ne Dominican amber
 210. *Mimetus bituberculatus* Wunderlich, 1988 Ne Dominican amber
 211. *Mimetus brevipes* Wunderlich, 2004q Pa Baltic amber
 [synonymised by Harms & Dunlop (2009), but resurrected by Wunderlich (2012c)]
 212. ? *Mimetus longipes* Wunderlich, 2004q Pa Baltic amber
- † **Protomimetus Wunderlich, 2011** **Palaeogene**
 213. ? *Protomimetus breviclypeus* Wunderlich, 2011h Pa Baltic amber
 214. *Protomimetus longicypeus* Wunderlich, 2011h* Pa Baltic amber
- ERESOIDEA C. L. Koch, 1851** **Cretaceous – Recent**

- ERESIDAE C. L. Koch, 1851** **?Miocene – Recent**
 no body fossil record, but a web attributed to the extant genus *Seothyra* was described by Pickford (2000) from Miocene aeolianites in the Namib Desert of Namibia
- ‘OECOBIOIDEA’**
- Oecobioidea fam. indet. *in* Wunderlich (2008d) K Myanmar amber
- OECOBIIDAE Blackwall, 1862** **Cretaceous – Recent**
 = UROCTEIDAE Thorell, 1869
- † **Lebanoecobius Wunderlich, 2004e** **Cretaceous**
215. *Lebanoecobius schleei* Wunderlich, 2004e* K Lebanese amber
- † **Mizalia C. L. Koch & Berendt, 1854** **Palaeogene**
 = † *Paruroctea* Petrunkevitch, 1942
216. *Mizalia blauvelti* (Petrunkevitch, 1942) Pa Baltic amber
217. *Mizalia gemini* Wunderlich, 2004e Pa Baltic amber
218. *Mizalia rostrata* C. L. Koch & Berendt, 1854* Pa Baltic amber
 i. = *Mizalia pilosula* C. L. Koch & Berendt, 1854 Pa Baltic amber
219. *Mizalia spirembolus* Wunderlich, 2004e Pa Baltic amber
Mizalia sp. *in* Wunderlich (2011h) Pa Baltic/Bltter. amber
- Oecobius Lucas, 1846** **?Cretaceous – Recent**
220. *Oecobius piliformis* Wunderlich, 1988 Ne Dominican amber
 ?*Oecobius* sp. indet *in* Penney (2002) K New Jersey amber
- Uroctea Dufour, 1820** **Palaeogene – Recent**
221. *Uroctea galloprovincialis* Gourret, 1887 Pa Aix-en-Provence
- † **Zamilia Wunderlich, 2008d** **Cretaceous**
222. *Zamilia antecessor* Wunderlich, 2008d K Myanmar amber
- HERSILIIDAE Thorell, 1870a** **Cretaceous – Recent**
 = CHALINUROIDAE Thorell, 1873
- Hersiliidae sp. 1–3 *in* Wunderlich (2004d) Pa Baltic amber
- Hersiliidae sp. *in* Wunderlich (2011f) Qt Madagascar copal
- † **Burmesiola Wunderlich, 2011i** **Cretaceous**
223. *Burmesiola cretacea* Wunderlich, 2011i* K Myanmar amber
- † **“Fictotama Petrunkevitch, 1963 (nomen dubium)”** **Neogene**
 [Wunderlich 2011f placed a new species in this genus, which was previously considered a *nomen dubium*. He did not formally revalidate the genus]
224. *“Fictotama” maculosa* Wunderlich, 2011g Ne Dominican amber
- † **Gerdia Menge, 1869** **Palaeogene**
225. *Gerdia myura* Menge, 1869* Pa Baltic amber
- † **Gerdiopsis Wunderlich, 2004e** **Palaeogene**
226. *Gerdiopsis infrigens* Wunderlich, 2004e* Pa Baltic amber
- † **Gerdiorum Wunderlich 2004e** **Palaeogene**

227. <i>Gerdiorum inflexum</i> Wunderlich 2004e*	Pa Baltic amber
Hersilia Audouin, 1826	Palaeogene – Recent
= † <i>Hersiliopsis</i> Wunderlich, 2004e	
228. <i>Hersilia aquisextana</i> Gourret, 1887	Pa Aix-en-Provence
229. <i>Hersilia longipes</i> Giebel, 1856	Pa Baltic amber
230. <i>Hersilia madagascarensis</i> (Wunderlich, 2004e)	Qt–R Madagas. copal
231. ? <i>Hersilia miranda</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† Hersiliana Wunderlich, 2004e	Quaternary – Recent
232. <i>Hersiliana brevipes</i> Wunderlich, 2004e*	Qt Madagascan copal
† Prototama Petrunkevitch, 1971	Neogene
= † <i>Priscotama</i> Petrunkevitch, 1971	
233. <i>Prototama antiqua</i> (Petrunkevitch, 1971)	Ne Chiapas amber
234. <i>Prototama maior</i> (Wunderlich, 1988)	Ne Dominican amber
235. <i>Prototama media</i> (Wunderlich, 1988)	Ne Dominican amber
236. <i>Prototama minor</i> (Wunderlich, 1987)	Ne Dominican amber
237. <i>Prototama succinea</i> Petrunkevitch, 1971*	Ne Chiapas amber
<i>Prototama</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
Superfamily uncertain	
† BURMASCUTIDAE Wunderlich, 2008d	Cretaceous
† <i>Burmascutum</i> Wunderlich, 2008d	Cretaceous
238. <i>Burmascutum aenigma</i> Wunderlich, 2008d*	K Myanmar amber
† SALTICOIDIDAE Wunderlich, 2008d	Cretaceous
† <i>Salticoidus</i> Wunderlich, 2008d	Cretaceous
239. <i>Salticoidus kaddumiorum</i> Wunderlich, 2008d*	K Jordanian amber
'CANOE TAPETUM' CLADE	Triassic – Recent
ORBICULARIAE Walckenaer, 1802	Triassic – Recent
DEINOPOIDEA C. L. Koch, 1851	Cretaceous – Recent
DEINOPIDAE C. L. Koch, 1851	Cretaceous – Recent
<i>Deinopsis</i> MacLeay, 1839	Quaternary – Recent
240. <i>Deinopsis</i> ? <i>madagascariensis</i> Lenz, 1886 [Recent]	Qt Madagascar copal
Menneus Simon, 1876b	Palaeogene – Recent
241. ? <i>Menneus pietrzeniukae</i> Wunderlich, 2004g	Pa Baltic amber
? <i>Menneus</i> sp. 1–3 <i>in</i> Wunderlich (2004g)	Pa Baltic amber
† <i>Palaeomicromennus</i> Penney, 2003b	Cretaceous
242. <i>Palaeomicromennus lebanensis</i> Penney, 2003b*	K Lebanese amber
ULOBORIDAE Thorell, 1869	Cretaceous – Recent
Uloboridae indet. <i>in</i> Wunderlich (2011f)	Qt Madagascar copal
† <i>Burmuloborus</i> Wunderlich, 2008d	Cretaceous

243. *Burmuloborus parvus* Wunderlich, 2008d* K Myanmar amber
- † **Eomiagrammopes Wunderlich, 2004f** **Palaeogene**
244. *Eomiagrammopes maior* Wunderlich, 2004f Pa Baltic amber
245. *Eomiagrammopes minor* Wunderlich, 2004f Pa Baltic amber
246. *Eomiagrammopes semiapertus* Wunderlich, 2011h Pa Baltic amber
247. *Eomiagrammopes singularis* Wunderlich, 2004f* Pa Baltic amber
248. *Eomiagrammopes spinipes* Wunderlich, 2004f Pa Baltic amber
- Eomiagrammopes* sp. 1–2 in Wunderlich (2004f) Pa Baltic amber
- ?*Eomiagrammopes* sp. in Wunderlich (2004f) Pa Baltic amber
- † **Hyptiomopes Wunderlich, 2004f** **Palaeogene**
249. *Hyptiomopes bitterfeldensis* Wunderlich 2004f* Pa Bitterfeld amber
- ?*Hyptiomopes* sp. in Wunderlich (2004f) Pa Bitterfeld amber
- Hyptiotes Walckenaer, 1837** **Palaeogene – Recent**
- = † *Androgeus* C. L. Koch & Berendt, 1854
250. *Hyptiotes convexus* Wunderlich, 2004f Pa Baltic amber
251. *Hyptiotes glaber* Wunderlich, 2004f Pa Baltic amber
252. *Hyptiotes saetosus* Wunderlich, 2004f Pa Baltic amber
253. *Hyptiotes stellatus* Wunderlich, 2004f Pa Baltic amber
254. *Hyptiotes triqueter* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- † **Jerseyuloborus Wunderlich, 2011i** **Cretaceous**
255. *Jerseyuloborus longisoma* Wunderlich, 2011i* K New Jersey amber
- Miagrammopes O. P.-Cambridge, 1870** **Neogene – Recent**
256. *Miagrammopes dominicanus* Wunderlich, 2004e Ne Dominican amber
- Miagrammopes* sp. in Penney (2001) Ne Dominican amber
- Miagrammopes* sp. in Wunderlich (2011f) Qt Madagascar copal
- † **Ocululoborus Wunderlich, 2012d** **Cretaceous**
257. *Ocululoborus curvatus* Wunderlich, 2012d* K Myanmar amber
- † **Opellianus Wunderlich, 2004f** **Palaeogene**
258. *Opellianus excellens* Wunderlich, 2004f* Pa Baltic amber
259. *Opellianus kazimierasi* Wunderlich 2004f Pa Baltic amber
260. *Opellianus ludwigi* Wunderlich 2004f Pa Baltic amber
- † **Palaeomiagrammopes Wunderlich, 2008d** **Cretaceous**
261. *Palaeomiagrammopes vesica* Wunderlich, 2008d* K Myanmar amber
- † **Palaeouloborus Selden, 1990** **Cretaceous**
262. *Palaeouloborus lacasae* Selden, 1990* K Sierra de Montsech
- † **Paramiagrammopes Wunderlich, 2008d** **Cretaceous**
263. *Paramiagrammopes cretaceus* Wunderlich, 2008d* K Myanmar amber
- Paramiagrammopes* sp. in Wunderlich (2008d) K Myanmar amber
- † **Ulobomopes Wunderlich, 2004f** **Palaeogene**
264. *Ulobomopes unicus* Wunderlich, 2004f* Pa Baltic amber

ARANEOIDEA Latreille, 1806	Jurassic – Recent
Araneoidea fam indet. <i>in</i> Wunderlich (2008 <i>d</i>)	K Myanmar amber
† Mesarania Hong, 1984	Jurassic
265. <i>Mesarania hebeiensis</i> Hong, 1984*	J Hebei, China
CYATHOLIPIDAE Simon, 1894	Palaeogene – Recent
= TEEMENAARIDAE Davies, 1978	
† Balticolipus Wunderlich, 2004<i>m</i>	Palaeogene
266. <i>Balticolipus kruemmeri</i> Wunderlich, 2004 <i>m</i> *	Pa Baltic / Bitt. amber
† Cyathosuccinus Wunderlich, 2004<i>m</i>	Palaeogene
267. <i>Cyathosuccinus elongatus</i> Wunderlich, 2004 <i>m</i> *	Pa Baltic amber
† Erigolipus Wunderlich, 2004<i>m</i>	Palaeogene
268. <i>Erigolipus griswoldi</i> Wunderlich, 2004 <i>m</i> *	Pa Baltic amber
† Spinilipus Wunderlich, 1993<i>b</i>	Palaeogene
269. <i>Spinilipus bispinosus</i> Wunderlich, 2004 <i>m</i>	Pa Bitterfeld amber
270. <i>Spinilipus curvatus</i> Wunderlich, 2004 <i>m</i>	Pa Bitterfeld amber
271. <i>Spinilipus glinki</i> Wunderlich, 2004 <i>m</i>	Pa Baltic amber
272. <i>Spinilipus kerneggeri</i> Wunderlich, 1993 <i>b</i> *	Pa Baltic amber
273. <i>Spinilipus longembolus</i> Wunderlich, 2004 <i>m</i>	Pa Baltic amber
† Succinilipus Wunderlich, 1993<i>b</i>	Palaeogene
274. <i>Succinilipus abditus</i> Wunderlich, 2004 <i>m</i>	Pa Baltic / Bitt. amber
275. <i>Succinilipus aspinosus</i> Wunderlich, 2004 <i>m</i>	Pa Bitterfeld amber
276. <i>Succinilipus saxoniensis</i> Wunderlich, 1993 <i>b</i>	Pa Bitterfeld amber
277. <i>Succinilipus similis</i> Wunderlich, 2004 <i>m</i>	Pa Bitterfeld amber
278. <i>Succinilipus teuberi</i> Wunderlich, 1993 <i>b</i> *	Pa Baltic amber
<i>Succinilipus</i> sp. <i>in</i> Wunderlich (2004 <i>m</i>)	Pa Baltic / Bitt. amber
SYNOTAXIDAE Simon, 1894	Palaeogene – Recent
† Acrometa Petrunkevitch, 1942	Palaeogene
= † <i>Eogonatium</i> Petrunkevitch, 1942	
= † <i>Liticen</i> Petrunkevitch, 1942	
= † <i>Theridiometa</i> Petrunkevitch, 1942	
= † <i>Viocurus</i> Petrunkevitch, 1958	
279. <i>Acrometa clava</i> Wunderlich, 2004 <i>n</i>	Pa Baltic amber
280. <i>Acrometa cristata</i> Petrunkevitch, 1942*	Pa NE Europe ambers
i. = <i>Theridiometa edwardsi</i> Petrunkevitch, 1942	Pa Baltic amber
ii. = <i>Viocurus fossilis</i> Petrunkevitch, 1958	Pa Baltic amber
281. <i>Acrometa eichmanni</i> Wunderlich, 2004 <i>n</i>	Pa Baltic amber
282. <i>Acrometa incidens</i> Wunderlich, 2004 <i>n</i>	Pa Baltic amber
283. <i>Acrometa minutum</i> (Petrunkevitch, 1942)	Pa Baltic amber

284. *Acrometa pala* Wunderlich, 2004*n* Pa Baltic amber
285. *Acrometa robusta* (Petrunkevitch, 1942) Pa Baltic amber
286. *Acrometa pseudorobusta* Dunlop & Jekel, 2009 Pa Baltic amber
 i. = *Acrometa robusta* (Petrunkevitch, 1946) [preoccupied]
287. *Acrometa samlandica* (Petrunkevitch, 1942) Pa Baltic amber
288. *Acrometa setosus* (Petrunkevitch, 1942) Pa Baltic amber
289. *Acrometa succini* Petrunkevitch, 1942 Pa Baltic amber
- † **Anandrus Menge, 1856** **Palaeogene**
 = † *Elucus* Petrunkevitch, 1942
290. *Anandrus inermis* (Petrunkevitch, 1942) Pa Baltic amber
291. *Anandrus infelix* (Petrunkevitch, 1950)* Pa Baltic amber
292. *Anandrus quaesitus* (Petrunkevitch, 1958) Pa Baltic amber
293. *Anandrus redemptus* (Petrunkevitch, 1958) Pa Baltic amber
- † **Chelicerinus Wunderlich, 2008a** **Palaeogene**
294. *Chelicerinus abnormis* Wunderlich, 2008a Pa Bitterfeld amber
- † **Cornuanandrus Wunderlich, 1986** **Palaeogene**
295. *Cornuanandrus bifurcatus* Wunderlich, 2004*n* Pa Bitterfeld amber
296. *Cornuanandrus bitterfeldensis* Wunderlich, 2004*n* Pa Bitterfeld amber
297. *Cornuanandrus corniculans* Wunderlich, 2004*n* Pa Baltic amber
298. *Cornuanandrus maior* Wunderlich, 1986* Pa Baltic amber
299. *Cornuanandrus minor* Wunderlich, 2004*n* Pa Baltic amber
- † **Dubiosynotaxus Wunderlich, 2004n** **Palaeogene**
300. *Dubiosynotaxus perfectus* Wunderlich, 2004*n** Pa Baltic amber
- † **Eosynotaxus Wunderlich, 2004n** **Palaeogene**
301. *Eosynotaxus bispinosus* Wunderlich, 2004*n* Pa Baltic amber
302. *Eosynotaxus bitterfeldensis* Wunderlich, 2004*n* Pa Bitterfeld amber
303. *Eosynotaxus custodens* Wunderlich, 2004*n* Pa Baltic amber
304. *Eosynotaxus fastigatus* Wunderlich, 2004*n* Pa Baltic amber
305. *Eosynotaxus paucispina* Wunderlich, 2004*n* Pa Baltic amber
306. *Eosynotaxus spinipes* Wunderlich, 2004*n* Pa Baltic amber
307. *Eosynotaxus wegneri* Wunderlich, 2004*n** Pa Baltic amber
- † **Gibbersynotaxus Wunderlich, 2004n** **Palaeogene**
308. *Gibbersynotaxus parvus* Wunderlich, 2004*n** Pa Baltic amber
- † **Protophysoglenes Wunderlich, 2004n** **Palaeogene**
309. *Protophysoglenes impressum* Wunderlich, 2004*n** Pa Baltic amber
- † **Pseudoacrometa Wunderlich, 1986** **Palaeogene**
310. *Pseudoacrometa gracilipes* Wunderlich, 1986* Pa Baltic amber
311. *Pseudoacrometa wittmanni* Wunderlich, 2004*n* Pa Baltic amber
- † **Succinitaxus Wunderlich, 2004n** **Palaeogene**
312. *Succinitaxus brevis* Wunderlich, 2004*n** Pa Baltic, Bitterfeld &
 Rovno amber

313. ? <i>Succinitaxus minutus</i> Wunderlich, 2004n	Pa Baltic amber
† <i>Sulcosynotaxus</i> Wunderlich, 2004n	Palaeogene
314. <i>Sulcosynotaxus cavatus</i> Wunderlich, 2004n*	Pa Baltic amber
NESTICIDAE Simon, 1894	Palaeogene – Recent
† <i>Balticonesticus</i> Wunderlich, 1986	Palaeogene
315. <i>Balticonesticus flexuosus</i> Wunderlich, 1986*	Pa Baltic amber
<i>Eidmanella</i> Roewer, 1935	Quaternary
316. <i>Eidmanella pallida</i> (Emerton, 1875) [Recent]	Qt Madagascar copal
† <i>Eopopino</i> Petrunkevitch, 1942	Palaeogene
317. <i>Eopopino budrysi</i> Eskov & Marusik, 1992	Pa Baltic amber
318. <i>Eopopino inopinatus affinis</i> Wunderlich, 1986	Pa Baltic amber
319. <i>Eopopino inopinatus inopinatus</i> Wunderlich, 1986	Pa Baltic amber
320. <i>Eopopino longipes</i> Petrunkevitch, 1942*	Pa Baltic amber
321. <i>Eopopino palanga</i> Eskov & Marusik, 1992	Pa Baltic amber
322. <i>Eopopino rarus rarus</i> Wunderlich, 1986	Pa Baltic amber
323. <i>Eopopino rarus solitarius</i> Wunderlich, 1986	Pa Baltic amber
324. <i>Eopopino rudloffii</i> Wunderlich, 2004o	Pa Bitterfeld amber
<i>Eopopino</i> sp. in Wunderlich (1986)	Pa Bitterfeld amber
† <i>Heteronesticus</i> Wunderlich, 1986	Palaeogene
325. <i>Heteronesticus magnoparacymbialis</i> Wunderlich, 1986*	Pa Baltic amber
† <i>Hispanonesticus</i> Wunderlich, 1986	Neogene
326. <i>Hispanonesticus latopalpus</i> Wunderlich, 1986*	Ne Dominican amber
THERIDIIDAE Sundevall, 1833	?Cretaceous – Recent
= PHYCOIDAE Thorell, 1873	
= EPISINIDAE O. P.-Cambridge, 1879a	
= HADROTARSIDAE Thorell, 1881	
?Theridiidae gen. et sp. indet in McAlpine & Martin (1969)	K Canadian amber
Theridiidae gen. et sp. in Nishikawa (1974)	Qt Mizunami copal
<i>Achaearana</i> Strand, 1929	Neogene – Recent
327. <i>Achaearana extincta</i> Wunderlich, 1988	Ne Dominican amber
<i>Achaearana</i> sp. in Wunderlich (1988)	Ne Dominican amber
<i>Argyrodes</i> Simon, 1864	Neogene – Recent
328. <i>Argyrodes (Ariamnes) copalis</i> Wunderlich, 2008b	Qt Colombian copal
329. <i>Argyrodes (Ariamnes) resina</i> Wunderlich, 2011f	Qt Madagascar copal
330. <i>Argyrodes (Rhomphaea) gibbifera</i> Wunderlich, 2004as	Qt Madagascar copal
331. <i>Argyrodes parvipatellaris</i> Wunderlich, 1988	Ne Dominican amber
<i>Argyrodes</i> sp. in Wunderlich (1988)	Ne Dominican amber
† <i>Balticoridion</i> Wunderlich, 2008b	Palaeogene
332. <i>Balticoridion dubium</i> Wunderlich, 2008b*	Pa Baltic / Bitt. amber
† <i>Balticpholcomma</i> Wunderlich, 2008b	Palaeogene

333. <i>Balticpholcomma scutatum</i> Wunderlich, 2008b*	Pa Baltic amber
† Caudasinus Wunderlich, 2008b	Palaeogene
334. <i>Caudasinus bispinosus</i> Wunderlich, 2008b	Pa Baltic amber
335. <i>Caudasinus caudatus</i> Wunderlich, 2008b*	Pa Baltic amber
336. <i>Caudasinus regeneratus</i> Wunderlich, 2008b	Pa Baltic amber
<i>Caudasinus</i> sp. in Wunderlich (2008b)	Pa Baltic amber
Chrosiothes Simon, 1894	Neogene – Recent
337. <i>Chrosiothes biconigerus</i> Wunderlich, 1988	Ne Dominican amber
338. <i>Chrosiothes curvispinosus</i> Wunderlich, 1988	Ne Dominican amber
339. <i>Chrosiothes emulgatus</i> Wunderlich, 1988	Ne Dominican amber
340. <i>Chrosiothes longispinosus</i> Wunderlich, 1988	Ne Dominican amber
341. <i>Chrosiothes monoceros</i> Wunderlich, 1988	Ne Dominican amber
342. <i>Chrosiothes tumulus</i> Wunderlich, 1988	Ne Dominican amber
343. <i>Chrosiothes unicornis</i> Wunderlich, 1988	Ne Dominican amber
Chryso O. P.-Cambridge, 1882a	Neogene – Recent
344. <i>Chryso conspicua</i> Wunderlich, 1988	Ne Dominican amber
345. <i>Chryso dubia</i> Wunderlich, 1988	Ne Dominican amber
† Clavibertus Wunderlich, 2008b	Palaeogene
346. <i>Clavibertus parvus</i> Wunderlich, 2008b	Pa Baltic amber
347. <i>Clavibertus prominens</i> Wunderlich, 2008b*	Pa Baltic amber
† Clya C. L. Koch & Berendt, 1854	Palaeogene
348. <i>Clya abdita</i> Wunderlich, 2008b	Pa Baltic amber
349. <i>Clya lugubris</i> C. L. Koch & Berendt, 1854*	Pa Baltic / Rovno amber
350. <i>Clya calefacta</i> Wunderlich, 2008b	Pa Baltic amber
351. <i>Clya gracilis</i> (Petrunkevitch, 1958)	Pa Baltic amber
352. <i>Clya granulata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
353. <i>Clya obscura</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
354. <i>Clya rotata</i> Wunderlich, 2008b	Pa Baltic amber
355. <i>Clya supercalefacta</i> Wunderlich, 2008b	Pa Baltic amber
356. <i>Clya superspiralis</i> Wunderlich, 2008b	Pa Baltic amber
357. <i>Clya tricurvata</i> Wunderlich, 2008b	Pa Baltic amber
† Cornutidion Wunderlich, 1988	Neogene
358. <i>Cornutidion elongatum</i> Wunderlich, 1988*	Ne Dominican amber
Craspedisia Simon, 1894	Neogene – Recent
359. <i>Craspedisia yapchoontecki</i> Penney & Marusik in Penney <i>et al.</i> (2012b)	Ne Dominican amber
† Cymbiopholcomma Wunderlich, 2008b	Palaeogene
360. <i>Cymbiopholcomma dudum</i> Wunderlich, 2008b*	Pa Baltic amber
361. <i>Cymbiopholcomma spiculum</i> Wunderlich, 2008b	Pa Baltic amber
† Dipoenata Wunderlich, 1988	Neogene
362. <i>Dipoenata altiocolata</i> Wunderlich, 1988	Ne Dominican amber

363.	<i>Dipoenata cala</i> Wunderlich, 1988	Ne Dominican amber
364.	<i>Dipoenata clypeata</i> Wunderlich, 1988	Ne Dominican amber
365.	<i>Dipoenata globulus</i> Wunderlich, 1988	Ne Dominican amber
366.	<i>Dipoenata praedominicana</i> (Wunderlich, 1986)	Qt Dominican copal
367.	<i>Dipoenata stipes</i> Wunderlich, 1988*	Ne Dominican amber
368.	<i>Dipoenata yolandae</i> Wunderlich, 1988	Ne Dominican amber
	<i>Dipoenata</i> sp. in Wunderlich (1988)	Ne Dominican amber
†	Eoasagena Wunderlich, 2008b	Palaeogene
	369. <i>Eoasagena scutata</i> Wunderlich, 2008b*	Pa Baltic amber
†	Eolyrifer Wunderlich, 2008b	Palaeogene
	370. <i>Eolyrifer longitibialis</i> Wunderlich, 2008b*	Pa Baltic amber
†	Eomysmena Petrunkevitch, 1942	Palaeogene – Neogene
	= † <i>Antopia</i> Menge, 1854 [tentative synonymy]	
	= † <i>Astodipoena</i> Petrunkevitch, 1958	
	= † <i>Eodipoena</i> Petrunkevitch, 1942	
371.	<i>Eomysmena asta</i> Petrunkevitch, 1971	Ne Chiapas amber
372.	<i>Eomysmena aviceps</i> Wunderlich, 2008b	Pa Baltic amber
373.	<i>Eomysmena calefacta</i> Wunderlich, 2008b	Pa Baltic amber
374.	<i>Eomysmena crassa</i> (Petrunkevitch, 1958)	Pa Baltic amber
375.	<i>Eomysmena baltica</i> Petrunkevitch, 1946	Pa Baltic amber
376.	' <i>Eomysmena</i> ' <i>bassleri</i> (Petrunkevitch, 1942)	Pa Baltic amber
377.	? <i>Eomysmena kaestneri</i> (Petrunkevitch, 1958)	Pa Baltic amber
378.	<i>Eomysmena militaris</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
379.	<i>Eomysmena moritura</i> Petrunkevitch, 1942*	Pa Baltic amber
	i. = <i>Eomysmena consulta</i> (Petrunkevitch, 1958)	
	[tentative synonymy]	Pa Baltic amber
380.	<i>Eomysmena nielseni</i> (Petrunkevitch, 1958)	Pa Baltic amber
381.	<i>Eomysmena oculata</i> (Petrunkevitch, 1942)	Pa Baltic amber
382.	<i>Eomysmena punctulata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
383.	<i>Eomysmena recta</i> Wunderlich, 2008b	Pa Baltic amber
384.	<i>Eomysmena tenera</i> (Menge in C. L. Koch & Berendt, 1854)	Pa Baltic amber
	<i>Eomysmena</i> spp. in Wunderlich 2008b	Pa Baltic / Bitt. Amber
†	Eoteutana Wunderlich, 2008b	Palaeogene
	385. <i>Eoteutana hirsuta</i> Wunderlich, 2008b*	Pa Baltic amber
	Episinus Latreille, 1809	Palaeogene – Recent
	= † <i>Flegia</i> C. L. Koch & Berendt, 1854	
	= † <i>Impulsor</i> Petrunkevitch, 1942	
	= † <i>Malleator</i> Petrunkevitch, 1942	
	= † <i>Mictodipoena</i> Petrunkevitch, 1958	
	= † <i>Municeps</i> Petrunkevitch, 1942 [tentative synonymy]	
386.	<i>Episinus anapidaeque</i> Wunderlich, 2008b	Pa Baltic amber
387.	<i>Episinus antecognatus</i> Wunderlich, 1986	Qt Dominican copal

388.	<i>Episinus appendix</i> Wunderlich, 2008b	Pa	Baltic amber
389.	<i>Episinus arrodens</i> Wunderlich, 2008b	Pa	Baltic amber
390.	<i>Episinus balticus</i> Marusik & Penney, 2004	Pa	Baltic / Bitt. amber
391.	<i>Episinus brevipalpus</i> Wunderlich, 1988	Ne	Dominican amber
392.	<i>Episinus bulla</i> Wunderlich, 2008b	Pa	Baltic amber
393.	<i>Episinus chiapasanus</i> (Petrunkevitch, 1971)	Ne	Chiapas amber
394.	<i>Episinus clunis</i> Wunderlich, 2008b	Pa	Baltic amber
395.	<i>Episinus cochlear</i> Wunderlich, 2008b	Pa	Baltic amber
396.	<i>Episinus cornutus</i> Wunderlich, 1988	Ne	Dominican amber
397.	<i>Episinus cymbialis</i> Wunderlich, 2008b	Pa	Baltic amber
398.	<i>Episinus dimidius</i> Wunderlich, 2008b	Pa	Baltic amber
399.	<i>Episinus eskovi</i> Marusik & Penney, 2004	Pa	Baltic amber
400.	<i>Episinus isopteraque</i> Wunderlich, 2008b	Pa	Baltic amber
401.	<i>Episinus latus</i> Wunderlich, 2008b	Pa	Baltic amber
402.	<i>Episinus longimanus</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
	i. = <i>Malleator niger</i> Petrunkevitch, 1942	Pa	Baltic amber
403.	<i>Episinus longisoma</i> Wunderlich, 2008b	Pa	Baltic amber
404.	<i>Episinus minutus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
405.	<i>Episinus mordellidaeque</i> Wunderlich, 2008b	Pa	Baltic amber
406.	<i>Episinus musculus</i> Wunderlich, 2008b	Pa	Baltic amber
407.	<i>Episinus mutilus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
408.	<i>Episinus nausticymbium</i> Wunderlich, 2008b	Pa	Baltic amber
409.	<i>Episinus neglectus</i> (Petrunkevitch, 1942)	Pa	Baltic amber
410.	<i>Episinus penneyi</i> Garcia-Villafuerte, 2006a	Ne	Chiapas amber
411.	<i>Episinus praecognatus</i> Wunderlich, 1982	Ne	Dominican amber
412.	<i>Episinus pulcher</i> (Petrunkevitch, 1942)	Pa	Baltic amber
413.	<i>Episinus regalis</i> (Petrunkevitch, 1958)	Pa	Baltic amber
414.	<i>Episinus stridulus</i> (Petrunkevitch, 1958)	Pa	Baltic amber
415.	<i>Episinus tibiassetta</i> Wunderlich, 2011g	Ne	Dominican amber
416.	<i>Episinus transversus</i> Wunderlich, 2008b	Pa	Baltic amber
417.	<i>Episinus tuberosus</i> Wunderlich, 1988	Ne	Dominican amber
	<i>Episinus</i> spp. in Wunderlich (2008b)	Pa	Baltic amber
	<i>Euryopsis</i> Menge, 1868		Palaeogene – Recent
418.	? <i>Euryopsis araneoides</i> Wunderlich, 2008b	Pa	Baltic amber
419.	<i>Euryopsis bitterfeldensis</i> Wunderlich, 2008b	Pa	Baltic / Bitt. amber
420.	<i>Euryopsis nexus</i> Wunderlich, 2008b	Pa	Baltic amber
421.	<i>Euryopsis streyi</i> Wunderlich, 2008b	Pa	Baltic / Bitt. Amber
	<i>Euryopsis/Emertonella</i> complex in Penney <i>et al.</i> (2012c)	Qt	Colombian copal
†	<i>Euryopus</i> Menge in C. L. Koch & Berendt, 1854		Palaeogene
422.	<i>Euryopus gracilipes</i> Menge in C. L. Koch & Berendt, 1854*	Pa	Baltic amber
	<i>Faiditus</i> Keyserling, 1884		Neogene – Recent

423. <i>Faiditus crassipatellaris</i> (Wunderlich, 1988)	Ne Dominican amber
† Femurraptor Wunderlich, 2011g	Neogene
424. <i>Femurraptor dominicanus</i> Wunderlich, 2011g*	Ne Dominican amber
† Globulidion Wunderlich, 2008b	Palaeogene
425. <i>Globulidion cochlea</i> Wunderlich, 2008b*	Pa Baltic amber
† Hirsutipalpus Wunderlich, 2008b	Palaeogene
426. <i>Hirsutipalpus varipes</i> Wunderlich, 2008b*	Pa Baltic / Bitt. Amber
† Kochiuridion Wunderlich, 2008b	Palaeogene
427. <i>Kochiuridion scutatum</i> Wunderlich, 2008b*	Pa Baltic / Bitt. amber
Lasaeola Simon, 1881	Palaeogene – Recent
= † <i>Nactodipoena</i> Petrunkevitch, 1942 [a subgenus <i>in</i> Wunderlich (2008b)]	
428. <i>Lasaeola acumen</i> Wunderlich, 2008b	Pa Baltic amber
429. <i>Lasaeola baltica</i> (Marusik & Penney, 2004)	Pa Baltic amber
430. <i>Lasaeola bitterfeldensis</i> Wunderlich, 2008b	Pa Bitterfeld amber
431. <i>Lasaeola communis</i> Wunderlich, 2008b	Pa Baltic amber
432. <i>Lasaeola (Nactodipoena) dunbari</i> (Petrunkevitch, 1942)	Pa Baltic amber
433. ? <i>Lasaeola furca</i> Wunderlich, 2008b	Pa Baltic amber
434. <i>Lasaeola germanica</i> (Petrunkevitch, 1958)	Pa Baltic amber
435. <i>Lasaeola (Phycosoma) inclinata</i> Wunderlich, 2012a	Qt Madagascan copal
436. <i>Lasaeola infulata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic / Bitt. Amber
437. <i>Lasaeola larvaque</i> Wunderlich, 2008b	Pa Baltic amber
438. <i>Lasaeola latusulci</i> Wunderlich, 2008b	Pa Baltic amber
439. <i>Lasaeola pristina</i> (Wunderlich, 1986)	Ne Dominican amber
440. <i>Lasaeola puta</i> Wunderlich, 1988	Ne Dominican amber
441. <i>Lasaeola sexsaetosa</i> Wunderlich, 2008b	Pa Baltic amber
442. ? <i>Lasaeola sigillata</i> Wunderlich, 2008b	Pa Bitterfeld amber
443. <i>Lasaeola vicina</i> (Wunderlich, 1982)	Ne Dominican amber
444. <i>Lasaeola vicinoides</i> Wunderlich, 1988	Ne Dominican amber
<i>Lasaeola</i> sp. <i>in</i> Wunderlich (1988)	Ne Dominican amber
<i>Lasaeola</i> spp. <i>in</i> Wunderlich (2008b)	Pa Baltic / Bitt. amber
† Medela Petrunkevitch, 1942 [?Theridiidae, cf. Wunderlich (2008b)]	Palaeogene
445. <i>Medela baltica</i> Petrunkevitch, 1942*	Pa Baltic amber
† Mimetidion Wunderlich, 2008b	Palaeogene
446. <i>Mimetidion furca</i> Wunderlich, 2008b*	Pa Baltic amber
† Nanomysmena Petrunkevitch, 1958	Palaeogene
447. <i>Nanomysmena aculeata</i> Petrunkevitch, 1958	Pa Baltic amber
448. <i>Nanomysmena munita</i> Petrunkevitch, 1958	Pa Baltic amber
449. <i>Nanomysmena palanga</i> Marusik & Penney, 2004	Pa Baltic amber
450. <i>Nanomysmena petrunkevitchi</i> Marusik & Penney, 2004	Pa Baltic amber
451. <i>Nanomysmena pseudogracilis</i> Marusik & Penney, 2004	Pa Baltic amber
† Nanosteatoda Wunderlich, 2008b	Palaeogene

452. <i>Nanosteatoda breviscutum</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
453. <i>Nanosteatoda trisetae</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
† Obscuropholcomma Wunderlich, 2008<i>b</i>	Palaeogene
454. <i>Obscuropholcomma</i> sp. <i>in</i> Wunderlich (2012 <i>b</i>)	Pa Rovno amber
455. <i>Obscuropholcomma tegens</i> Wunderlich, 2008 <i>b</i> *	Pa Baltic amber
Phoroncidia Westwood, 1835	Quaternary – Recent
456. <i>Phoroncidia ?aculeata</i> Westwood, 1835 [Recent]	Qt Madagascan copal
Platnickina Koçak & Kemal, 2008	Quaternary – Recent
457. <i>Platnickina duosetae</i> Wunderlich, 2012 <i>a</i>	Qt Madagascan copal
† Praetereuryopsis Wunderlich, 2008<i>b</i>	Palaeogene
458. <i>Praetereuryopsis phoroncidoides</i> Wunderlich, 2008 <i>b</i> *	Pa Baltic amber
† Pronepos Petrunkevitch, 1963	Neogene
459. <i>Pronepos exilis</i> Petrunkevitch, 1963*	Ne Chiapas amber
460. <i>Pronepos fossilis</i> Petrunkevitch, 1963	Ne Chiapas amber
† Protosteatoda Wunderlich, 2008<i>b</i>	Palaeogene
461. <i>Protosteatoda gutta</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
† Pseudoteutana Wunderlich, 2008<i>b</i>	Palaeogene
462. <i>Pseudoteutana stigmata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Eomysmena stridens</i> Petrunkevitch, 1958.....	Pa Baltic amber
ii. = <i>Flegia succini</i> Petrunkevitch, 1942	Pa Baltic amber
† Rugapholcomma Wunderlich, 2008<i>b</i>	Palaeogene
463. <i>Rugapholcomma patellaris</i> Wunderlich, 2008 <i>b</i> *	Pa Baltic amber
† Spinisinus Wunderlich, 2008<i>b</i>	Palaeogene
464. <i>Spinisinus parvioculi</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
465. <i>Spinisinus splendidus</i> Wunderlich, 2008 <i>b</i> *	Pa Baltic amber
† Spinitharinus Wunderlich, 2008<i>b</i>	Palaeogene
466. <i>Spinitharinus bulbosus</i> Wunderlich, 2008 <i>b</i> *	Pa Baltic / Bitt. amber
467. <i>Spinitharinus cheliceratus</i> Wunderlich, 2008 <i>b</i>	Pa Baltic / Bitt. amber
468. <i>Spinitharinus coniectens</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
469. <i>Spinitharinus curvatus</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
470. <i>Spinitharinus cymbioseta</i> Wunderlich, 2008 <i>b</i>	Pa Baltic amber
<i>Spinitharinus</i> spp. <i>in</i> Wunderlich (2008 <i>b</i>)	Pa Baltic amber
Spintharus Hentz, 1850	Neogene – Recent
471. <i>Spintharus longisoma</i> Wunderlich, 1988	Ne Dominican amber
Steatoda Sundevall, 1833	?Palaeogene – Recent
472. ‘ <i>Steatoda</i> ’ <i>anticus</i> (Berland, 1939)	Pa Baltic amber
Stemmops O. P.-Cambridge, 1894	Neogene – Recent
473. <i>Stemmops incertus</i> Wunderlich, 1988	Ne Dominican amber
474. <i>Stemmops prominens</i> Wunderlich, 1988	Ne Dominican amber
Styposis Simon, 1894	Neogene – Recent
475. <i>Styposis pholcoides</i> Wunderlich, 1988	Ne Dominican amber

† <i>Succinobertus</i> Wunderlich, 2008b	Palaeogene
476. <i>Succinobertus adjacens</i> Wunderlich, 2008b*	Pa Baltic / Bitt. Amber
† <i>Succinura</i> Wunderlich, 2008b	Palaeogene
477. <i>Succinura aciesaeta</i> Wunderlich, 2008b	Pa Baltic amber
478. <i>Succinura bellavista</i> Wunderlich, 2008b*	Pa Baltic amber
479. <i>Succinura circuita</i> Wunderlich, 2008b	Pa Baltic amber
480. <i>Succinura dubia</i> Wunderlich, 2008b	Pa Baltic amber
481. <i>Succinura fuscoruber</i> Wunderlich, 2008b	Pa Baltic amber
482. <i>Succinura ovalis</i> Wunderlich, 2008b	Pa Baltic amber
<i>Succinura</i> sp. in Wunderlich (2008b)	Pa Baltic amber
<i>Theridion</i> Walckenaer, 1805	?Cretaceous – Recent
483. ' <i>Theridion</i> ' <i>alutaceum</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
484. <i>Theridion annulipes</i> Heer, 1865	Ne Öhningen
485. <i>Theridion atalus</i> Chang, 2004 [both generic and familial assignment unreliable!]	K Jehol Biota
486. ' <i>Theridion</i> ' <i>berendti</i> Marusik & Penney, 2004	Pa Baltic amber
i. = <i>Theridion globosa</i> C. L. Koch & Berendt, 1854 [preoccupied]	
487. <i>Theridion bucklandi</i> Thorell, 1870a	Pa Aix-en-Provence
488. <i>Theridion contrarium</i> Wunderlich, 1988	Ne Dominican amber
489. <i>Theridion crassipalpus</i> Berland, 1939	Pa Aix-en-Provence
490. ' <i>Theridion</i> ' <i>detersum</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
491. <i>Theridion erectoides</i> Wunderlich, 1988	Ne Dominican amber
492. <i>Theridion erectum</i> Wunderlich, 1988	Ne Dominican amber
493. ' <i>Theridion</i> ' <i>globosus</i> (Presl, 1822)	Pa Baltic amber
494. <i>Theridion globulus</i> Heer, 1865	Ne Öhningen
495. ' <i>Theridion</i> ' <i>hirtum</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
496. <i>Theridion inversum</i> Wunderlich, 1988	Ne Dominican amber
497. <i>Theridion maculipes</i> Heer, 1865	Ne Öhningen
498. ' <i>Theridion</i> ' <i>oblongum</i> (Presl, 1822)	Pa Baltic amber
499. ' <i>Theridion</i> ' <i>ovale</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
500. ' <i>Theridion</i> ' <i>ovatum</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
501. ' <i>Theridion</i> ' <i>simplex</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
502. <i>Theridion variosoma</i> Wunderlich, 1988	Ne Dominican amber
503. <i>Theridion wunderlichi</i> Penney, 2001	Ne Dominican amber
i. = <i>Theridion ovale</i> Wunderlich, 1988 [preoccupied]	
† <i>Thyelia</i> C. L. Koch & Berendt, 1854	Palaeogene
504. <i>Thyelia anomala</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
505. <i>Thyelia convexa</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
506. <i>Thyelia fossula</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
507. <i>Thyelia marginata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
508. <i>Thyelia pallida</i> C. L. Koch & Berendt, 1854	Pa Baltic amber

509.	<i>Thyelia scotina</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
510.	<i>Thyelia tristis</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
511.	<i>Thyelia villosa</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
Ulesanis L. Koch, 1872		Palaeogene – Recent
512.	<i>Ulesanis antecessor</i> Wunderlich, 2008b	Pa Baltic Amber
513.	<i>Ulesanis frontprocera</i> Wunderlich, 2008b	Pa Baltic Amber
514.	<i>Ulesanis longicymbium</i> Wunderlich, 2008b	Pa Baltic Amber
515.	<i>Ulesanis ovalis</i> Wunderlich, 2008b	Pa Baltic / Bitt. amber
516.	<i>Ulesanis parva</i> Wunderlich, 2008b	Pa Baltic / Bitt. amber
†	Unispinatoda Wunderlich, 2008b	Palaeogene
517.	<i>Unispinatoda aculeata</i> Wunderlich, 2008b*	Pa Baltic / Bitt. Amber
†	Vicipholcomma Wunderlich, 2008b	Palaeogene
518.	<i>Vicipholcomma spiralis</i> Wunderlich, 2008b*	Pa Baltic Amber
Theridiidae incertae sedis		
519.	' <i>Eomysmena</i> ' <i>succini</i> (Petrunkevitch, 1942)	Pa Baltic amber
520.	' <i>Anelosimus</i> ' <i>clypeatus</i> Wunderlich, 1988	Ne Dominican amber
THERIDIOSOMATIDAE Simon, 1881		Cretaceous – Recent
	Theridiosomatidae gen. et sp. indet <i>in</i> Wunderlich (2004i)	Pa Baltic amber
	Theridiosomatidae gen. et sp. indet <i>in</i> Wunderlich (2011f)	Qt Madagascar copal
†	Eocoddingtonia Selden, 2010	Cretaceous
521.	<i>Eocoddingtonia eskovi</i> Selden, 2010*	K Baissa, Transbaikalia
†	Eoepeirotypus Wunderlich, 2004j	Palaeogene
522.	<i>Eoepeirotypus retrobulbus</i> Wunderlich, 2004j*	Pa Baltic amber
	<i>Eoepeirotypus</i> sp. <i>in</i> Wunderlich (2004)	Pa Bitterfeld amber
†	Eotheridiosoma Wunderlich, 2004j	Palaeogene
523.	? <i>Eotheridiosoma hamatum</i> Wunderlich, 2011e	Pa Baltic amber
524.	<i>Eotheridiosoma tuber</i> Wunderlich, 2004j*	Pa Bitterfeld amber
525.	<i>Eotheridiosoma volutum</i> Wunderlich, 2004j	Pa Bitterfeld amber
†	Hypotheridiosoma Wunderlich, 2012d	Cretaceous
526.	<i>Hypotheridiosoma paracymbium</i> Wunderlich, 2012d*	K Myanmar amber
†	Leviunguis Wunderlich, 2012d	Cretaceous
527.	<i>Leviunguis bruckschi</i> Wunderlich, 2012d*	K Myanmar amber
†	Palaeoepeirotypus Wunderlich, 1988	Neogene
528.	<i>Palaeoepeirotypus iuvenis</i> Wunderlich, 1988*	Ne Dominican amber
529.	<i>Palaeoepeirotypus iuvenoides</i> Wunderlich, 1988	Ne Dominican amber
†	Spinitheridiosoma Wunderlich, 2004j	Palaeogene
	NB: type species designated from the wrong genus!	
530.	<i>Spinitheridiosoma balticum</i> Wunderlich, 2004j	Pa Baltic amber
531.	<i>Spinitheridiosoma bispinosum</i> Wunderlich, 2004j	Pa Bitterfeld amber
532.	<i>Spinitheridiosoma rima</i> Wunderlich, 2004j	Pa Baltic amber

<i>Theridiosoma</i> O. P.-Cambridge, 1879b	Neogene – Recent
533. <i>Theridiosoma incompletum</i> Wunderlich, 1988	Ne Dominican amber
† <i>Umerosoma</i> Wunderlich, 2004j	Palaeogene
534. <i>Umerosoma multispina</i> Wunderlich, 2004j*	Pa Baltic amber
SYMPHYTOGNATHIDAE Hickman, 1931	Recent
no fossil record	
ANAPIDAE Simon, 1895	Palaeogene – Recent
= TEXTRICELLIDAE Hickman, 1945	
† <i>Balticonopsis</i> Wunderlich, 2004k	Palaeogene
535. <i>Balticonopsis bispina</i> Wunderlich, 2004k	Pa Baltic amber
536. <i>Balticonopsis bitterfeldensis</i> Wunderlich, 2004k	Pa Bitterfeld amber
537. <i>Balticonopsis bulbosa</i> Wunderlich, 2004k	Pa Baltic amber
538. <i>Balticonopsis ceranowiczae</i> Wunderlich, 2004k	Pa Baltic amber
539. <i>Balticonopsis holti</i> Wunderlich, 2004k*	Pa Baltic amber
540. <i>Balticonopsis perkovskyi</i> Wunderlich, 2004ar	Pa Rovno amber
541. <i>Balticonopsis thomasi</i> Wunderlich, 2004k	Pa Baltic amber
<i>Balticonopsis</i> sp. in Wunderlich (2004k)	Pa Baltic amber
† <i>Dubianapis</i> Wunderlich, 2004k	Palaeogene
542. <i>Dubianapis obscura</i> Wunderlich, 2004k*	Pa Baltic amber
† <i>Flagellanapis</i> Wunderlich, 2004k	Palaeogene
543. <i>Flagellanapis voighti</i> Wunderlich, 2004k*	Pa Baltic/Bitt. Amber
† <i>Fossilanapis</i> Wunderlich, 2004k	Palaeogene
544. <i>Fossilanapis anderseri</i> Wunderlich, 2004k	Pa Baltic amber
545. <i>Fossilanapis baetcheri</i> Wunderlich, 2004k*	Pa Baltic amber
546. <i>Fossilanapis eichmanni</i> Wunderlich, 2004k	Pa Baltic amber
547. <i>Fossilanapis flexiotarsus</i> Wunderlich, 2004k	Pa Baltic amber
548. <i>Fossilanapis multispinae</i> Wunderlich, 2011h	Pa Baltic amber
549. <i>Fossilanapis saltans</i> Wunderlich, 2004k	Pa Baltic amber
550. <i>Fossilanapis unispinum</i> Wunderlich, 2004k	Pa Baltic amber
<i>Fossilanapis</i> sp. in Wunderlich (2004k)	Pa Bitterfeld amber
<i>Fossilanapis</i> sp. in Wunderlich (2011h)	Pa Baltic amber
† <i>Palaeoanapis</i> Wunderlich, 1988	Neogene
551. <i>Palaeoanapis nana</i> Wunderlich, 1988*	Ne Dominican amber
† <i>Ruganapis</i> Wunderlich, 2004k	Palaeogene
552. <i>Ruganapis scutata</i> Wunderlich, 2004k*	Pa Baltic amber
† <i>Saxonanapis</i> Wunderlich, 2004k	Palaeogene
553. <i>Saxonanapis grabenhorsti</i> Wunderlich, 2004k*	Pa Baltic/Bitt. Amber
† <i>Tuberanapis</i> Wunderlich, 2004k	Palaeogene
554. <i>Tuberanapis parvibulbus</i> Wunderlich, 2004k*	Pa Baltic amber

COMAROMIDAE Wunderlich, 2004 [stat. nov. 2011].....	Palaeogene – Recent
† Balticoroma Wunderlich, 2004k	Palaeogene
= † <i>Balticorma</i> [sic] Weitschat & Wichard, 2002 [<i>nomen nudum</i>]	
555. <i>Balticoroma damzeni</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
556. <i>Balticoroma ernstorum</i> Wunderlich, 2004 <i>k</i>	Pa Baltic/Bitt. amber
557. <i>Balticoroma gracilipes</i> Wunderlich 2004 <i>k</i>	Pa Baltic/Bitt. amber
558. <i>Balticoroma reschi</i> Wunderlich, 2004 <i>k</i> *	Pa Baltic amber
559. <i>Balticoroma serafinorum</i> Wunderlich, 2004 <i>k</i>	Pa Baltic/Bitt. amber
560. <i>Balticoroma tibialis</i> Wunderlich, 2004 <i>k</i>	Pa Baltic amber
561. <i>Balticoroma wheateri</i> Penney & Marusik, 2011 <i>in</i> Penney <i>et al.</i>	Pa Baltic amber
MYSMENIDAE Petrunkevitch, 1928	Palaeogene – Recent
Mysmeninae sp. <i>in</i> Wunderlich (2004 <i>ar</i>)	Pa Rovno amber
† Dominicanopsis Wunderlich, 2004k	Neogene
562. <i>Dominicanopsis grimaldii</i> Wunderlich, 2004 <i>k</i> *	Ne Dominican amber
† Eomysmenopsis Wunderlich, 2004k	Palaeogene
563. <i>Eomysmenopsis spinipes</i> Wunderlich, 2004 <i>k</i> *	Pa Baltic / Bitt. Amber
Mysmena Simon, 1894	Palaeogene – Recent
<i>Mysmena</i> (s. l.) sp. indet <i>in</i> Wunderlich (2012a)	Qt Madagascan copal
564. <i>Mysmena</i> (s.l.) <i>copalis</i> Wunderlich, 2011 <i>f</i>	Qt Madagascan copal
565. <i>Mysmena curvata</i> Wunderlich, 2011 <i>h</i>	Pa Baltic amber
566. <i>Mysmena dominicana</i> Wunderlich, 1998	Qt Madagascan copal
567. <i>Mysmena fossilis</i> Petrunkevitch, 1971	Ne Chiapas amber
568. <i>Mysmena groehni</i> Wunderlich, 2004 <i>k</i>	Pa Baltic / Bitt. amber
569. <i>Mysmena grotae</i> Wunderlich, 2004 <i>k</i>	Pa Baltic amber
Mysmenopsis Simon, 1897b	Neogene – Recent
570. <i>Mysmenopsis lissycolleyae</i> Penney, 2000	Ne Dominican amber
† Palaeomysmena Wunderlich, 2004k	Palaeogene
571. <i>Palaeomysmena hoffeinsorum</i> Wunderlich, 2004 <i>k</i> *	Pa Baltic amber
† BALTSUCCINIDAE Wunderlich, 2004l	Palaeogene
† Baltsuccinus Wunderlich, 2004l	Palaeogene
572. <i>Baltsuccinus flagellaceus</i> Wunderlich, 2004 <i>l</i> *	Pa Baltic amber
573. <i>Baltsuccinus similis</i> Wunderlich, 2004 <i>l</i>	Pa Baltic amber
† PROTHERIDIIDAE Wunderlich, 2004l	Cretaceous – Palaeo.
† Protheridion Wunderlich, 2004l	Palaeogene
574. <i>Protheridion bitterfeldensis</i> Wunderlich, 2004 <i>l</i>	Pa Bitterfeld amber
575. <i>Protheridion detritus</i> Wunderlich, 2004 <i>l</i>	Pa Baltic amber
576. <i>Protheridion obscurum</i> Wunderlich, 2004 <i>l</i>	Pa Baltic amber
577. <i>Protheridion punctatum</i> Wunderlich, 2004 <i>l</i>	Pa Baltic amber
578. <i>Protheridion tibialis</i> Wunderlich, 2004 <i>l</i> *	Pa Baltic amber

† Zarqaraneus Wunderlich, 2008d	Cretaceous
579. <i>Zarqaraneus hudaе</i> Wunderlich, 2008d*	K Jordanian amber
† PRAETHERIDIIDAE Wunderlich, 2012c	Palaeogene
† <i>Praetheridion</i> Wunderlich, 2004l	Palaeogene
580. <i>Praetheridion fleissneri</i> Wunderlich, 2004l*	Pa Baltic amber
SYNAPHRIDAE Wunderlich, 1986	Palaeogene – Recent
† <i>Iardinidis</i> Wunderlich 2004k	Palaeogene
581. <i>Iardinidis brevipes</i> Wunderlich, 2004k*	Pa Baltic amber
PIMOIDAE Wunderlich, 1986	Palaeogene – Recent
<i>Pimoa</i> Chamberlin & Ivie, 1943	Palaeogene – Recent
582. <i>Pimoa expandens</i> Wunderlich, 2004r	Pa Baltic amber
583. <i>Pimoa (Eopimoa) hormigai</i> Wunderlich, 2004r	Pa Baltic amber
584. <i>Pimoa inopinata</i> Wunderlich, 2004r	Pa Baltic amber
585. <i>Pimoa liedtkei</i> Wunderlich, 2004r	Pa Baltic amber
586. <i>Pimoa lingua</i> Wunderlich, 2004r	Pa Baltic amber
587. <i>Pimoa (Eopimoa) longiscapus</i> Wunderlich, 2008a	Pa Baltic amber
588. <i>Pimoa multicuspuli</i> Wunderlich, 2004r	Pa Baltic amber
589. <i>Pimoa (Eopimoa) obruens</i> Wunderlich, 2008a	Pa Baltic amber
<i>Pimoa</i> sp. <i>in</i> Wunderlich (2004r)	Pa Baltic amber
<i>Pimoa (Eopimoa)</i> sp. <i>in</i> Wunderlich (2008a)	Pa Baltic amber
PUMILIOPIMOIDAE Wunderlich, 2008a	Palaeogene – Recent
† <i>Pumiliopimoa</i> Wunderlich, 2008a	Palaeogene
590. <i>Pumiliopimoa parma</i> Wunderlich, 2008a*	Pa Baltic amber
SINOPIMOIDAE Li & Wunderlich, 2008	Recent
no fossil record	
LINYPHIIDAE Blackwall, 1859	Cretaceous – Recent
= MICRYPHANTIDAE Bertkau, 1878a	
= ERIGONIDAE Simon, 1884c	
?Linyphiidae gen. et sp. indet <i>in</i> McAlpine & Martin (1969)	K Canadian amber
Linyphiidae gen. et sp. indet <i>in</i> Penney (2002)	K New Jersey amber
Linyphiidae gen. et sp. indet <i>in</i> Schmidt <i>et al.</i> (2010)	K Ethiopian amber
Linyphiinae gen. et sp. indet <i>in</i> Penney & Selden (2002)	K Lebanese amber
[NB: Wunderlich (2012d) questioned the veracity of these Cretaceous linyphiids.]	
† <i>Agynetiophantes</i> Wunderlich, 2004s	Palaeogene
591. <i>Agynetiophantes gibbiferus</i> Wunderlich, 2004s*	Pa Baltic amber
<i>Ceratinopsis</i> Emerton, 1882	Quaternary – Recent

592. <i>Ceratinopsis deformans</i> (Wunderlich, 1998)	Qt	Madagascan copal
Cnephalocotes Simon, 1884c	Quaternary – Recent	
593. <i>Cnephalocotes obscurus</i> (Blackwall, 1834b) [Recent]	Qt	England
† Custodela Petrunkevitch, 1942	Palaeogene	
= † <i>Obnisis</i> Petrunkevitch, 1942 [tentative synonymy]		
594. <i>Custodela acuta</i> Wunderlich, 2004s	Pa	Baltic amber
595. <i>Custodela acutula</i> Wunderlich, 2004s	Pa	Bitterfeld amber
596. <i>Custodela bispina</i> Wunderlich, 2004s	Pa	Bitterfeld amber
597. <i>Custodela bispinosa</i> Wunderlich, 2004s	Pa	Bitterfeld amber
598. <i>Custodela cheiracantha</i> (C. L. Koch & Berendt, 1854)*	Pa	Baltic amber
599. <i>Custodela clava</i> Wunderlich, 2004s	Pa	Baltic amber
600. <i>Custodela curva</i> Wunderlich, 2004s	Pa	Baltic amber
601. <i>Custodela curvata</i> Wunderlich, 2004s	Pa	Bitterfeld amber
602. <i>Custodela divergens</i> Wunderlich, 2004s	Pa	Baltic amber
603. <i>Custodela expandens</i> Wunderlich, 2004s	Pa	Baltic amber
604. <i>Custodela falcata</i> Wunderlich, 2004s	Pa	Baltic amber
605. <i>Custodela femurspinosa</i> Wunderlich, 2004s	Pa	Bitterfeld amber
606. <i>Custodela henningseni</i> Wunderlich, 2004s	Pa	Baltic amber
607. <i>Custodela kochi</i> Wunderlich, 2004s	Pa	Baltic amber
608. <i>Custodela lamellata</i> (Wunderlich, 1988)	Pa	Baltic amber
609. <i>Custodela lanx</i> Wunderlich, 2004s	Pa	Baltic amber
610. <i>Custodela oblonga</i> (C. L. Koch & Berendt, 1854)	Pa	Baltic amber
611. <i>Custodela obtusa</i> Wunderlich, 2004s	Pa	Baltic amber
612. ? <i>Custodela parva</i> Wunderlich, 2004s	Pa	Bitterfeld amber
613. <i>Custodela pseudokochi</i> Wunderlich, 2004s	Pa	Baltic amber
614. <i>Custodela stridulans</i> Wunderlich, 2004s	Pa	Bitterfeld amber
615. <i>Custodela tenuipes</i> (Petrunkevitch, 1942)	Pa	Baltic amber
616. <i>Custodela tibialis</i> Wunderlich, 2004s	Pa	Baltic amber
<i>Custodela</i> sp. <i>in</i> Wunderlich (2004s)	Pa	Bitterfeld amber
† Custodelela Wunderlich, 2004s	Palaeogene	
617. <i>Custodelela hamata</i> Wunderlich, 2004s*	Pa	Bitterfeld amber
† Eolabulla Wunderlich, 2004s	Palaeogene	
618. <i>Eolabulla falcata</i> Wunderlich, 2004s	Pa	Baltic amber
619. <i>Eolabulla gladiformis</i> Wunderlich, 2004s	Pa	Baltic amber
620. <i>Eolabulla laminata</i> Wunderlich, 2004s*	Pa	Baltic amber
621. <i>Eolabulla perforata</i> Wunderlich, 2004s	Pa	Baltic amber
622. <i>Eolabulla sagitta</i> Wunderlich, 2004s	Pa	Baltic amber
623. <i>Eolabulla similis</i> Wunderlich, 2004s	Pa	Baltic amber
<i>Eolabulla</i> sp. 1–2 <i>in</i> Wunderlich (2004s)	Pa	Baltic amber
† Eophantes Wunderlich, 2004s	Palaeogene	
624. <i>Eophantes complicatus</i> Wunderlich, 2004s*	Pa	Baltic amber

625. ? <i>Eophantes seorsum</i> Wunderlich, 2012c	Pa Baltic amber
<i>Erigone</i> Audouin, 1826	Neogene – Recent
<i>Erigone</i> sp. in Hopkins <i>et al.</i> (1976)	Qt Alaska
626. <i>Erigone atra</i> Blackwall, 1833 [Recent]	Qt England
627. ? <i>Erigone dechenii</i> Bertkau, 1878b	Ne Rott, Germany
<i>Floricomus</i> Crosby & Bishop, 1925	Neogene – Recent
628. <i>Floricomus fossilis</i> Penney, 2005c	Ne Dominican amber
<i>Gonatium</i> Menge, 1868	Quaternary – Recent
629. <i>Gonatium rubens</i> (Blackwall, 1833) [Recent]	Qt England
<i>Hypselistes</i> Simon, 1894	Quaternary – Recent
630. <i>Hypselistes jacksoni</i> (O. P.-Cambridge, 1902) [Recent]	Qt England
<i>Linyphia</i> Latreille, 1804a	Palaeogene – Recent
631. <i>Linyphia andraei</i> Bertkau, 1878b	Ne Rott, Germany
632. <i>Linyphia byrami</i> Cockerell, 1925	Pa Green River
633. <i>Linyphia florissantii</i> Petrunkevitch, 1922	Pa Florissant
634. <i>Linyphia pachygnathoides</i> Petrunkevitch, 1922	Pa Florissant
635. <i>Linyphia quievreuxi</i> Berland, 1939	Pa Aix-en-Provence
636. <i>Linyphia retensa</i> Scudder, 1890a	Pa Florissant
637. <i>Linyphia rottensis</i> Bertkau, 1878b	Ne Rott, Germany
638. <i>Linyphia seclusa</i> (Scudder, 1890a)	Pa Florissant
† <i>Madagascarphantes</i> Wunderlich, 2012a	Quaternary
639. <i>Madagascarphantes vomerans</i> Wunderlich, 2012a*	Qt Madagascan copal
† <i>Malepellis</i> Petrunkevitch, 1971	Neogene
640. <i>Malepellis extincta</i> Petrunkevitch, 1971*	Ne Chiapas amber
<i>Meioneta</i> Hull, 1920	Neogene – Recent
641. <i>Meioneta bigibber</i> (Wunderlich, 1988)	Ne Dominican amber
642. <i>Meioneta fastigata</i> (Wunderlich, 1988)	Ne Dominican amber
643. <i>Meioneta separata</i> (Wunderlich, 1988)	Ne Dominican amber
<i>Meioneta</i> sp. in Wunderlich (1988)	Ne Dominican amber
<i>Micryphantes</i> C. L. Koch, 1833	Palaeogene
644. <i>Micryphantes molybdinus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
645. <i>Micryphantes regularis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† <i>Mystagogus</i> Petrunkevitch, 1942 ... [Wunderlich suggests possibly in Cyatholipidae]	Palaeogene
646. <i>Mystagogus dubius</i> Petrunkevitch, 1958	Pa Baltic amber
647. <i>Mystagogus glaber</i> Petrunkevitch, 1942*	Pa Baltic amber
† <i>Paralabulla</i> Wunderlich, 2004s	Palaeogene
648. <i>Paralabulla bitterfeldensis</i> Wunderlich, 2004s*	Pa Bitterfeld amber
649. ? <i>Paralabulla dubia</i> Wunderlich, 2004s	Pa Baltic amber
650. <i>Paralabulla succinifera</i> Wunderlich, 2004s	Pa Baltic amber
<i>Paralabulla</i> sp. in Wunderlich (2004s, 2012c)	Pa Bitterfeld amber
<i>Pocadicnemis</i> Simon, 1884c	Quaternary – Recent

651. <i>Pocadicnemis pumila</i> (Blackwall, 1841) [Recent]	Qt	England
Savignia Blackwall, 1833	Quaternary – Recent	
652. <i>Savignia frontata</i> Blackwall, 1833 [Recent]	Qt	England
Selenyphantes Gertsch & Davis, 1946	Neogene – Recent	
= † <i>Palaeolinyphia</i> Wunderlich, 1986		
653. <i>Selenyphantes flagellifera</i> (Wunderlich, 1986)	Ne	Dominican amber
† Succineta Wunderlich, 2004s	Palaeogene	
654. <i>Succineta brevispina</i> Wunderlich, 2004s	Pa	Baltic amber
655. <i>Succineta discoidalis</i> Wunderlich, 2004s*	Pa	Baltic amber
<i>Succineta</i> sp. in Wunderlich (2004s)	Pa	Baltic amber
† Succiphantes Wunderlich, 2004s	Palaeogene	
656. <i>Succiphantes tanasevitchi</i> Wunderlich, 2004s	Pa	Baltic amber
657. <i>Succiphantes velteni</i> Wunderlich, 2004s*	Pa	Baltic amber
Toschia Caporiacco, 1949	Quaternary – Recent	
658. ? <i>Toschia fossilis</i> Wunderlich, 2004as	Qt	Madagascan copal
TETRAGNATHIDAE Menge, 1866	Cretaceous – Recent	
= PACHYGNATHIDAE Menge, 1866		
= METIDAE Simon, 1894		
= NANOMETIDAE Forster & Forster, 1999		
† Anameta Wunderlich, 2004h	Palaeogene	
659. <i>Anameta distenda</i> Wunderlich, 2004h*	Pa	Bitterfeld amber
660. <i>Anameta kuntneri</i> Wunderlich, 2008a	Pa	Baltic amber
Azilia Keyserling, 1882	Neogene – Recent	
661. <i>Azilia hispaniolensis</i> Wunderlich, 1988	Ne	Dominican amber
i. = <i>Azilia muellenmeisteri</i> Wunderlich, 1988	Ne	Dominican amber
<i>Azilia</i> sp. in Wunderlich (1988)	Ne	Dominican amber
† Balticgnatha Wunderlich, 2011h	Palaeogene	
662. <i>Balticgnatha projectens</i> Wunderlich 2011h*	Pa	Baltic amber
† Baltleucauge Wunderlich, 2008a	Palaeogene	
663. <i>Baltleucauge gillespieae</i> Wunderlich 2008a*	Pa	Baltic amber
664. <i>Baltleucauge propinqua</i> Wunderlich, 2012c	Pa	Baltic amber
† Corneometa Wunderlich, 2004h	Palaeogene	
665. <i>Corneometa baltica</i> Wunderlich 2004h*	Pa	Baltic amber
666. <i>Corneometa pilosipes</i> Wunderlich 2004h	Pa	Baltic amber
Cyrtognatha Keyserling, 1882	Neogene – Recent	
667. <i>Cyrtognatha weitschati</i> Wunderlich, 1988	Ne	Dominican amber
† Eometa Petrunkevitch, 1958	Palaeogene	
668. <i>Eometa calefacta</i> Wunderlich, 2004h	Pa	Baltic amber
669. <i>Eometa longipes</i> Petrunkevitch, 1958	Pa	Baltic amber
670. <i>Eometa occulta</i> Wunderlich, 2004h	Pa	Baltic amber
671. <i>Eometa perfecta</i> Wunderlich, 2004h	Pa	Baltic amber

672. <i>Eometa samlandica</i> Petrunkevitch, 1958*	Pa Baltic amber
<i>Eometa</i> sp. 1–2 in Wunderlich (2004h)	Pa Baltic amber
Homalometa Simon, 1897b	Neogene – Recent
673. <i>Homalometa fossilis</i> Wunderlich, 1988	Ne Dominican amber
† Huergina Selden & Penney, 2003	Cretaceous
674. <i>Huergina diazromerali</i> Selden & Penney, 2003*	K Las Hoyas, Spain
† Macryphantes Selden, 1990	Cretaceous
675. <i>Macryphantes cowdeni</i> Selden, 1990*	K Sierra de Montsech
Meta C. L. Koch, 1836	Palaeogene – Recent
676. <i>Meta (Praetermeta) maculosa</i> Wunderlich, 2008a	Pa Baltic amber
677. <i>Meta (Praetermeta) velans</i> (Wunderlich, 2004h)	Pa Baltic amber
† Palaeometa Petrunkevitch, 1922	Palaeogene
678. <i>Palaeometa opertanea</i> (Scudder, 1890a)*	Pa Florissant
† Palaeopachygnatha Petrunkevitch, 1922	Palaeogene
679. <i>Palaeopachygnatha cockerelli</i> Petrunkevitch, 1922	Pa Florissant
680. <i>Palaeopachygnatha scudderi</i> Petrunkevitch, 1922*	Pa Florissant
† Priscometa Petrunkevitch, 1958	Palaeogene
681. <i>Priscometa capta</i> Wunderlich, 2004h	Pa Baltic amber
682. <i>Priscometa minor</i> Wunderlich, 2004h	Pa Baltic amber
683. <i>Priscometa tenuipes</i> Petrunkevitch, 1958*	Pa Baltic amber
† Samlandicmeta Wunderlich, 2012c	Palaeogene
684. <i>Samlandicmeta mutila</i> Wunderlich, 2012c	Pa Baltic amber
Tetragnatha Latreille, 1804a	Palaeogene – Recent
685. <i>Tetragnatha parva</i> (Hong, 1985)	Ne Shanwang
686. <i>Tetragnatha pristina</i> Schawaller, 1982c	Ne Dominican amber
687. <i>Tetragnatha tertiaria</i> Scudder, 1885	Pa Florissant
NEPHILIDAE Simon, 1894	Jurassic – Recent
Nephilidae indet. in Wunderlich (2012c)	Pa Baltic amber
† Cretaraneus Selden, 1990	Cretaceous
688. <i>Cretaraneus liaoningensis</i> Cheng, Meng & Wang in Cheng <i>et al.</i> , 2008	K Jehol biota
689. <i>Cretaraneus martensnetoi</i> Mesquita, 1996	K Crato Formation
690. <i>Cretaraneus vilaltae</i> Selden, 1990*	K Sierra de Montsech
† Eonephila Wunderlich, 2004i	Palaeogene
691. <i>Eonephila bitterfeldensis</i> Wunderlich, 2004i	Pa Bitterfeld amber
692. <i>Eonephila excellens</i> Wunderlich, 2004i*	Pa Baltic amber
693. <i>Eonephila longembolus</i> Wunderlich, 2004i	Pa Baltic amber
† Geratonephila Poinar in Poinar & Buckley, 2012	Cretaceous
694. <i>Geratonephila burmanica</i> Poinar in Poinar & Buckley, 2012*	K Myanmar amber
† Luxurionephila Wunderlich, 2004i	Palaeogene

695. <i>Luxurionephila spinifera</i> Wunderlich, 2004 <i>i</i>	Pa Baltic amber
† Minutunguis Wunderlich, 2011<i>f</i>	Quaternary
696. <i>Minutunguis silvestris</i> Wunderlich, 2011 <i>f</i> *	Qt Madagascar copal
Nephila Leach, 1815	Jurassic – Recent
697. <i>Nephila breviembolus</i> Wunderlich, 1986	Ne Dominican amber
698. <i>Nephila dommeli</i> Wunderlich, 1982	Ne Dominican amber
699. <i>Nephila furca</i> Wunderlich, 1986	Ne Dominican amber
700. <i>Nephila longembolus</i> Wunderlich, 1986	Ne Dominican amber
701. <i>Nephila jurassica</i> Selden, Shih & Ren, 2011	J Daohugou
702. <i>Nephila pennatipes</i> Scudder, 1885	Pa Florissant
703. <i>Nephila tenuis</i> Wunderlich, 1986	Ne Dominican amber
<i>Nephila</i> sp. in Dunlop & Penney (2012)	K Crato Formation
† Palaeonephila Wunderlich, 2004<i>i</i>	Palaeogene
704. <i>Palaeonephila brevis</i> Wunderlich, 2004 <i>i</i>	Pa Baltic amber
705. <i>Palaeonephila curvata</i> Wunderlich, 2004*	Pa Baltic amber
706. <i>Palaeonephila dilitans</i> Wunderlich, 2004 <i>i</i>	Pa Baltic amber
707. <i>Palaeonephila fibula</i> Wunderlich, 2004 <i>i</i>	Pa Baltic amber
708. <i>Palaeonephila longipes</i> Wunderlich, 2004 <i>i</i>	Pa Baltic amber
† JURARANEIDAE Eskov, 1984	Jurassic
† Juraraneus Eskov, 1984	Jurassic
709. <i>Juraraneus rasnitsyni</i> Eskov, 1984	J Transbaikalia
ARANEIDAE Simon, 1895	Cretaceous – Recent
= EPEIRIDAE Sundevall, 1833 [based on a generic synonym]	
= EUETRIIDAE Thorell, 1887 [based on a generic synonym]	
= ARGIOPIDAE Simon, 1890	
= ZYGIELLIDAE Simon, 1929	
?Araneinae sp. in Wunderlich (2004 <i>h</i>)	Pa Baltic amber
Araneidae gen. et sp. indet. in Ribera (2003)	Qt Girona, Spain
?Mangorini indet. in Wunderlich (2011 <i>a</i>)	Pa Baltic amber
† Anepeira Wunderlich, 2004<i>i</i>	Palaeogene
710. <i>Anepeira complicata</i> Wunderlich, 2004*	Pa Baltic amber
† Araneometa Wunderlich, 1988	Neogene
711. <i>Araneometa excelsa</i> Wunderlich, 1988	Ne Dominican amber
712. <i>Araneometa herrlingi</i> Wunderlich, 1988*	Ne Dominican amber
713. <i>Araneometa spirembolus</i> Wunderlich, 1988	Ne Dominican amber
<i>Araneometa</i> sp. in Wunderlich (1988)	Ne Dominican amber
Araneus Clerck, 1757	?Cretaceous – Recent
714. ? <i>Araneus</i> sp. in Wunderlich (2012 <i>c</i>)	Pa Baltic amber
715. <i>Araneus absconditus</i> (Scudder, 1890 <i>a</i>)	Pa Florissant
716. <i>Araneus aethus</i> Chang, 2004 [generic assignment unreliable!]	K Jehol biota

717. <i>Araneus beipiaoensis</i> Chang, 2004 [generic assignment unreliable!] K Jehol biota
718. <i>Araneus carbonaceous</i> Zhang, Sun & Zhang, 1994 Ne Shanwang
719. <i>Araneus cinctus</i> (Scudder, 1890a) Pa Florissant
720. <i>Araneus defunctus</i> Petrunkevitch, 1958 Pa Baltic amber
721. <i>Araneus delitus</i> (Scudder, 1890a) Pa Florissant
722. <i>Araneus emertoni</i> (Scudder, 1890a) Pa Florissant
723. <i>Araneus exustus</i> Petrunkevitch, 1963 Ne Chiapas amber
724. <i>Araneus kinchloeae</i> Dunlop & Jekel, 2009 Pa Florissant
i. = <i>Araneus indistinctus</i> (Petrunkevitch, 1922) [preoccupied]	
725. <i>Araneus inelegans</i> Zhang, Sun & Zhang, 1994 Ne Shanwang
726. <i>Araneus leptopodus</i> Zhang, Sun & Zhang, 1994 Ne Shanwang
727. <i>Araneus liaoxiensis</i> Chang, 2004 [generic assignment unreliable!] K Jehol biota
728. <i>Araneus longimanus</i> (Petrunkevitch, 1922) Pa Florissant
729. <i>Araneus (Calinurus) longipes</i> Dalman, 1826 Qt Copal
730. <i>Araneus luianus</i> Zhang, Sun & Zhang, 1994 Ne Shanwang
731. <i>Araneus meeki</i> (Scudder, 1890a) Pa Florissant
732. <i>Araneus molassicus</i> (Heer, 1865) Ne Öhningen
733. <i>Araneus nanus</i> Wunderlich, 1988 Ne Dominican amber
734. <i>Araneus piceus</i> Lin, Zhang & Wang, 1989 Ne Shanwang
735. <i>Araneus reheensis</i> Chang, 2004 [generic assignment unreliable!] K Jehol biota
736. <i>Araneus ruidipedalis</i> Zhang, Sun & Zhang, 1994 Ne Shanwang
737. <i>Araneus troschelii</i> (Bertkau, 1878b) Ne Rott, Germany
738. <i>Araneus vulcanalis</i> (Scudder, 1890a) Pa Florissant
Argiope Audouin, 1826	Neogene – Recent
= † <i>Magnaranea</i> Hong, 1985	
739. <i>Argiope furva</i> (Hong, 1985) Ne Shanwang
† Bararaneus Wunderlich, 2004i	Palaeogene
740. ? <i>Bararaneus annulatus</i> Wunderlich, 2004i Pa Baltic amber
741. <i>Bararaneus evolvens</i> Wunderlich, 2004i* Pa Baltic amber
† Chrysometata Wunderlich, 2004h	Palaeogene
742. <i>Chrysometata palaeartica</i> Wunderlich, 2004h* Pa Baltic amber
† Cyclososoma Petrunkevitch, 1958	Palaeogene
743. <i>Cyclososoma succini</i> Petrunkevitch, 1958* Pa Baltic amber
Enacrosoma Mello-Leitão, 1932	Neogene – Recent
744. <i>Enacrosoma verrucosa</i> (Wunderlich, 1988) Ne Dominican amber
† Eoaraneus Wunderlich, 2004i	Palaeogene
745. <i>Eoaraneus complexus</i> Wunderlich, 2004i* Pa Baltic amber
† Eochorizopes Wunderlich, 2008a	Palaeogene
746. <i>Eochorizopes szeklinskiae</i> Wunderlich, 2008a* Pa Baltic amber
† Eozygiella Wunderlich, 2004h	Palaeogene
747. <i>Eozygiella compacta</i> Wunderlich, 2004h* Pa Baltic amber

† Fossilaraneus Wunderlich, 1988	Neogene
748. <i>Fossilaraneus incertus</i> Wunderlich, 1988*	Ne Dominican amber
Gea C. L. Koch, 1843a	Palaeogene – Recent
749. <i>Gea krantzi</i> von Heyden, 1859	Ne Rott, Germany
† Graea Thorell, 1869	Palaeogene
= † <i>Eustaloides</i> Petrunkevitch, 1942	
750. ? <i>Graea aberrans</i> Wunderlich, 2004 <i>h</i>	Pa Baltic amber
751. <i>Graea bitterfeldensis</i> Wunderlich, 2004 <i>h</i>	Pa Bitterfeld amber
752. <i>Graea breviembolus</i> Wunderlich, 2004 <i>h</i>	Pa Baltic amber
753. <i>Graea brevis</i> Wunderlich, 2004 <i>h</i>	Pa Baltic amber
754. <i>Graea calceatus</i> (Petrunkevitch, 1950)	Pa Baltic amber
755. <i>Graea epeiroides</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
756. <i>Graea impudica</i> Wunderlich, 2004 <i>h</i>	Pa Baltic amber
757. <i>Graea lingula</i> Wunderlich, 2004 <i>h</i>	Pa Baltic amber
758. <i>Graea magnocoli</i> Wunderlich, 2012 <i>c</i>	Pa Baltic amber
759. <i>Graea minor</i> (Petrunkevitch, 1950)	Pa Baltic amber
760. <i>Graea setosa</i> (Petrunkevitch, 1942)	Pa Baltic amber
761. <i>Graea succini</i> Petrunkevitch, 1942	Pa Baltic amber
Hypognatha Guérin, 1839	Quaternary – Recent
762. <i>Hypognatha testudinaria</i> (Taczanowski, 1879) [Recent]	Qt Colombian copal
† Meditrina Petrunkevitch, 1942	Palaeogene
763. <i>Meditrina circumvallata</i> Petrunkevitch, 1942*	Pa Baltic amber
† Mesozygiella Penney & Ortuño, 2006	Cretaceous
764. <i>Mesozygiella dunlopi</i> Penney & Ortuño, 2006*	K Álava amber
† Miraraneus Wunderlich, 2004<i>i</i>	Palaeogene
765. <i>Miraraneus peregrinus</i> Wunderlich, 2004*	Pa Baltic amber
† Mirometa Petrunkevitch, 1963	Neogene
766. <i>Mirometa valdespinosa</i> Petrunkevitch, 1963	Ne Chiapas amber
Molinaranea Mello-Leitão, 1940	Neogene – Recent
767. <i>Molinaranea mitnickii</i> Saupe, Selden & Penney, 2010	Ne Dominican amber
† Pycnosinga Wunderlich, 1988	Neogene
768. <i>Pycnosinga fossilis</i> Wunderlich, 1988*	Ne Dominican amber
† Testudinaroides Dunlop & Jekel, 2008	Neogene
= † <i>Testudinaria</i> Zhang, Sun & Zhang, 1994 [preoccupied]	
769. <i>Testudinaroides papposa</i> (Zhang, Sun & Zhang, 1994)	Ne Shanwang
† Tethneus Scudder, 1885	Palaeogene
= † <i>Melanites</i> Hong, 1985	
770. <i>Tethneus guyoti</i> Scudder, 1890 <i>a</i>	Pa Florissant
771. <i>Tethneus hentzi</i> Scudder, 1885*	Pa Florissant
772. <i>Tethneus obduratus</i> Scudder, 1890 <i>a</i>	Pa Florissant
773. <i>Tethneus orbiculatus</i> (Hong, 1985)	Ne Shanwang

774. <i>Tethneus provectus</i> Scudder, 1890a	Pa Florissant
775. <i>Tethneus robustus</i> Petrunkevitch, 1922	Pa Florissant
776. <i>Tethneus twenhofeli</i> Petrunkevitch, 1922	Pa Florissant
Zilla C. L. Koch, 1834	Palaeogene – Recent
777. <i>Zilla gracilis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
778. <i>Zilla porrecta</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
779. <i>Zilla veterana</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
RETROLATERAL TIBIAL APOPHYSIS CLADE	Cretaceous – Recent
?RTA-clade <i>in</i> Wunderlich (2008 <i>d</i>)	K Myanmar amber
LYCOSOIDEA Sundevall, 1833	Cretaceous – Recent
† Korearachne Selden, Nam, Kim & Kim, 2012	Cretaceous
780. <i>Korearachne jinju</i> Selden, Nam, Kim & Kim, 2012*	K Sacheon, S. Korea
[Tentative assignment to Lycosoidea; disputed by Wunderlich (2012 <i>d</i>) who suggested it could be a haplogyne spider in Pholcoidea or Leptonetoidea]	
LYCOSIDAE Sundevall, 1833	?Cretaceous – Recent
Lycosidae gen. et sp. <i>in</i> Bottali (1975)	Qt Italy
Lycosidae gen. et sp. <i>in</i> Schawaller (1982 <i>d</i>)	Ne Willershausen
Lycosidae gen. et sp. <i>in</i> Penney (2001)	Ne Dominican amber
Lycosidae gen. et sp. <i>in</i> Kim & Nam (2012) [unreliable record]	K Lioyuan, China
Alopecosa Simon, 1885<i>b</i>	Quaternary – Recent
781. <i>Alopecosa ?pulverulenta</i> (Clerck, 1757) [Recent]	Qt England
† Dryadia Zhang, Sun & Zhang, 1994	Palaeogene
782. <i>Dryadia acanthopoda</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
Lycosa Latreille, 1804<i>a</i>	Palaeogene – Recent
783. <i>Lycosa florissanti</i> Petrunkevitch, 1922	Pa Florissant
784. <i>Lycosa lithographica</i> Schawaller & Ono, 1979	Ne Randecker Maar
785. <i>Lycosa malleata</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
786. <i>Lycosa miocaena</i> Schawaller & Ono, 1979	Ne Randecker Maar
787. <i>Lycosa subterranea</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
Pardosa C. L. Koch, 1847	Quaternary – Recent
788. <i>Pardosa pullata</i> (Clerck, 1757) [Recent]	Qt England
<i>Pardosa</i> sp. <i>in</i> Scott (2003)	Qt England
Pirata Sundevall, 1833	Quaternary – Recent
789. <i>Pirata ?piraticus</i> (Clerck, 1757) [Recent]	Qt England
Trochosa C. L. Koch, 1847	Quaternary – Recent
790. <i>Trochosa terricola</i> Thorell, 1856 [Recent]	Qt England
† PARATTIDAE Petrunkevitch, 1922	Palaeogene
† Parattus Petrunkevitch, 1922	Palaeogene
791. <i>Parattus evocatus</i> (Scudder, 1890 <i>a</i>)	Pa Florissant

792. <i>Parattus latitatus</i> (Scudder, 1890a)	Pa Florissant
793. <i>Parattus oculatus</i> Petrunkevitch, 1922	Pa Florissant
794. <i>Parattus resurrectus</i> (Scudder, 1890a)*	Pa Florissant
TRECHALEIDAE Simon, 1890	Palaeogene – Recent
= TRICLARIDAE O. P.-Cambridge, 1877 [<i>nomen oblitum</i>]	
= PERISSOBLEMMATIDAE O. P.-Cambridge, 1882b [based on a synonym]	
Trechaleidae sp. <i>in</i> Wunderlich (2004aa)	Pa Baltic amber
† <i>Eotrechalea</i> Wunderlich, 2004aa	Palaeogene
795. <i>Eotrechalea annulata</i> Wunderlich, 2004aa*	Pa Baltic amber
† <i>Esuritor</i> Petrunkevitch, 1942	Palaeogene
796. <i>Esuritor aculeatus</i> Petrunkevitch, 1958	Pa Baltic amber
797. <i>Esuritor spinipes</i> Petrunkevitch, 1942*	Pa Baltic amber
† <i>Linoptes</i> Menge, 1854	Palaeogene
798. ?' <i>Linoptes</i> ' <i>oculeus</i> Menge <i>in</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
NB: <i>Linoptes</i> mentioned as a <i>nomen nudum</i> by Wunderlich (2004z); this species listed by Wunderlich (2004aa) under Trechaleidae and another species under Pisauridae (see below)	
PISAURIDAE Simon, 1890	Palaeogene – Recent
= BRADYSTICHIDAE Simon, 1884	
= DOLOMEDIDAE Simon, 1898a	
= HALIDAE Jocqué, 1994	
Pisauridae sp. <i>in</i> Wunderlich (1988)	Pa Dominican amber
Pisauridae sp. <i>in</i> Wunderlich (2004z)	Pa Baltic amber
<i>Dolomedes</i> Latreille, 1804a	Quaternary – Recent
799. <i>Dolomedes fimbriatus</i> (Clerck, 1757) [Recent]	Qt England
† '<i>Linoptes</i>' Menge, 1854	Palaeogene
= † <i>Eopisaurella</i> Petrunkevitch, 1958	
NB: See notes on <i>Linoptes</i> under Trechaleidae above!	
800. ?' <i>Linoptes</i> ' <i>valdespinosa</i> (Petrunkevitch, 1958)*	Pa Baltic amber
?' <i>Linoptes</i> ' sp. 1–8 <i>in</i> Wunderlich (2004z)	Pa Baltic amber
† <i>Palaeoperenethis</i> Selden & Penney, 2009	Palaeogene
801. <i>Palaeoperenethis thaleri</i> Selden & Penney, 2009*	Pa British Columbia
OXYOPIDAE Thorell, 1870a	Palaeogene – Recent
= SPHASIDAE O. P.-Cambridge, 1871	
= HAMATALIVIDAE Marx, 1890b	
Oxyopidae sp. <i>in</i> Wunderlich 2004ab	Pa Bitterfeld amber
<i>Oxyopes</i> Latreille, 1804a	Palaeogene – Recent
802. <i>Oxyopes defectus</i> Wunderlich, 1988	Ne Dominican amber
803. ' <i>Oxyopes</i> ' <i>succini</i> Petrunkevitch, 1958	Pa Baltic amber
<i>Oxyopes</i> sp. <i>in</i> Wunderlich (1988, 2004ab)	Ne Dominican amber
† <i>Planoxyopes</i> Petrunkevitch, 1963	Neogene

804. *Planoxyopes eximius* Petrunkevitch, 1963* Ne Chiapas amber
 i. = *Planoxyopes fossilis* Wunderlich, 1988 [*lapsus*] Ne Chiapas amber
- SENOCULIDAE Simon, 1890** **Recent**
 = NEOTHEREUTOIDAE Holmberg, 1883 [based on a generic synonym]
- no fossil record
- STIPHIDIIDAE Dalmas, 1917** **Recent**
- no fossil record
- ZOROCRATIDAE Dahl, 1913** **Recent**
- no fossil record
- PSECHRIDAE Simon, 1890** **Recent**
- no fossil record
- ZOROPSIDAE Bertkau, 1882** **Palaeogene – Recent**
 Zoropsidae sp. *in* Wunderlich (2004x) Pa Baltic / Bitt. amber
- † ***Eomatachia* Petrunkevitch, 1942** **Palaeogene**
805. *Eomatachia barbarus* Wunderlich, 2004x Pa Baltic amber
 806. *Eomatachia bipartita* Wunderlich, 2004x Pa Baltic amber
 807. *Eomatachia divergens* Wunderlich, 2004x Pa Baltic amber
 808. *Eomatachia duplex* Wunderlich, 2004x Pa Baltic amber
 809. *Eomatachia latifrons* Petrunkevitch, 1942* Pa Baltic amber
 810. *Eomatachia recedens* Wunderlich, 2004x Pa Baltic amber
 811. *Eomatachia succini* (Petrunkevitch, 1942) Pa Baltic amber
 812. *Eomatachia wegneri* Wunderlich, 2004x Pa Baltic amber
 813. *Eomatachia xanthippe* Wunderlich, 2004x Pa Baltic amber
- † ***Eoprychia* Petrunkevitch, 1958** **Palaeogene**
814. *Eoprychia succini* Petrunkevitch, 1958* Pa Baltic amber
 815. *Eoprychia succinopsis* Wunderlich, 2004x Pa Baltic amber
 816. *Eoprychia vicina* Wunderlich, 2004x Pa Baltic amber
 Eoprychia sp. *in* Wunderlich (2004x) ?Pa not specified
- † ***Succiniropsis* Wunderlich, 2004x** **Palaeogene**
817. *Succiniropsis kutscheri* Wunderlich, 2004x* Pa Baltic / Bitt. Amber
 818. *Succiniropsis runcinata* Wunderlich, 2012c Pa Baltic amber
 819. *Succiniropsis samlandica* Wunderlich, 2004x Pa Baltic amber
- † **INSECUTORIDAE Petrunkevitch, 1942** **Palaeogene**
- † ***Insecutor* Petrunkevitch, 1942** **Palaeogene**
820. *Insecutor aculeatus* Petrunkevitch, 1942* Pa Baltic amber
 821. *Insecutor mandibulatus* Petrunkevitch, 1942 Pa Baltic amber

822. ? <i>Insecutor pecten</i> Wunderlich, 2004y	Pa Baltic amber
823. <i>Insecutor rufus</i> Petrunkevitch, 1942	Pa Baltic amber
824. ? <i>Insecutor spinifer</i> Wunderlich, 2004y	Pa Baltic amber
? <i>Insecutor</i> sp. in Wunderlich (2004y)	Pa Baltic amber
ZORIDAE F. O. P.-Cambridge, 1893	Palaeogene – Recent
† Zorapostenus Wunderlich, 2008c	Palaeogene
825. <i>Zorapostenus raveni</i> Wunderlich, 2008c	Pa Baltic amber
† SUCCINOMIDAE Wunderlich, 2012c	Palaeogene
† Eohalinobius Wunderlich, 2008c	Palaeogene
826. <i>Eohalinobius calefactus</i> Wunderlich, 2012c	Pa Baltic amber
827. <i>Eohalinobius hiddenseensis</i> Wunderlich, 2012c	Pa Baltic amber
828. <i>Eohalinobius patina</i> Wunderlich, 2012c	Pa Baltic amber
829. <i>Eohalinobius scutatus</i> Wunderlich, 2008c	Pa Baltic amber
† Succinomus Wunderlich, 2008c	Palaeogene
830. <i>Succinomus duomammillae</i> Wunderlich, 2008c	Pa Baltic amber
831. ? <i>Succinomus gibbosus</i> Wunderlich, 2012c	Pa Baltic amber
CTENIDAE Keyserling, 1877	Neogene – Recent
= ACANTHOCTENIDAE Simon, 1892b	
† Nanoctenus Wunderlich, 1988	Neogene
832. <i>Nanoctenus longipes</i> Wunderlich, 1988*	Ne Dominican amber
AGELENIDAE C. L. Koch, 1837	Palaeogene – Recent
= TEGENARIDAE Prach, 1860	
= † INCEPTORIDAE Petrunkevitch, 1942	
Agelena Walckenaer, 1805	Palaeogene – Recent
833. <i>Agelena tabida</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
Histopona Thorell, 1869	Palaeogene – Recent
834. ? <i>Histopona anthracina</i> Bertkau, 1878b	Ne Rott, Germany
† Inceptor Petrunkevitch, 1942	Palaeogene
835. <i>Inceptor aculeatus</i> Petrunkevitch, 1942*	Pa Baltic amber
836. <i>Inceptor dubius</i> Petrunkevitch, 1946	Pa Baltic amber
Tegenaria Latreille, 1804a	Palaeogene – Recent
837. ? <i>Tegenaria fragmentum</i> Wunderlich, 2004w	Pa Baltic amber
838. <i>Tegenaria lacazei</i> Gourret, 1887	Pa Aix-en-Provence
839. ? <i>Tegenaria obtusa</i> Wunderlich, 2004w	Pa Baltic amber
840. <i>Tegenaria virilis</i> Menge in C. L. Koch & Berendt, 1854	Pa Baltic amber
DICTYNOIDEA O. P.-Cambridge, 1871	Palaeogene – Recent
Dictynoidea incertae sedis	

- † *Sinodictyna* Hong, 1982 Palaeogene
 841. *Sinodictyna fushunensis* Hong, 1982* Pa Fu Shun amber
- CYBAEIDAE Simon, 1898a** Palaeogene – Recent
 = ARGYRONETIDAE Thorell, 1870a [both family names protected by usage]
- Argyroneta* Latreille, 1804a ?Neogene – Recent
 842. *Argyroneta aquatica* (Clerck, 1757) [Recent] Qt England
 843. ?*Argyroneta longipes* Heer, 1865 Ne Öhningen
- † *Vectaraneus* Selden, 2001 Palaeogene
 844. *Vectaraneus yulei* Selden, 2001* Pa Bembridge Marls
- DESIDAE Pocock, 1895** Palaeogene – Recent
Myro O. P.-Cambridge, 1876 Palaeogene – Recent
845. *Myro extinctus* Petrunkevitch, 1958 ...[possibly belongs in Dictynidae]..... Pa Baltic amber
 846. *Myro hirsutus* Petrunkevitch, 1942 Pa Baltic amber
- AMPHINECTIDAE Forster & Wilton, 1973** Recent
 = NEOLANIDAE Forster & Wilton, 1973
 no fossil record
- CYCLOCTENIDAE Simon, 1898a** Recent
 no fossil record
- HAHNIIDAE Bertkau, 1878a** Palaeogene – Recent
- † *Cymbiohahnia* Wunderlich, 2004v Palaeogene
 847. *Cymbiohahnia parens* Wunderlich, 2004v Pa Baltic, Bitterfeld & Rovno amber
- † *Eohahnia* Petrunkevitch, 1958 Palaeogene
 848. *Eohahnia succini* Petrunkevitch, 1958* Pa Baltic amber
- † *Protohahnia* Wunderlich, 2004v Palaeogene
 849. *Protohahnia antiqua* Wunderlich, 2004v* Pa Baltic amber
 850. *Protohahnia tripartita* Wunderlich, 2004v Pa Baltic amber
- genus uncertain**
 851. ‘*Tegenaria*’ *obscura* C. L. Koch & Berendt, 1854 Pa Baltic amber
- DICTYNIDAE O. P.-Cambridge, 1871** Cretaceous – Recent
 = RHIOIDAE Thorell, 1873
 = † ARTHRODICTYNIDAE Petrunkevitch, 1942
- Dictynidae gen. et sp. indet *in* Penney (2002) K New Jersey amber
 Dictynidae sp. 1–2 *in* Wunderlich (2004v) Pa Baltic amber
 Dictynidae sp. 1–5 *in* Wunderlich (2008d) K Myanmar amber
 Dictyninae indet *in* Wunderlich (2012b) Pa Rovno amber
- Argenna* Thorell, 1870a Neogene – Recent

852. <i>Argenna fossilis</i> Petrunkevitch in Palmer, 1957	Ne Mojave Desert
† Arthrodictyna Petrunkevitch, 1942	Palaeogene
853. <i>Arthrodictyna segmentata</i> Petrunkevitch, 1942*	Pa Baltic amber
† Balticocryphoeca Wunderlich, 2004v	Palaeogene
854. <i>Balticocryphoeca curvitaris</i> Wunderlich, 2004v*	Pa Baltic / Bitt. amber
† Brommellina Wunderlich, 2004v	Palaeogene
855. <i>Brommellina longungulae</i> Wunderlich, 2004v*	Pa Baltic amber
† Burmadietyna Wunderlich, 2008d	Cretaceous
856. <i>Burmadietyna pecten</i> Wunderlich, 2008d*	K Myanmar amber
† Chelicirrum Wunderlich, 2004v	Palaeogene
857. <i>Chelicirrum stridulans</i> Wunderlich, 2004v*	Pa Baltic amber
† Cryphoezaga Wunderlich, 2004v	Palaeogene
858. <i>Cryphoezaga dubia</i> Wunderlich, 2004v*	Pa Baltic amber
Dictyna Sundevall, 1833	Quaternary – Recent
859. <i>Dictyna rufa</i> Wunderlich, 2012a	Qt Madagascar copal
† Eobrommella Wunderlich, 2004v	Palaeogene
860. <i>Eobrommella scutata</i> Wunderlich, 2004v*	Pa Baltic amber
† Eocryphoeca Petrunkevitch, 1946	Palaeogene
861. <i>Eocryphoeca bitterfeldensis</i> Wunderlich, 2004v	Pa Bitterfeld amber
862. <i>Eocryphoeca electrina</i> Wunderlich, 2004v	Pa Baltic amber
863. <i>Eocryphoeca falcata</i> Wunderlich, 2004v	Pa Baltic amber
864. <i>Eocryphoeca gibbifera</i> Wunderlich, 2004v	Pa Baltic amber
865. <i>Eocryphoeca gracilipes</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
866. <i>Eocryphoeca ligula</i> Wunderlich, 2004v	Pa Baltic amber
867. <i>Eocryphoeca mammilla</i> Wunderlich, 2004v	Pa Baltic amber
868. <i>Eocryphoeca splendens</i> Wunderlich, 2004v	Pa Baltic amber
<i>Eocryphoeca</i> sp. in Wunderlich (2004v)	Pa Baltic amber
† Eocryphoecara Wunderlich, 2004v	Palaeogene
869. <i>Eocryphoecara abicera</i> Wunderlich, 2004v*	Pa Baltic amber
† Eodictyna Wunderlich, 2004v	Palaeogene
870. <i>Eodictyna communis</i> Wunderlich, 2004v*	Pa Baltic amber
† Eolathys Petrunkevitch, 1950	Palaeogene
871. <i>Eolathys debilis</i> Petrunkevitch, 1950	Pa Baltic amber
872. <i>Eolathys succini</i> Petrunkevitch, 1950*	Pa Baltic amber
† Flagelldictyna Wunderlich, 2012a	Quaternary
873. <i>Flagelldictyna copalis</i> Wunderlich, 2012a*	Qt Madagascar copal
† Gibbermastigusa Wunderlich, 2004v	Palaeogene
874. <i>Gibbermastigusa lateralis</i> Wunderlich, 2004v*	Pa Baltic amber
† Hispaniolyna Wunderlich, 1988	Neogene
875. <i>Hispaniolyna hirsuta</i> Wunderlich, 1988	Ne Dominican amber
876. <i>Hispaniolyna magna</i> Wunderlich, 1988*	Ne Dominican amber

† Mastigusa Menge in C. L. Koch & Berendt, 1854	Palaeogene
= † <i>Eotetrilus</i> Wunderlich, 1982 [<i>nomen nudum</i>]	
877. <i>Mastigusa acuminata</i> Menge in C. L. Koch & Berendt, 1854*	Pa Baltic amber
878. <i>Mastigusa arcuata</i> Wunderlich, 2004v	Pa Baltic amber
879. <i>Mastigusa bitterfeldensis</i> Wunderlich, 2004v	Pa Bitterfeld amber
880. <i>Mastigusa laticymbium</i> Wunderlich, 2004v	Pa Baltic amber
881. <i>Mastigusa magnibulbus</i> Wunderlich, 2004v	Pa Bitterfeld amber
882. <i>Mastigusa media</i> Wunderlich, 1986	Pa Baltic amber
883. <i>Mastigusa modesta</i> Wunderlich, 1986	Pa Baltic amber
884. <i>Mastigusa scutata</i> Wunderlich, 2004v	Pa Baltic amber
<i>Mastigusa</i> sp. in Wunderlich (2004v)	Pa Baltic amber
† Mizagalla Wunderlich, 2004v	Palaeogene
885. <i>Mizagalla quattuor</i> Wunderlich, 2004v*	Pa Baltic amber
886. <i>Mizagalla tuberculata</i> Wunderlich, 2004v	Pa Baltic amber
† Palaeodictyna Wunderlich, 1988	Neogene
887. <i>Palaeodictyna intermedia</i> Wunderlich, 1988	Ne Dominican amber
888. <i>Palaeodictyna longispina</i> Wunderlich, 1988	Ne Dominican amber
889. <i>Palaeodictyna singularis</i> Wunderlich, 1988	Ne Dominican amber
890. <i>Palaeodictyna spiculum</i> Wunderlich, 1988	Ne Dominican amber
891. <i>Palaeodictyna termitophila</i> Wunderlich, 1988*	Ne Dominican amber
892. <i>Palaeodictyna unispina</i> Wunderlich, 1988	Ne Dominican amber
† Palaeolathys Wunderlich, 1986	Neogene
893. <i>Palaeolathys circumductus</i> Wunderlich, 1988	Ne Dominican amber
894. <i>Palaeolathys copalis</i> Wunderlich, 1986	Qt Dominican copal
895. <i>Palaeolathys quadruplex</i> Wunderlich, 1988	Ne Dominican amber
896. <i>Palaeolathys similis</i> Wunderlich, 1988	Ne Dominican amber
897. <i>Palaeolathys spinosa</i> Wunderlich, 1986*	Ne Dominican amber
<i>Palaeolathys</i> sp. in Wunderlich (1988)	Ne Dominican amber
† Protomastigusa Wunderlich, 2004v	Palaeogene
898. <i>Protomastigusa composita</i> Wunderlich, 2004v	Pa Baltic amber
† Scopulyna Wunderlich, 2004v	Palaeogene
899. <i>Scopulyna cursor</i> Wunderlich, 2004v	Pa Baltic amber
† Succinya Wunderlich, 1988	Neogene
900. <i>Succinya longembolus</i> Wunderlich, 1988	Ne Dominican amber
901. <i>Succinya pulcher</i> Wunderlich, 1988*	Ne Dominican amber
902. <i>Succinya spinipalpus</i> Wunderlich, 1988	Ne Dominican amber
Thallumetus Simon, 1892b	Subrecent – Recent
903. <i>Thallumetus copalis</i> Wunderlich, 2004at	Qt Colombian copal
AMAUROBIIDAE Thorell, 1870a	Palaeogene – Recent
= CINIFLONIDAE Blackwall, 1841	
[partly also Dictynidae; based on a generic synonym]	

Amaurobiinae sp. <i>in</i> Wunderlich (2004 <i>u</i>)	Pa Baltic amber
PHYXELIDIDAE Lehtinen, 1967	Recent
no fossil record	
TITANOECIDAE Lehtinen, 1967	Quaternary – Recent
† <i>Copaldictyna</i> Wunderlich, 2004<i>v</i>	Quaternary
Tentative transfer by Wunderlich (2012 <i>a</i>)	
904. <i>Copaldictyna madagascariensis</i> Wunderlich, 2004 <i>v</i> *	Qt Madagascan copal
NICODAMIDAE Simon, 1898	Recent
= MEGADICTYNIDAE Lehtinen, 1967	
no fossil record	
TENGELLIDAE Dahl, 1908	Recent
no fossil record	
MITURGIDAE Simon, 1885<i>a</i>	Neogene – Recent
= CHEIRACANTHIDAE Wagner, 1887	
<i>Strotarchus</i> Simon, 1888	Neogene – Recent
= † <i>Mimeutychurus</i> Petrunkevitch, 1963 [tentative synonymy]	
905. <i>Strotarchus heidti</i> Wunderlich, 1988	Ne Dominican amber
906. <i>Strotarchus paradoxus</i> (Petrunkevitch, 1963)	Ne Chiapas amber
ANYPHAENIDAE Bertkau, 1878<i>a</i>	Palaeogene – Recent
= AMAUROBIOIDIDAE Hickman, 1949	
<i>Anyphaena</i> Sundevall, 1833	Palaeogene – Recent
907. ' <i>Anyphaena</i> ' <i>fuscata</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
<i>Anyphaenoides</i> Berland, 1913	Neogene – Recent
908. <i>Anyphaenoides bulla</i> (Wunderlich, 1988)	Ne Dominican amber
<i>Lupettiana</i> Brescovit, 1997	Neogene – Recent
909. <i>Lupettiana ligula</i> (Wunderlich, 1988)	Ne Dominican amber
<i>Wulfila</i> O. P.-Cambridge, 1895	Neogene – Recent
910. <i>Wulfila spinipes</i> Wunderlich, 1988	Ne Dominican amber
LIOCRANIDAE Simon, 1897<i>a</i>	Palaeogene – Recent
?Liocranidae <i>in</i> Wunderlich (1988)	Ne Dominican amber
<i>Apostenus</i> Westring, 1851	Palaeogene – Recent
911. <i>Apostenus arnoldorum</i> Wunderlich, 2004 <i>ag</i>	Pa Baltic amber
912. <i>Apostenus bigibber</i> Wunderlich, 2004 <i>ag</i>	Pa Baltic / Bitt. amber
913. <i>Apostenus spinimanus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
<i>Donuea</i> Strand, 1932	Quaternary – Recent
914. <i>Donuea collustrata</i> Bosselaers & Dierick, 2010 [Recent]	Qt – R Madagascar

- † ***Palaeospinisoma* Wunderlich, 2004ag** **Palaeogene**
 915. *Palaeospinisoma femoralis* Wunderlich, 2004ag* Pa Baltic amber

CLUBIONOIDEA *incertae sedis*

Wunderlich (2011d) proposed removing almost all the amber fossils from the clubionids *sensu stricto*. We follow this in part for the two genera below, but would prefer a more formal treatment before accepting all these transfers. In general the delimitation of even modern clubionids, and related forms, is problematic.

- † ***Concursator* Petrunkevitch, 1958** **Palaeogene**
 916. *Concursator nudipes* Petrunkevitch, 1958* Pa Baltic amber
- † ***Systariella* Wunderlich, 2004af** **Palaeogene**
 917. *Systariella magniocoli* Wunderlich, 2004af* Pa Baltic amber

- CLUBIONIDAE Simon, 1895** **Palaeogene – Recent**
 Clubionidae gen. et sp. *in* Nishikawa (1974) Qt Mizunami copal

- Clubiona* Latreille, 1804a** **Palaeogene – Recent**
918. *Clubiona arcana* Scudder, 1890a Pa Florissant
 919. *Clubiona attenuata* C. L. Koch & Berendt, 1854 Pa Baltic amber
 920. *Clubiona curvispinosa* Petrunkevitch, 1922 Pa Florissant
 921. *Clubiona florissantii* Petrunkevitch, 1922 Pa Florissant
 922. *Clubiona lanata* C. L. Koch & Berendt, 1854 Pa Baltic amber
 923. *Clubiona microphthalma* C. L. Koch & Berendt, 1854 Pa Baltic amber
 924. *Clubiona pubescens* C. L. Koch & Berendt, 1854 Pa Baltic amber
 925. *Clubiona sericea* C. L. Koch & Berendt, 1854 Pa Baltic amber
 926. *Clubiona tomentosa* C. L. Koch & Berendt, 1854 Pa Baltic amber

- † ***Desultor* Petrunkevitch, 1942** **Palaeogene**
 927. *Desultor depressus* Petrunkevitch, 1942 Pa Baltic amber

- Elaver* O. P.-Cambridge, 1898** **Neogene – Recent**
 928. *Elaver nutua* (Wunderlich, 1988) Ne Dominican amber

- † ***Eobumbatrix* Petrunkevitch, 1922** **Palaeogene**
 929. *Eobumbatrix latebrosa* (Scudder, 1890a)* Pa Florissant

- † ***Eodoter* Petrunkevitch, 1958** **Palaeogene**
930. *Eodoter eopala* Wunderlich, 2004af Pa Baltic amber
 931. *Eodoter lonimammillae* Wunderlich, 2012c Pa Baltic amber
 932. *Eodoter magnificus* Petrunkevitch, 1958* Pa Baltic amber
 933. *Eodoter scutatus* Wunderlich, 2011d Pa Baltic amber
 934. *?Eodoter tibialis* Wunderlich, 2011d Pa Baltic amber

- † ***Eostentatrix* Petrunkevitch, 1922** **Palaeogene**
935. *Eostentatrix cockerelli* Petrunkevitch, 1922 Pa Florissant
 936. *Eostentatrix ostentata* (Scudder, 1890a)* Pa Florissant

- † ***Eoversatrix* Petrunkevitch, 1922** **Palaeogene**
 937. *Eoversatrix eversa* (Scudder, 1890a)* Pa Florissant

- † ***Machilla* Petrunkevitch, 1958** [family uncertain] **Palaeogene**
 938. *Machilla setosa* Petrunkevitch, 1958* Pa Baltic amber
- † ***Massula* Petrunkevitch, 1942** [family uncertain] **Palaeogene**
 939. *Massula klebsi* Petrunkevitch, 1942* Pa Baltic amber
- † ***Prosocer* Petrunkevitch, 1963** **Neogene**
 940. *Prosocer mollis* Petrunkevitch, 1963* Ne Chiapas amber
- Clubionidae incertae sedis**
- † ***Chiapasona* Petrunkevitch, 1963** **Neogene**
 941. *Chiapasona defuncta* Petrunkevitch, 1963* Ne Chiapas amber
- CORINNIDAE Karsch, 1880a** **Palaeogene – Recent**
 = MYRMECIIDAE C. L. Koch, 1851 [name already used for ants]
- † ***Ablator* Petrunkevitch, 1942** **Palaeogene**
 = † *Abliquritor* Petrunkevitch, 1942
942. *Ablator biguttatus* Wunderlich, 2004ah Pa Baltic amber
 943. *Ablator curvatus* Wunderlich, 2004ah Pa Baltic amber
 944. *Ablator deminuens* Wunderlich, 2004ah Pa Baltic amber
 945. *Ablator depressus* Wunderlich, 2004ah Pa Baltic amber
 946. *Ablator duomammillae* Wunderlich, 2004ah Pa Baltic amber
 947. *Ablator felix* (Petrunkevitch, 1958) Pa Baltic amber
 948. *Ablator inevolvens* Wunderlich, 2004ah Pa Baltic amber
 949. *Ablator longus* Wunderlich, 2004ah Pa Baltic amber
 950. *Ablator nonguttatus* Wunderlich, 2004ah Pa Baltic amber
 951. *Ablator parvus* Wunderlich, 2004ah Pa Baltic amber
 952. *Ablator plumosus* (Petrunkevitch, 1950) Pa Baltic amber
 953. *Ablator robustus* Wunderlich, 2004ah Pa Baltic amber
 954. *Ablator scutatus* Wunderlich, 2004ah Pa Baltic amber
 955. *Ablator splendens* Wunderlich, 2004ah Pa Baltic amber
 956. *Ablator triguttatus* (C. L. Koch & Berendt, 1854)* Pa Baltic amber
 i. = *Philodromus microcephalus* C. L. Koch & Berendt,
 1854 Pa Baltic amber
 ii. = *Philodromus squamiger* C. L. Koch & Berendt, 1854 ..Pa Baltic amber
 iii. = *Abliqurator niger* Petrunkevitch, 1942 Pa Baltic amber
- † ***Alterphrurolithus* Wunderlich, 2004ah** **Palaeogene**
 957. *Alterphrurolithus longipes* Wunderlich, 2004ah Pa Baltic amber
- Castianeira* Keyserling, 1880b** **Neogene – Recent**
 958. *Castianeira tenebricosa* Wunderlich, 1988 Ne Dominican amber
- † ***Chemmisomma* Wunderlich, 1988** **Neogene**
 959. *Chemmisomma dubia* Wunderlich, 1988* Ne Dominican amber
- Corinna* C. L. Koch, 1842a** **Neogene – Recent**
 960. *Corinna flagelliformis* Wunderlich, 1988 Ne Dominican amber

† Cornucymbium Wunderlich, 2004ah	Palaeogene
961. <i>Cornucymbium insolens</i> Wunderlich, 2004ah*	Pa Baltic amber
† Cryptoplanus Petrunkevitch, 1958	Palaeogene
962. <i>Cryptoplanus bulbosus</i> Wunderlich, 2004ah	Pa Baltic amber
963. <i>Cryptoplanus complicatus</i> Wunderlich, 2004ah	Pa Baltic amber
964. <i>Cryptoplanus incidens</i> Wunderlich, 2004ah	Pa Baltic amber
965. <i>Cryptoplanus lanatus</i> (Petrunkevitch, 1958)	Pa Baltic amber
966. <i>Cryptoplanus paradoxus</i> Petrunkevitch, 1958*	Pa Baltic amber
967. <i>Cryptoplanus sericatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
968. <i>Cryptoplanus sinuosus</i> Wunderlich, 2004ah	Pa Baltic amber
<i>Cryptoplanus</i> sp. <i>in</i> Wunderlich (2004ah)	Pa Baltic amber
† Eomazax Petrunkevitch, 1958	Palaeogene
969. <i>Eomazax pulcher</i> Petrunkevitch, 1958*	Pa Baltic amber
Megalostrata Karsch, 1880a	Neogene – Recent
970. <i>Megalostrata grandis</i> Wunderlich, 1988	Ne Dominican amber
† Myrmecorinna Wunderlich, 2004ah	Palaeogene
971. <i>Myrmecorinna gracilis</i> Wunderlich, 2004ah*	Pa Baltic amber
† Palpiraptor Wunderlich, 2011f	Quaternary
972. <i>Palpiraptor myrmarachnoides</i> Wunderlich, 2011f*	Qt Madagascar copal
Phrurolithus C. L. Koch, 1839b	Palaeogene
973. <i>Phrurolithus extinctus</i> Petrunkevitch, 1958	Pa Baltic amber
974. <i>Phrurolithus fossilis</i> Petrunkevitch, 1958	Pa Baltic amber
975. <i>Phrurolithus ipseni</i> Petrunkevitch, 1958	Pa Baltic amber
† Protoorthobula Wunderlich, 2004ah	Palaeogene
976. <i>Protoorthobula bifida</i> Wunderlich, 2004ah*	Pa Baltic amber
977. <i>Protoorthobula deelemani</i> Wunderlich, 2004ah	Pa Baltic / Bitt. amber
Trachelas L. Koch, 1872	Neogene
978. <i>Trachelas poinari</i> Penney, 2001	Ne Dominican amber
ZODARIIDAE Thorell, 1881	Palaeogene – Recent
= CRYPTOTHELIDAE L. Koch, 1872 [younger name protected by useage]	
= † ADJUTORIDAE Petrunkevitch, 1942	
Zodariidae gen. et sp. indet 1–4 <i>in</i> Wunderlich (2004ae)	Pa Baltic amber
† Adjutor Petrunkevitch, 1942	Palaeogene
979. <i>Adjutor deformis</i> Petrunkevitch, 1958	Pa Baltic amber
980. <i>Adjutor mirabilis</i> Petrunkevitch, 1942*	Pa Baltic amber
† Admissor Petrunkevitch, 1942	Palaeogene
981. <i>Admissor aculeatus</i> Petrunkevitch, 1942*	Pa Baltic amber
† Adorator Petrunkevitch, 1942	Palaeogene
982. <i>Adorator hispidus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic / Rovno amber
i. = <i>Segestria cylindrica</i> C. L. Koch & Berendt, 1854	Pa Baltic amber

ii.	= <i>Eresus curtipes</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
iii.	= <i>Eresus monachus</i> C. L. Koch & Berendt, 1854	Pa	Baltic amber
iv.	= <i>Adorator brevipes</i> Petrunkevitch, 1942*	Pa	Baltic amber
983.	<i>Adorator samlandicus</i> Petrunkevitch, 1942	Pa	Baltic amber
†	Angusdarion Wunderlich, 2004ae		Palaeogene
984.	<i>Angusdarion humilis</i> Wunderlich, 2004ae*	Pa	Baltic amber
†	Anniculus Petrunkevitch, 1942		Palaeogene
985.	<i>Anniculus balticus</i> Petrunkevitch, 1942*	Pa	Baltic amber
†	Eocydrele Petrunkevitch, 1958		Palaeogene
986.	<i>Eocydrele mortua</i> Petrunkevitch, 1958*	Pa	Baltic amber
†	Propago Petrunkevitch, 1963		Neogene
987.	<i>Propago debilis</i> Petrunkevitch, 1963*	Ne	Chiapas amber
†	Spinizodarion Wunderlich, 2004ae		Palaeogene
988.	<i>Spinizodarion ananulum</i> Wunderlich, 2004ae*	Pa	Baltic amber
†	Zodariodamus Wunderlich 2004ae		Palaeogene
989.	<i>Zodariodamus recurvatus</i> Wunderlich 2004ae*	Pa	Baltic amber

PENESTOMIDAE Simon, 1903 **Recent**

no fossil record

† **EPHALMATORIDAE Petrunkevitch, 1950** **Palaeogene**

† **Ephalmator Petrunkevitch, 1950** **Palaeogene**

990.	<i>Ephalmator bitterfeldensis</i> Wunderlich, 2004ad	Pa	Bitterfeld amber
991.	<i>Ephalmator calidus</i> Wunderlich, 2004ad	Pa	Baltic amber
992.	<i>Ephalmator debilis</i> Wunderlich, 2004ad	Pa	Baltic amber
993.	<i>Ephalmator distinctus</i> Wunderlich, 2004ad	Pa	Baltic amber
994.	<i>Ephalmator ellwangeri</i> Wunderlich, 2004ad	Pa	Baltic amber
995.	? <i>Ephalmator eximius</i> Petrunkevitch, 1958	Pa	Baltic amber
996.	<i>Ephalmator fossilis</i> Petrunkevitch, 1950*	Pa	Baltic amber
997.	<i>Ephalmator kerneggeri</i> Wunderlich, 2004ad	Pa	Baltic amber
998.	<i>Ephalmator petrunkevitchi</i> Wunderlich, 2004ad	Pa	Baltic amber
999.	<i>Ephalmator ruthildae</i> Wunderlich, 2004ad	Pa	Baltic amber
1000.	<i>Ephalmator tredecim</i> Wunderlich, 2012c	Pa	Baltic amber
1001.	<i>Ephalmator trudis</i> Wunderlich, 2004ad	Pa	Baltic amber
1002.	<i>Ephalmator turpiculus</i> Wunderlich, 2004ad	Pa	Baltic amber
	<i>Ephalmator</i> sp. in Wunderlich (2004ad)	Pa	Baltic amber

CHUMMIDAE Jocqué, 2001 **Recent**

no fossil record

HOMALONYCHIDAE Simon, 1893 **Recent**

no fossil record

GNAPHOSOIDEA Simon, 1893	Palaeogene – Recent
AMMOXENIDAE Simon, 1893	Recent
no fossil record	
CITHAERONIDAE Simon, 1893	Recent
no fossil record	
GALLIENIELLIDAE Millot, 1947	Recent
no fossil record	
TROCHANTERIIDAE Karsch, 1879	Palaeogene – Recent
= PLATORIDAE Simon, 1890	
† <i>Eotrochanteria</i> Wunderlich, 2004am	Palaeogene
1003. <i>Eotrochanteria kruegeri</i> Wunderlich, 2004am*	Pa Baltic amber
† <i>Sosybius</i> C. L. Koch & Berendt, 1854	Palaeogene
= † <i>Adamator</i> Petrunkevitch, 1942	
= † <i>Adjunctor</i> Petrunkevitch, 1942	
= † <i>Adulatrix</i> Petrunkevitch, 1942	
1004. <i>Sosybius berendti</i> Wunderlich, 2004am	Pa Baltic amber
1005. <i>Sosybius decumana</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1006. <i>Sosybius falcatus</i> Wunderlich, 2004am	Pa Baltic amber
1007. <i>Sosybius fusca</i> (Petrunkevitch, 1942)	Pa Baltic amber
1008. <i>Sosybius kochi</i> Wunderlich, 2004am	Pa Baltic amber
1009. <i>Sosybius lateralis</i> Wunderlich, 2004am	Pa Baltic amber
1010. <i>Sosybius longipes</i> Wunderlich, 2004am	Pa Baltic amber
1011. <i>Sosybius major</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1012. <i>Sosybius minor</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
1013. <i>Sosybius mizgirisi</i> Wunderlich, 2004am	Pa Baltic amber
1014. <i>Sosybius parva</i> (Petrunkevitch, 1942)	Pa Baltic amber
1015. <i>Sosybius perniciosus</i> Wunderlich, 2004am	Pa Baltic amber
1016. <i>Sosybius rufa</i> (Petrunkevitch, 1942)	Pa Baltic amber
1017. <i>Sosybius similis</i> Petrunkevitch, 1942	Pa Baltic amber
1018. <i>Sosybius succineus</i> (Petrunkevitch, 1942)	Pa Baltic amber
1019. <i>Sosybius tibialis</i> Wunderlich, 2004am	Pa Baltic amber
1020. <i>Sosybius unispinosus</i> Wunderlich, 2004am	Pa Baltic amber
<i>Sosybius</i> sp. <i>in</i> Wunderlich (2004am, ar)	Pa Baltic / Rovno amber
† <i>Thereola</i> Petrunkevitch, 1955	Palaeogene
= † <i>Therea</i> Koch & Berendt, 1854 [preoccupied]	
1021. <i>Thereola petiolata</i> (C. L. Koch & Berendt, 1854)* [♀ = ? <i>Dasuminia</i> sp. according to Wunderlich 2004b]	Pa Baltic amber
1022. <i>Thereola pubescens</i> (Menge <i>in</i> C. L. Koch & Berendt, 1854)	Pa Baltic amber

† <i>Trochanteridromulus</i> Wunderlich, 2004am	Palaeogene
1023. <i>Trochanteridromulus glabripes</i> Wunderlich, 2004am*	Pa Baltic amber
† <i>Trochanteridromus</i> Wunderlich, 2004am	Palaeogene
1024. <i>Trochanteridromus scutatus</i> Wunderlich, 2004am*	Pa Baltic amber
† <i>Veterator</i> Petrunkevitch, 1963	Neogene
1025. <i>Veterator angustus</i> Wunderlich, 1988	Ne Dominican amber
1026. <i>Veterator ascutum</i> Wunderlich, 1988	Ne Dominican amber
1027. <i>Veterator extinctus</i> Petrunkevitch, 1963*	Ne Chiapas amber
1028. <i>Veterator incompletus</i> Wunderlich, 1982	Ne Dominican amber
1029. <i>Veterator longipes</i> Wunderlich, 1988	Ne Dominican amber
1030. <i>Veterator loricatus</i> Wunderlich, 1988	Ne Dominican amber
1031. <i>Veterator porrectus</i> Wunderlich, 1988	Ne Dominican amber
1032. <i>Veterator viduus</i> Wunderlich, 1988	Ne Dominican amber
<i>Veterator</i> sp. 1–2 in Wunderlich (1988)	Ne Dominican amber
LAMPONIDAE Simon, 1893	Recent
no fossil record	
PRODIDOMIDAE Simon, 1884a	Quaternary – Recent
= MILTIIDAE Thorell, 1873 [based on a generic synonym]	
<i>Prodidomus</i> Hentz, 1847	Quaternary – Recent
1033. <i>Prodidomus madagascariensis</i> Wunderlich, 2011c	Qt Madagascar copal
GNAPHOSIDAE Pocock, 1898	?Cretaceous – Recent
= DRASSIDAE Sundevall, 1833 [based on a generic synonym]	
† <i>Captrix</i> Petrunkevitch, 1942	Palaeogene
1034. <i>Captrix lineata</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
<i>Drassodes</i> Westring, 1851	Palaeogene – Recent
1035. <i>Drassodes cupreus</i> (Blackwall, 1834a) [Recent]	Qt England
1036. ? <i>Drassodes femurus</i> Lin, Zhang & Wang, 1989	Ne Shanwang
1037. ? <i>Drassodes sextii</i> Berland, 1939	Pa Aix-en-Provence
† <i>Drassyllinus</i> Wunderlich, 1988	Neogene
1038. <i>Drassyllinus aliter</i> Wunderlich, 1988*	Ne Dominican amber
† <i>Eognaphosops</i> Wunderlich, 2011b	Palaeogene
1039. <i>Eognaphosops cryptoplanoides</i> Wunderlich 2011b*	Pa Baltic amber
† <i>Eomactator</i> Petrunkevitch, 1958	Palaeogene
1040. <i>Eomactator hamatus</i> Wunderlich, 2011b	Pa Baltic amber
1041. <i>Eomactator hirsutipes</i> Wunderlich, 2011b	Pa Baltic amber
1042. <i>Eomactator mactatus</i> Petrunkevitch, 1958*	Pa Baltic amber
1043. <i>Eomactator obscurior</i> Wunderlich, 2011b	Pa Baltic amber
Gnaphosa Latreille, 1804a	?Cretaceous – Recent
1044. <i>Gnaphosa affinis</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber

i. = <i>Philodromus dubius</i> C. L. Koch & Berendt, 1854	
1045. <i>Gnaphosa ambigua</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1046. <i>Gnaphosa liaoningensis</i> Chang, 2004	
[generic assignment unreliable!]	K Jehol biota
Micaria Westring, 1851	Palaeogene – Recent
1047. <i>Micaria procera</i> C. L. Koch & Berendt, 1954	Pa Baltic amber
1048. <i>Micaria tenella</i> Heer, 1865	Ne Öhningen
† Palaeodrassus Petrunkevitch, 1922	Palaeogene
1049. <i>Palaeodrassus cockerelli</i> Petrunkevitch, 1922	Pa Florissant
1050. <i>Palaeodrassus florissanti</i> Petrunkevitch, 1922	Pa Florissant
1051. <i>Palaeodrassus hesternus</i> (Scudder, 1890a)	Pa Florissant
1052. <i>Palaeodrassus ingenuus</i> (Scudder, 1890a)*	Pa Florissant
1053. <i>Palaeodrassus interitus</i> (Scudder, 1890a)	Pa Florissant
Scopoides Platnick, 1989	Palaeogene – Recent
1054. <i>Scopoides dominicanus</i> Wunderlich, 2011g	Ne Dominican amber
Zelotes Gistel, 1848	Palaeogene
1055. <i>Zelotes concinna</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1056. <i>Zelotes mundula</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Melanophora nobilis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1057. <i>Zelotes regalis</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
† Zelotetis Wunderlich, 2011b	Palaeogene
1058. <i>Zelotetis calefacta</i> Wunderlich, 2011b	Pa Baltic amber
SELENOPIDAE Simon, 1897a	Palaeogene – Recent
† Garcorops Corronca, 2003	Quaternary – Recent
1059. <i>Garcorops jadis</i> Bosselaers, 2004	Qt Madagascar copal
i. = ? <i>Anyphops cortex</i> Wunderlich, 2004as	Qt Madagascar copal
Selenops Latreille, 1819	Palaeogene – Recent
1060. <i>Selenops benoiti</i> Wunderlich, 2004as	Qt Madagascar copal
1061. <i>Selenops beynai</i> Schawaller, 1984	Ne Dominican amber
1062. <i>Selenops dominicanus</i> Wunderlich, 2004an	Ne Dominican amber
<i>Selenops</i> sp. in Wunderlich (1988)	Ne Dominican amber
<i>Selenops</i> sp. in García-Villafuerte (2006b)	Ne Chiapas amber
<i>Selenops</i> sp. in Penney (2007)	Pa Le Quesnoy amber
SPARASSIDAE Bertkau, 1872	Palaeogene – Recent
= HETEROPODIDAE Thorell, 1873	
= MICROMMATIDAE Bertkau, 1878a	
= EUSPARASSIDAE Järvi, 1912	
Sparassidae sp. 1–2 in (Wunderlich 2008c)	Pa Baltic amber
† Caduceator Petrunkevitch, 1942	Palaeogene
1063. <i>Caduceator minutus</i> Petrunkevitch, 1942*	Pa Baltic amber

1064. *Caduceator quadrimaculatus* Petrunkevitch, 1950 Pa Baltic amber
- † ***Collacteus* Petrunkevitch, 1942** **Palaeogene**
1065. *Collacteus captivus* Petrunkevitch, 1942* Pa Baltic amber
- † ***Eostaianus* Petrunkevitch, 1950** **Palaeogene**
1066. *Eostaianus succini* Petrunkevitch, 1950* Pa Baltic amber
- † ***Eostasina* Petrunkevitch, 1942** **Palaeogene**
1067. *Eostasina aculeata* Petrunkevitch, 1942* Pa Baltic amber
- Eusparassus* Simon 1903** **Palaeogene – Recent**
1068. *Eusparassus crassipes* (C. L. Koch & Berendt, 1854) Pa Baltic amber
- Heteropoda* Latreille, 1804a** **Palaeogene – Recent**
- = † *Retina* Hong, 1985
1069. *Heteropoda rpbusta* [sic] (Hong, 1985) Ne Shanwang
- [NB: as '*H. robusta*' this would be a junior homonym of a living species.]
- Pseudosparianthis* Simon, 1887** **Neogene – Recent**
1070. *Pseudosparianthis pfeifferi* (Wunderlich, 1988) Ne Dominican amber
- Zachria* L. Koch, 1875** **Palaeogene – Recent**
- [NB: An Australian genus; Wunderlich (2012c) regarded at least *Z. desiderabilis* as gen. indet.]
1071. *Zachria desiderabilis* Petrunkevitch, 1950 Pa Baltic amber
1072. *Zachria peculiata* Petrunkevitch, 1946 Pa Baltic amber
1073. *Zachria restincta* Petrunkevitch, 1958 Pa Baltic amber
- PHILODROMIDAE Thorell, 1870a** **Cretaceous – Recent**
- Philodromidae sp. *in* Wunderlich (1988) Ne Dominican amber
- Philodromidae sp. *in* Wunderlich (2004ae) Ne Baltic amber
- † ***Cretadromus* Cheng, Shen & Gao, 2009** **Cretaceous**
1074. *Cretadromus liaoningensis* Cheng, Shen & Gao, 2009 K Liaoning Province
- [NB: Wunderlich (2012d) suggested this could be a Theridosomatidae]
- † ***Eoathanatus* Petrunkevitch, 1950** **Palaeogene – Recent**
1075. *Eoathanatus diritatis* Petrunkevitch, 1950* Pa Baltic amber
- THOMISIDAE Sundevall, 1833** **Palaeogene – Recent**
- = APHANTOCHILIDAE Thorell, 1873
- = MISUMENIDAE Thorell, 1887
- = STIPHROPODIDAE Simon, 1895
- = XYSTICIDAE Dahl, 1912
- = BORBOROPACTIDAE Wunderlich, 2004ao
- Thomisidae gen. et sp. *in* Nishikawa (1974) Qt Mizunami copal
- Thomisidae gen. et sp. *in* Bottali (1975) Qt Italy
- Thomisidae gen. et sp. *in* Schawaller (1982d) Ne Willershausen
- Thomisidae gen. et sp. *in* Wunderlich (1988) Ne Dominican amber
- Thomisidae gen. et sp. 1–2 *in* Wunderlich (2004ap) Pa Baltic amber
- Thomisidae gen. et sp. *in* Garcíá-Villafuerte (2006b) Ne Chiapas amber

Coriarachne Thorell, 1870b	Quaternary – Recent
<i>Coriarachne</i> sp. in Cutler (1970)	Qt Wyoming
† Ecotona Lin, Zhang & Wang, 1989 [ex Araneidae]	Neogene
1076. <i>Ecotona brunnea</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
1077. <i>Ecotona pilulifera</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
1078. <i>Ecotona transipeda</i> Lin, Zhang & Wang, 1989*	Ne Shanwang
† Facundia Petrunkevitch, 1942	Palaeogene
1079. <i>Facundia clara</i> Petrunkevitch, 1942*	Pa Baltic amber
† Fiducia Petrunkevitch, 1950	Palaeogene
1080. <i>Fiducia tenuipes</i> Petrunkevitch, 1950*	Pa Baltic amber
† Filiolella Petrunkevitch, 1955a	Palaeogene
= † <i>Filiola</i> Petrunkevitch, 1942 [preoccupied]	
1081. <i>Filiolella argentata</i> (Petrunkevitch, 1942)*	Pa Baltic amber
† Heterotmarus Wunderlich, 1988	Neogene
1082. <i>Heterotmarus altus</i> Wunderlich, 1988*	Ne Dominican amber
† Komisumena Ono, 1981	Neogene
1083. <i>Komisumena rosae</i> Ono, 1981*	Ne Dominican amber
† Miothomismus Zhang, Sun & Zhang, 1994	Neogene
1084. <i>Miothomismus subnudus</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
1085. <i>Miothomismus sylvaticus</i> Zhang, Sun & Zhang, 1994*	Ne Shanwang
Misumena Latreille, 1804a	Palaeogene – Recent
1086. <i>Misumena samlandica</i> Petrunkevitch, 1942	Pa Baltic amber
† Palaeoxysticus Wunderlich, 1985	Neogene
1087. <i>Palaeoxysticus extinctus</i> Wunderlich, 1985	Ne Randecker Maar
† Parvulus Zhang, Sun & Zhang, 1994	Neogene
1088. <i>Parvulus latissimus</i> Zhang, Sun & Zhang, 1994*	Ne Shanwang
† Succinaenigma Wunderlich, 2004ap	Palaeogene
1089. <i>Succinaenigma raptor</i> Wunderlich, 2004ap*	Pa Baltic amber
† Succiniraptor Wunderlich, 2004ao	Palaeogene
1090. <i>Succiniraptor radiatus</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
i. = <i>Succiniraptor paradoxus</i> Wunderlich, 2004ao*	Pa Baltic amber
Synema Simon, 1864	Palaeogene – Recent
1091. <i>Synema enigmaticum</i> Berland, 1939	Pa Aix-en-Provence
† Syphax C. L. Koch & Berendt, 1854	Palaeogene
1092. <i>Syphax asper</i> Petrunkevitch, 1950	Pa Baltic amber
1093. <i>Syphax crassipes</i> Petrunkevitch, 1942	Pa Baltic amber
1094. <i>Syphax fuliginosus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1095. <i>Syphax gracilis</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1096. <i>Syphax megacephalus</i> C. L. Koch & Berendt, 1854*	Pa Baltic amber
1097. <i>Syphax thoracicus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† Thomisiraptor Wunderlich, 2004ap	Palaeogene

1098. <i>Thomisiraptor liedtkei</i> Wunderlich, 2004ap*	Pa Baltic amber
Thomisus Walckenaer, 1805	Palaeogene – Recent
1099. <i>Thomisus defossus</i> Scudder, 1890a	Pa Florissant
1100. <i>Thomisus disjunctus</i> Scudder, 1890a	Pa Florissant
1101. <i>Thomisus lividus</i> Heer, 1865	Ne Öhningen
1102. <i>Thomisus resutus</i> Scudder, 1890a	Pa Florissant
1103. <i>Thomisus sulzeri</i> Heer, 1865	Ne Öhningen
Xysticus C. L. Koch, 1835	Palaeogene – Recent
1104. ? <i>Xysticus annulipes</i> Bertkau, 1878b	Ne Rott, Germany
1105. <i>Xysticus archaeopalpus</i> Leech & Matthews, 1971	Ne Alaska
1106. <i>Xysticus oeningensis</i> (Heer, 1865)	Ne Öhningen
<i>Xysticus</i> sp. in Protescu (1937)	Pa Romanian amber
SALTICIDAE Blackwall, 1841	Palaeogene – Recent
= ATTIDAE Sundevall, 1833 [based on a generic synonym]	
= LYSSOMANIDAE Peckham & Wheeler, 1889	
Salticidae gen. et sp. in Schawaller (1982d)	Ne Willershausen
† Almolinus Petrunkevitch, 1958	Palaeogene
1107. <i>Almolinus bitterfeldensis</i> Wunderlich, 2004aq	Pa Bitterfeld amber
1108. <i>Almolinus clarus</i> Petrunkevitch, 1958*	Pa Baltic amber
1109. <i>Almolinus ligula</i> Wunderlich, 2004aq	Pa Baltic amber
? <i>Almolinus</i> sp. in Wunderlich (2004aq)	Pa Baltic amber
† Attoides Brongniart, 1877	Palaeogene
1110. <i>Attoides eresiformis</i> Brongniart, 1877	Pa Aix-en-Provence
† Calilinus Wunderlich, 2004aq	Palaeogene
1111. <i>Calilinus fleissneri</i> Wunderlich, 2004aq*	Pa Baltic amber
† Cenattus Petrunkevitch, 1942	Palaeogene
1112. <i>Cenattus exophthalmicus</i> Petrunkevitch, 1942*	Pa Baltic amber
Corythalia C. L. Koch, 1851	Neogene – Recent
1113. <i>Corythalia ocululiter</i> Wunderlich, 1988	Ne Dominican amber
1114. <i>Corythalia pilosa</i> Wunderlich, 1982	Ne Dominican amber
1115. <i>Corythalia scissa</i> Wunderlich, 1988	Ne Dominican amber
† Descangeles Wunderlich, 1988	Neogene
1116. <i>Descangeles pygmaeus</i> Wunderlich, 1988*	Ne Dominican amber
<i>Descangeles</i> sp. 1–2 in Wunderlich (1988)	Ne Dominican amber
Descanso Peckham & Peckham, 1892	Neogene – Recent
<i>Descanso</i> sp. in Wunderlich (1988)	Ne Dominican amber
† Distanilinus Wunderlich, 2004aq	Palaeogene
1117. <i>Distanilinus filum</i> Wunderlich, 2004aq	Pa Baltic amber
1118. <i>Distanilinus nutus</i> Wunderlich, 2004aq*	Pa Baltic amber
1119. <i>Distanilinus paranutus</i> Wunderlich, 2004aq	Pa Baltic amber

1120. <i>Distanilinus pernutus</i> Wunderlich, 2004aq	Pa Baltic amber
† Eoattopsis Gourret, 1887	Palaeogene
1121. <i>Eoattopsis hirsutus</i> Gourret, 1887*	Pa Aix-en-Provence
† Eolinus Petrunkevitch, 1942	Palaeogene
1122. <i>Eolinus balticus</i> Žabka, 1988	Pa Baltic amber
1123. <i>Eolinus fungus</i> Wunderlich, 2004aq	Pa Baltic amber
1124. <i>Eolinus insuriens</i> Wunderlich, 2004aq	Pa Baltic amber
1125. <i>Eolinus prominens</i> Wunderlich, 2004aq	Pa Baltic amber
1126. <i>Eolinus samlandica</i> Wunderlich, 2004aq	Pa Baltic amber
1127. <i>Eolinus succineus</i> Petrunkevitch, 1942*	Pa Baltic amber
1128. <i>Eolinus theryi</i> Petrunkevitch, 1942	Pa Baltic amber
1129. <i>Eolinus theryoides</i> Wunderlich, 2004aq	Pa Baltic amber
1130. <i>Eolinus tystschenkoi</i> Proszynski & Žabka, 1980	Pa Baltic amber
1131. <i>Eolinus vates</i> Wunderlich, 2004aq	Pa Baltic amber
<i>Eolinus</i> sp. in Wunderlich (2004aq)	Pa Baltic amber
Euophrys C. L. Koch, 1834	Palaeogene – Recent
1132. <i>Euophrys gibberula</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1133. <i>Euophrys randeckensis</i> Schawaller & Ono, 1979	Ne Randecker Maar
† Evagoratus Zhang, Sun & Zhang, 1994	Neogene
1134. <i>Evagoratus longicruris</i> Zhang, Sun & Zhang, 1994	Ne Shanwang
† Gorgopsidis Wunderlich, 2004aq	Palaeogene
1135. <i>Gorgopsidis bechlyi</i> Wunderlich, 2004aq*	Pa Baltic amber
† Gorgopsina Petrunkevitch, 1955a	Palaeogene
1136. <i>Gorgopsina amabilis</i> Wunderlich, 2004aq	Pa Baltic amber
1137. <i>Gorgopsina constricta</i> Wunderlich, 2004aq	Pa Baltic amber
1138. <i>Gorgopsina expandens</i> Wunderlich, 2004aq	Pa Baltic amber
1139. ' <i>Gorgopsina</i> ' <i>fasciata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1140. <i>Gorgopsina flexuosa</i> Wunderlich, 2004aq	Pa Baltic amber
1141. <i>Gorgopsina formosa</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1142. <i>Gorgopsina fractura</i> Wunderlich, 2004ar	Pa Rovno amber
1143. <i>Gorgopsina frenata</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
1144. <i>Gorgopsina inclusa</i> Wunderlich, 2004aq	Pa Baltic amber
1145. <i>Gorgopsina jucunda</i> (Petrunkevitch, 1942)	Pa Baltic amber
1146. <i>Gorgopsina marginata</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1147. <i>Gorgopsina melanocephala</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1148. <i>Gorgopsina naumanni</i> Giebel, 1856	Pa Baltic amber
1149. <i>Gorgopsina paulula</i> (C. L. Koch & Berendt, 1854)	Pa Baltic amber
1150. <i>Gorgopsina rectangularis</i> Wunderlich, 2011h	Pa Baltic amber
1151. <i>Gorgopsina speciosa</i> Wunderlich, 2004aq	Pa Baltic amber
Heliophanus C. L. Koch, 1833	Palaeogene – Recent
1152. <i>Heliophanus extinctus</i> Berland, 1939	Pa Aix-en-Provence

<i>Hyllus</i> C. L. Koch, 1846	Quaternary – Recent
= † <i>Parevophrys</i> Petrunkevitch, 1942	
1153. <i>Hyllus succini</i> (Petrunkevitch, 1942)	Qt Copal
Originally described as Baltic amber	
<i>Lyssomanes</i> Hentz, 1845	Neogene – Recent
1154. <i>Lyssomanes pristinus</i> Wunderlich, 1986	Ne Dominican amber
i. = <i>Lyssomanes galianoae</i> Reiskind, 1989	Ne Dominican amber
1155. <i>Lyssomanes pulcher</i> Wunderlich, 1988	Ne Dominican amber
<i>Maevia</i> C. L. Koch, 1846	?Neogene – Recent
? <i>Maevia</i> sp. in Riquelme & Hill (2013)	Ne Chiapas amber
† <i>Microlinus</i> Wunderlich, 2004aq	Palaeogene
1156. <i>Microlinus calidus</i> Wunderlich, 2004aq	Pa Baltic amber
1157. <i>Microlinus folium</i> Wunderlich, 2004aq*	Pa Baltic amber
<i>Myrmarachne</i> MacLeay, 1839	Quaternary – Recent
= † <i>Entomocephalus</i> Holl, 1829 [suppressed; see ICZN Opinion 2258]	
1158. <i>Myrmarachne formicoides</i> (Holl, 1829)	?Qt Copal [?not amber]
<i>Neon</i> Simon, 1876a	Quaternary – Recent
1159. <i>Neon ?reticulatus</i> (Blackwall, 1853) [Recent]	Qt England
† <i>Paralinus</i> Petrunkevitch, 1942	Palaeogene
1160. <i>Paralinus crosbyi</i> Petrunkevitch, 1942*	Pa Baltic amber
† <i>Pensacolatus</i> Wunderlich, 1988	Neogene
1161. <i>Pensacolatus coxalis</i> Wunderlich, 1988*	Ne Dominican amber
1162. <i>Pensacolatus spinipes</i> Wunderlich, 1988	Ne Dominican amber
1163. ? <i>Pensacolatus tibialis</i> Wunderlich, 2004aq	Ne Dominican amber
<i>Pensacolatus</i> sp. in Wunderlich (1988)	Ne Dominican amber
<i>Phidippus</i> C. L. Koch, 1846	Palaeogene
1164. <i>Phidippus impressus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
1165. <i>Phidippus pusillus</i> C. L. Koch & Berendt, 1854	Pa Baltic amber
† <i>Phlegrata</i> Wunderlich, 1988	Neogene
1166. <i>Phlegrata pala</i> Wunderlich, 1988*	Ne Dominican amber
† <i>Prolinus</i> Petrunkevitch, 1958	Palaeogene
1167. <i>Prolinus fossilis</i> Petrunkevitch, 1958*	Pa Baltic amber
<i>Sarinda</i> Peckham & Peckham, 1892	Neogene – Recent
? <i>Sarinda</i> sp. in Wunderlich (2004aq)	Ne Dominican amber
† <i>Steneattus</i> Bronn, 1856	Palaeogene
= † <i>Leda</i> C. L. Koch & Berendt, 1854 [preoccupied]	
1168. <i>Steneattus promissa</i> (C. L. Koch & Berendt, 1854)*	Pa Baltic amber
<i>Thiodina</i> Simon, 1900	Neogene
1169. <i>Thiodina beugelorum</i> Wolff, 1990	Ne Dominican amber
Araneomorphae incertae sedis	
† <i>Elvina</i> Thorell, 1870b	Neogene

1170. *Elvina antiqua* (von Heyden, 1859) Ne Linz am Rhein

Araneae *incerate sedis*

- Araneae gen. et sp. nov. *in* Ansorge (2003) J Grimmen, Germany
- † ***Amphiclotho* Gourret, 1887** **Palaeogene**
1171. *Amphiclotho breviuscula* Gourret, 1887* Pa Aix-en-Provence
- † ***Amphithomismus* Gourret, 1887** **Palaeogene**
1172. *Amphithomismus barbatus* Gourret, 1887* Pa Aix-en-Provence
- † ***Atocatle* Feldmann, Vega, Applegate & Bishop, 1998** [really a spider?]..... **Cretaceous**
1173. *Atocatle ranulfoi* Feldmann, Vega, Applegate & Bishop, 1998* K Puebla, México
- † ***Cercidiella* Gourret, 1887** **Palaeogene**
1174. *Cercidiella aquisextana* Gourret, 1887* Pa Aix-en-Provence
- † ***Clubionella* Gourret, 1887** **Palaeogene**
1175. *Clubionella antiqua* Gourret, 1887* Pa Aix-en-Provence
- † ***Eresoides* Gourret, 1887** **Palaeogene**
1176. *Eresoides orbicularis* Gourret, 1887* Pa Aix-en-Provence
- † ***Hersilioides* Gourret, 1887** **Palaeogene**
1177. *Hersilioides thanatiformis* Gourret, 1887* Pa Aix-en-Provence
- † ***Opisthophylax* Menge, 1856** **Palaeogene**
1178. *Opisthophylax exarata* Menge, 1856* Pa Baltic amber
- † ***Prodysdera* Gourret, 1887** **Palaeogene**
1179. *Prodysdera intermedia* Gourret, 1887* Pa Aix-en-Provence
- † ***Protochersis* Gourret, 1887** **Palaeogene**
1180. *Protochersis spinosus* Gourret, 1887* Pa Aix-en-Provence
- † ***Protolachesis* Gourret, 1887** **Palaeogene**
1181. *Protolachesis annulata* Gourret, 1887* Pa Aix-en-Provence
- † ***Paralycosa* Dunlop & Jekel, 2009** **Palaeogene**
- = † *Protolycosa* Gourret, 1887 [preoccupied]
1182. *Paralycosa attiformis* (Gourret, 1887)* Pa Aix-en-Provence
- † ***Pseudothomismus* Gourret, 1887** **Palaeogene**
1183. *Pseudothomismus articulatus* Gourret, 1887* Pa Aix-en-Provence
- † ***Schellenbergia* Heer, 1865** **Neogene**
1184. *Schellenbergia rotundata* Heer, 1865* Ne Öhningen
- † ***Timeropus* Thorell, 1891** **Palaeogene**
- = † *Lycosoides* Gourret, 1887 [preoccupied]
1185. *Timeropus hersiliformis* (Gourret, 1887)* Pa Aix-en-Provence

NOMINA DUBIA

***Amaurobius* C. L. Koch, 1837** [no currently valid fossil species]

1. *Amaurobius faustus* C. L. Koch & Berendt, 1854 Pa Baltic amber
2. *Amaurobius rimosus* C. L. Koch & Berendt, 1854 Pa Baltic amber

- Auximus Simon, 1892** [now *Lathys* Simon, 1884: Dictynidae; no currently valid fossil species]
3. *Auximus fossilis* Petrunkevitch, 1950 Pa Baltic amber
4. *Auximus succini* Petrunkevitch, 1942 Pa Baltic amber
- † **Clythia C. L. Koch & Berendt, 1854 (*nomen dubium*)** **Palaeogene**
5. *Clythia alma* C. L. Koch & Berendt, 1854* Pa Baltic amber
- † **Corynitoides Dunlop & Jekel, 2009 (*nomen dubium*)** **Palaeogene**
 = † *Corynitis* Menge in C. L. Koch & Berendt, 1854 [preoccupied]
6. *Corynitoides spinosa* (Menge in C. L. Koch & Berendt, 1854)* Pa Baltic amber
7. *Corynitoides undulata* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
- † **Eocryphoeca Petrunkevitch, 1958** [also contains valid fossil species]
8. *Eocryphoeca distincta* Petrunkevitch, 1950 Pa Baltic amber
9. *Eocryphoeca fossilis* (Petrunkevitch, 1942) Pa Baltic amber
- † **Eometa Petrunkevitch, 1958** [also contains valid fossil species]
10. *Eometa aberrans* Petrunkevitch, 1958 Pa Baltic amber
11. *Eometa robusta* Petrunkevitch, 1958 Pa Baltic amber
- Ero C. L. Koch 1836** [also contains valid fossil species]
12. *Ero setulosa* C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Fictotama Petrunkevitch, 1963 (*nomen dubium*)** **Palaeogene**
13. *Fictotama extincta* Petrunkevitch, 1963* Ne Chiapas amber
- † **Memoratrix Petrunkevitch, 1942 (*nomen dubium*)** **Palaeogene**
 NB: Regarded by Wunderlich (2004p) as a possible pimoid or linyphiid
14. *Memoratrix rydei* Petrunkevitch, 1942 Pa Baltic amber
- † **Mimetarchaea Eskov, 1992** **Palaeogene**
15. *Mimetarchaea gintaras* Eskov, 1992* Pa Baltic amber
 NB: Name based on a subadult male
- † **Miropholcus Petrunkevitch, 1942 (*nomen dubium*)** **Palaeogene**
 = † *Micropholcus* Petrunkevitch, 1942 [*lapsus*]
16. *Miropholcus heteropus* Petrunkevitch, 1942* Pa Baltic amber
- † **Perturbator Petrunkevitch, 1971 (*nomen dubium*)** **Neogene**
17. *Perturbator corniger* Petrunkevitch, 1971* Ne Chiapas amber
- † **Phalangopus Menge in C. L. Koch & Berendt, 1854 (*nomen dubium*)** **Palaeogene**
18. *Phalangopus subtilis* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- † **Praeoarces Wunderlich, 2004q** **Palaeogene**
19. *Praeoarces exitus* Wunderlich, 2004q* Pa Baltic amber
- Segestria Latreille, 1804** [also contains valid fossil species]
20. *Segestria elongata* C. L. Koch & Berendt, 1854 Pa Baltic amber
21. *Segestria nana* C. L. Koch & Berendt, 1854 Pa Baltic amber

NOMINA NUDA

Amaurobius C. L. Koch, 1837 [no currently valid fossil species]

1. *Amaurobius spinimanus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

- † **Anatone Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
2. *Anatone hirsuta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 3. *Anatone marginata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 4. *Anatone spinipes* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Aranea Clerck, 1757** [now *Araneus* Clerck, 1757; which also contains valid fossil species]
5. *Aranea fossilis* Keferstein, 1834 Pa Aix-en-Provence
- Archaea C. L. Koch & Berendt, 1854** [also contains valid fossil species]
6. *Archaea incomta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 7. *Archaea sphinx* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Athera Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
8. *Athera exilis* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Attus Walckenaer, 1805** [now *Salticus* Latreille, 1804; no currently valid fossil species]
9. *Attus fossilis* Walckenaer, 1837 Pa Baltic amber
- Clubiona Latreille, 1804** [also contains valid fossil species]
10. *Clubiona eseri* Heer, 1865 Ne Öhningen
 11. *Clubiona latifrons* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 12. *Clubiona parvula* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 13. *Clubiona pilosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Clythia C. L. Koch & Berendt, 1854** [also contains a *nomen dubium* fossil species]
14. *Clythia funestra* Koch & Berendt, 1854 Pa Baltic amber
 15. *Clythia gracilentata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 16. *Clythia leptocarena* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Dielacata Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
17. *Dielacata superba* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Drassus Walckenaer, 1805** [now *Gnaphosa* Latreille, 1804; which also contains valid fossil species]
18. *Drassus oblongus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Dysdera Latreille, 1804** [also contains valid fossil species]
19. *Dysdera hippopodium* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 20. *Dysdera glabrata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 21. *Dysdera scobiculata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
 22. *Dysdera tenera* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Eolinus Petrunkevitch, 1942** [also contains valid fossil species]
23. *Eolinus bitterfeldensis* Wunderlich, 2004aq Pa Baltic amber
 24. *Eolinus tystschenkoides* Wunderlich, 2004aq Pa Baltic amber
- Epeira Walckenaer, 1805** [now *Araneus* Clerck, 1757; which also contains valid fossil species]
25. *Epeira eocaenica* Giebel, 1856 Pa Baltic amber
 26. *Epeira eocena* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Epeiridion Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
27. *Epeiridion femoratum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Erithus Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
28. *Erithus applanatus* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Ero C. L. Koch & Berendt, 1836** [also contains valid fossil species]

29. *Ero coronata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
30. *Ero exculpta* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
31. *Ero sphaerica* C. L. Koch & Berendt, 1854 Pa Baltic amber
32. *Ero quadripunctata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Eyukselus Özdikmen, 2007 (*nomen nudum*)** **Palaeogene**
 = † *Propetes* Menge, 1854 [preoccupied]
33. *Eyukselus argutus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
34. *Eyukselus felinus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
35. *Eyukselus griseus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
36. *Eyukselus latifrons* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
37. *Eyukselus pumilus* (Menge in C. L. Koch & Berendt, 1854) Pa Baltic amber
- Gea C. L. Koch, 1843** [also contains valid fossil species]
38. *Gea pubescens* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Heteromma Menge, 1856 (*nomen nudum*)** **Palaeogene**
39. *Heteromma intersecta* Menge, 1856* Pa Baltic amber
- † **Idmonia Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
40. *Idmonia virginea* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Melanophora C. L. Koch, 1833** [now *Zelotes* Gistel, 1848; which also contains valid fossil species]
41. *Melanophora lepida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
42. *Melanophora nitida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Micaria Westring, 1851** [also contains valid fossil species]
43. *Micaria ovata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
44. *Micaria squamata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
45. *Micaria tenuis* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Micryphantes C. L. Koch, 1833** [also contains valid fossil species]
46. *Micryphantes globulus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
47. *Micryphantes turritus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Mizalia C. L. Koch & Berendt, 1854** [also contains valid fossil species]
48. *Mizalia truncata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Ocia Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
49. *Ocia hirsuta* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Ocypete C. L. Koch, 1836** [now *Heteropoda* Latreille, 1804; which also contains valid fossil species]
50. *Ocypete angustifrons* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
51. *Ocypete marginata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Onca Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
52. *Onca lepida* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
53. *Onca pumila* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- Philodromus Walckenaer, 1826** [also contains valid fossil species]
54. *Philodromus griseus* Menge, 1856 Pa Baltic amber
55. *Philodromus marginatus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
56. *Philodromus reptans* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
57. *Philodromus redogradus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

58. *Philodromus spinipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Pythonissa C. L. Koch, 1837** [now *Gnaphosa* Latreille, 1804; which also contains valid fossil species]
59. *Pythonissa bipunctata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
60. *Pythonissa discophora* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
61. *Pythonissa glabra* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
62. *Pythonissa villosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Segestria Latreille, 1804** [also contains valid fossil species]
63. *Segestria exarata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
64. *Segestria sulcata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
65. *Segestria undulata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Siga Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
66. *Siga crinita* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- † **Spheconia Menge in C. L. Koch & Berendt, 1854 (*nomen nudum*)** **Palaeogene**
67. *Spheconia brevipes* Menge in C. L. Koch & Berendt, 1854* Pa Baltic amber
- † **Syphax C. L. Koch & Berendt, 1854** [also contains valid fossil species]
68. *Syphax hirtus* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Theridium Walckenaer, 1805** [now *Theridion* Walckenaer, 1805; which also contains valid fossil species]
69. *Theridium bifurcum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
70. *Theridium chorius* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
71. *Theridium clavigerum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
72. *Theridium crassipes* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
73. *Theridium setulosum* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- Thomisus Walckenaer, 1805** [also contains valid fossil species]
74. *Thomisus matutinus* Menge, 1856 Pa Baltic amber
- † **Thyelia C. L. Koch & Berendt, 1854** [also contains valid fossil species]
75. *Thyelia mengei* Giebel, 1856 Pa Baltic amber
76. *Thyelia pectinata* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
77. *Thyelia spinosa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
- † **Zilla C. L. Koch & Berendt, 1834** [also contains valid fossil species]
78. *Zilla cornumana* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber
79. *Zilla spinipalpa* Menge in C. L. Koch & Berendt, 1854 Pa Baltic amber

MISIDENTIFICATIONS

- Aranea Clerck, 1757** [now *Araneus* Clerck, 1757; which also contains valid fossil species]
1. *Aranea fusca pilosa* Bloch, 1776 [*nomen dubium*; non Araneae?] Qt Copal
- † **Araneaovoius Dunlop & Braddy, 2011 [ichnogenus]** **Palaeogene**
2. *Araneaovoius columbiae* (Scudder 1878)* [fossil egg sac] Pa Canada / USA
- † **Archaeometa Pocock, 1911** **?Devonian – Carb.**
3. ?*Archaeometa devonica* Størmer, 1976 [unidentifiable] D Alken an der Mosel
4. *Archaeometa nephilina* Pocock, 1911* [not identified] C Coseley
- † **Arachnometa Petrunkevitch, 1949** **Carboniferous**

5. *Arachnometa tuberculata* Petrunkevitch, 1949* [not identified] C Coseley
- † ***Eopholcus* Frič, 1904** **Carboniferous**
6. *Eopholcus pedatus* Frič, 1904* [not identified] C Nýřany
- † ***Oichnus* Bromley 1981 [ichnogenus]** ??
7. *Oichnus bavincourti* (Vaillant, 1909) [at one stage placed in *Cteniza*] Pa Northern France
- † ***Palaeocteniza* Hirst, 1923** **Devonian**
8. *Palaeocteniza crassipes* Hirst, 1923* [juvenile trigonotarbid?] D Rhyne chert
- † ***Pleurolycosa* Frič, 1904** **Carboniferous**
9. *Pleurolycosa prolifera* (Frič, 1901)* [unidentifiable] C Nýřany

43,678 Recent species according to Platnick (2013)

HAPTOPODA

1 currently valid species of fossil haptopodid

† HAPTOPODA Pocock, 1911	Carboniferous
† PLESIOSIRONIDAE Pocock, 1911	Carboniferous
† <i>Plesiosiro</i> Pocock, 1911	Carboniferous
1. <i>Plesiosiro madeleyi</i> Pocock, 1911	C Coseley

no Recent species

AMBLYPYGI

9 currently valid species of fossil whip spider

AMBLYPYGI Thorell, 1882	Carbon. – Recent
= PHRYNÉIDES Walckenaer, 1837	
= PHRYNICHIDA Petrunkevitch, 1945a	
PALAEOAMBLYPYGI Weygoldt, 1996 (suborder)	Carbon. – Recent
family uncertain	
† Sorellophrynus Harvey, 2002	Carboniferous
= † <i>Protophrynus</i> Petrunkevitch, 1913 (preoccupied)	
1. <i>Sorellophrynus carbonarius</i> (Petrunkevitch, 1913)*	C Mazon Creek
† Thelyphrynus Petrunkevitch, 1913	Carboniferous
2. <i>Thelyphrynus elongatus</i> Petrunkevitch, 1913	C Mazon Creek
PARACHARONTIDAE Weygoldt, 1996	Carbon. – Recent
† Graeophonus Scudder, 1890b	Carboniferous
3. <i>Graeophonus anglicus</i> Pocock, 1911	C Coseley
4. <i>Graeophonus carbonarius</i> (Scudder, 1876)*	C Cape Breton
5. <i>Graeophonus scudderi</i> Pocock, 1911	C Mazon Creek
EUAMBLYPYGI Weygoldt, 1996 (suborder)	Cretaceous – Recent
CHARINIDAE Quintero, 1986	Recent
no fossil record	
NEOAMBLYPYGI Weygoldt, 1996 (infraorder)	Cretaceous – Recent
CHARONTIDAE Simon, 1892a	Recent
no fossil record	
PHRYNOIDEA Blanchard, 1852	Cretaceous – Recent
PHRYNICHIDAE Simon, 1892a	Recent
no fossil record	
PHRYNIDAE Blanchard, 1852	Cretaceous – Recent
= † ELECTROPHRYNIDAE Petrunkevitch, 1971	
† Britopygus Dunlop & Martill, 2002	Cretaceous
6. <i>Britopygus weygoldti</i> Dunlop & Martill, 2002	K Crato Formation
† Electrophrynus Petrunkevitch, 1971	Neogene
7. <i>Electrophrynus mirus</i> Petrunkevitch, 1971	Ne Chiapas amber
Phrynus Lamarck, 1801	Neogene – Recent

8. *Phrynus mexicana* Poinar & Brown, 2004 Ne Chiapas amber
 9. *Phrynus resinae* (Schawaller, 1979*b*) Ne Dominican amber

NOMINA DUBIA

1. *Phrynus fossilis* Keferstein, 1834 Pa Aix-en-Provence
 i. = *Phrynus marioni* Gourret, 1887 Pa Aix-en-Provence

136 Recent species according to Harvey (2003)

UROPYGI

7 currently valid species of fossil whip scorpion

UROPYGI Thorell, 1882	Carbon. - Recent
= THELYPHONIDA Latreille, 1804b	
= UROTRICHA C. L. Koch, 1851	
= OXOPOEI Thorell, 1888	
= HOLOPELTIDIA Börner, 1902	
plesion genera	
† Geralinura Scudder, 1884	Carboniferous
1. <i>Geralinura britannica</i> Pocock, 1911	C Coseley
2. <i>Geralinura carbonaria</i> Scudder, 1884*	C Mazon Creek
i. = <i>Geralinura gigantea</i> Petrunkevitch, 1913	C Mazon Creek
ii. = <i>Geralinura similis</i> Petrunkevitch, 1913	C Mazon Creek
† Parageralinura Tetlie & Dunlop, 2008	Carboniferous
3. <i>Parageralinura naufraga</i> (Brauckmann & Koch, 1983)	C Hagen-Vorhalle
4. <i>Parageralinura neerlandicus</i> Laurentiaux-Viera & Laurentiaux, 1961.....	C Limburg
† Proschizomus Dunlop & Horrocks, 1996	Carboniferous
5. <i>Proschizomus petrunkevitchi</i> Dunlop & Horrocks, 1996	C Coseley
† Prothelyphonus Frič, 1904	Carboniferous
6. <i>Prothelyphonus bohemicus</i> (Kušta, 1884 <i>b</i>)	C Rakovník
i. = <i>Prothelyphonus cordai</i> Frič, 1904	C Rakovník
ii. = <i>Geralinura crassa</i> Kušta, 1888	C Rakovník
iii. = <i>Geralinura noctua</i> Kušta, 1888	C Rakovník
iv. = <i>Geralinura scudderi</i> Kušta, 1888	C Rakovník
THELYPHONIDAE Lucas 1835	Cretaceous – Recent
† Mesoproctus Dunlop, 1988	Cretaceous
7. <i>Mesoproctus rowlandi</i> Dunlop, 1998	K Crato Formation
<i>Mesoproctus</i> sp. in Dunlop & Martill (2002)	K Crato Formation

MISIDENTIFICATIONS

1. *Thelyphonus hadleyi* Pierce, 1945 [unidentifiable, ?algal]Ne California

SCHIZOMIDA

6 currently valid species of fossil schizomid from 6 published names

- the fossil family Calcitronidae cannot be meaningfully compared to the Recent families

SCHIZOMIDA Petrunkevitch, 1945b	Palaeogene – Recent
= TARTARIDES Thorell, 1888 (tribe)	
= COLOPYGA Cook, 1899 (order)	
= SCHIZOPELTIDA Börner, 1902 (tribe)	
† CALCITRONIDAE Petrunkevitch, 1945b	Palaeogene – Neogene
† <i>Calcitro</i> Petrunkevitch, 1945b	Palaeogene – Neogene
1. <i>Calcitro fisheri</i> Petrunkevitch, 1945b*	Ne Onyx Marble
2. <i>Calcitro oplonis</i> Lin in Lin et al., 1988	Pa Shandong, China
HUBBARDIIDAE Cook, 1899	Neogene – Recent
<i>Antillostenochrus</i> Armas and Teruel, 2002	Neogene – Recent
3. <i>Antillostenochrus pseudoannulatus</i> (Krüger & Dunlop, 2010)	Ne Dominican Amber
† <i>Calcoschizomus</i> Pierce, 1951	Neogene
4. <i>Calcoschizomus latisternum</i> Pierce, 1951	Ne Onyx Marble
† <i>Onychothelyphonus</i> Pierce, 1950	Neogene
5. <i>Onychothelyphonus bonneri</i> Pierce, 1950	Ne Onyx Marble
<i>Rowlandius</i> Reddell & Cockendolpher, 1995	Neogene – Recent
6. <i>Rowlandius velteni</i> (Krüger & Dunlop, 2010)	Ne Dominican Amber
PROTOSCHIZOMIDAE Rowland, 1975	Recent
no fossil record	

267 Recent species according to Harvey (pers. comm. 2009)

References

- Absolon, K. & Kratochvíl, J. 1932. Zur Kenntnis der höhlenbewohnenden Araneae der illyrischen Karstgebiete. *Mitteilungen über Höhlen- und Karstforschung*, 3: 73–81.
- Agassiz, L. 1844. *Monographie des poisons fossils du Vieux Gres Rouge ou Systeme Devonian*. Neufchatel, folio: 171 pp.
- Allen, J. G. & Feldman, R. M. 2005. *Panduralimulus babcocki* n. gen. and sp., a new Limulacean horseshoe crab from the Permian of Texas. *Journal of Paleontology*, 79: 594–600.
- Ambrose, T. & Romano, M. 1972. New Upper Carboniferous Chelicerata (Arthropoda) from Somerset, England. *Palaeontology* 15: 569–578.
- Ambrus, B. & Hably, L. 1979. *Eriophyes daphnogene* sp. n. a fossil gall from the Upper Oligocene of Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici*, 71: 55–56.
- Amerling, C. 1862. Naturökonomie der von ihm beobachteten Milben, insbesondere der Trombidieen. *Sitzungsberichte der Königlich Böhmisches Gesellschaft der Wissenschaften in Prague*, 2: 54–56.
- Ammon, L. von 1901. Ueber *Anthracomartus* aus dem Pfälzischen Carbon. *Geognostische Jahreshefte*, 13: 1–6.
- Anderson, L. I., Dunlop, J. A. & Trewin, N. H. 2000. A Middle Devonian chasmataspid arthropod from Achanarras Quarry, Caithness, Scotland. *Scottish Journal of Geology*, 36: 151–158.
- Andrée, K. 1913. Ueber *Anthracophrynus tuberculatus* nov. gen. nov. spec. aus dem productiven Karbon von Dudweiler im Saar-Revier, nebst einer Liste der bisher im Karbon Deutschland gefundenen Arachnoiden-Reste. *Jahres-Bericht und Mitteilungen der Oberrheinischen Geologischen Vereins*, 3: 89–93.
- Ansorge, J. 2003. Insects from the Lower Toarcian of Middle Europe and England. *Acta zoologica cracoviensia*, 46 (suppl.–Fossil Insects): 291–310.
- Aoki, J. 1965. Oribatiden (Acarina) Thailands. I. *Nature and Life in Southeast Asia*, 4: 129–193.
- Aoki, J. 1966a. A remarkable new oribatid mite from South Japan (Cryptostigmata: Tokunocepheidae, fam. nov.). *Acarologia*, 8: 358–364.
- Aoki, J. 1966b. Epizoic symbiosis: an oribatid mite, *Symbioribates papuensis*, representing a new family, from cryptogamic plants growing on backs of Papuan weevils (Acari: Cryptostigmata). *Pacific Insects*, 8: 281–289.
- Aoki, J. 1974. [On the fossil mites in Mizunami amber from Gifu Prefecture, Central Japan.] *Bulletin of the Mizunami Fossil Museum*, 1: 397–399 [in Japanese with English summary].
- Aoki, J. 1976. Oribatid mites from the IBP Study Area, Pasoh Forest Reserve, West Malaysia. *Nature and Life in Southeast Asia*, 7: 39–59.
- Aoki, J., Takaku, G. & Ito, F. 1994. Aribatidae, a new myrmecophilous oribatid mite family from Java. *International Journal of Acarology*, 20: 3–10.

- Arillo, A. & Subías, L.S. 2000. A new fossil oribatid mite, *Arachaeocheustes minguezae* n. gen. n. sp. from Spanish Lower Cretaceous amber. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 84: 231–236.
- Arillo, A. & Subías, L.S. 2002. Second fossil oribatid mite from the Spanish Lower Cretaceous amber. *Eupterotegaeus bitranslamellatus* n. sp. (Acariformes, Oribatida, Cepheidae). *Acarologia*, 42: 403–406.
- Arillo, A., Subías, L. S. & Shtanchaeva, U. 2008. A new fossil oribatid mite, *Ommatocephus nortoni* sp. nov. (Acariformes, Oribatida, Cepheidae), from a new outcrop of Lower Cretaceous Álava amber (northern Spain). *Systematic and Applied Acarology*, 13: 252–255.
- Arillo, A., Subías, L. S. & Shtanchaeva, U. 2009. A new fossil species of oribatid mite, *Ametroproctus valeriae* sp. nov. (Acariformes, Oribatida, Ametroproctidae), from the Lower Cretaceous amber of San Just, Teruel Province, Spain. *Cretaceous Research*, 30: 322–324.
- Armas L. F. de & Teruel, R. 2002. Un género nuevo de Hubbardiidae (Arachnida: Schizomida) de las Antillas Mayores. *Revista Ibérica de Aracnología*, 6: 45–52.
- Atyeo, W. T. & Baker, E. W. 1964. Tarsocheylidae, a new family of prostigmatic mites (Acarina). *Bulletin of the University of Nebraska State Museum*, 4: 243–256.
- Atyeo W. T. & Gaud, J. 1979. Ptyssalgidae, a new family of analgoid feather mites (Acarina, Acaridida). *Journal of Medical Entomology*, 16: 306–308.
- Atyeo, W. T. & Peterson, P. C. 1972. The feather mite family Alloptidae Gaud, new status, I. The subfamilies Trouessartiinae Gaud and Thysanocercinae, new subfamily (Analgoidea). *Zoologischer Anzeiger*, 188: 56–60.
- Atyeo W. T., Baker, E. W. & Delfinado M. D. 1974. *Gaudiella minuta*, a new genus and species of mite (Acarina: Acaridida) belonging to the new family Gaudiellidae. *Journal of the Washington Academy of Sciences*, 64: 295–298.
- Audouin, V. 1826. Explication sommaire des planches d'arachnides de l'Égypte et de la Syrie. *In Description de l'Égypte ou Recueil des Observations et des Recherches qui ont été Faites en Égypte Pendant l'Expédition de l'Armée Française, 1st edition, 1(4)*, 99–186. C. L. F. Panckoucke, Paris.
- Ausserer, A. 1867. Die Arachniden Tirols nach ihrer horizontalen und verticalen Verbreitung; 1. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 17: 137–170.
- Ausserer, A. 1875. Zweiter Beitrag zur Kenntniss der Arachniden-Familie der Territelariae Thorell (Mygalidae Autor). *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 25: 125–206.
- Ayyildiz, N. & Luxton, M. 1989. Epimerellidae (Acari, Oribatida), a new mite family. *Journal of Natural History*, 23: 1381–1386.
- Baily, W. H. 1863. Remarks on some Coal Measures Crustacea belonging to the genus *Belinurus*, König, with description of two new species from Queen's County, Ireland. *Annals and Magazine of Natural History*, 11: 107–114.

- Baily, W. H. 1869. On fossils obtained at Kiltorcan Quarry, Co. Kilkenny. *British Association Report*, pp. 73–75.
- Baker, E. W. 1949. Pomerantziidae, a new family of prostigmatic mites. *Journal of the Washington Academy of Science*, 39: 269–271.
- Baker, E. W. & Pritchard, A. E. 1953. The family categories of tetranychoid mites, with a review of the new families Linotetranae and Tuckerellidae. *Annals of the Entomological Society of America*, 46: 243–258.
- Baldwin W. & Sutcliffe, W. H. 1904. *Eoscorpius sparthensis* n. sp. from the Middle Coal Measures of Lancashire. *Quarterly Journal of the Geological Society of London*, 60: 395–398.
- Balogh, J. 1958. Oribatides nouvelles de l'Afrique tropicale. *Revue Zoologie Botanique Africaines*, 58: 1–34.
- Balogh, J. 1968. New oribatids (Acari) from New Guinea. *Acta Zoologica Academiae Scientiarum Hungaricae*, 14: 259–285.
- Balogh, J. 1970. New oribatids (Acari) from New Guinea. II. *Acta Zoologica Academiae Scientiarum Hungaricae*, 16: 291–344.
- Balogh, J. 1972. *The oribatid genera of the world*. Akadémiai Kiadó, Budapest, 188 pp.
- Balogh, J. 1983. A partial revision of the Oppiidae Grandjean, 1954 (Acari: Oribatei). *Acta Zoologica Academiae Scientiarum Hungaricae*, 29: 1–79.
- Balogh, J. & Balogh, P. 1984. A review of the Oribatuloidea Thor, 1929 (Acari: Oribatei). *Acta Zoologica Hungarica*, 30: 257–313.
- Balogh, J. & Balogh, P. 1992. *The oribatid mites genera of the world. I*. Hungarian National Museum Press, 263 pp.
- Balzan, L. 1888. *Chernetidae Nonnullae Sud-Americanae, III*. Asuncion.
- Balzan, L. 1892. Voyage de M. E. Simon au Venezuela (Décembre 1887 – Avril 1888). Arachnides. Chernetes (Pseudoscorpiones). *Annales de la Société Entomologique de France*, 60: 497–552.
- Bamber, R. 2007. A holistic re-interpretation of the phylogeny of the Pycnogonida Latreille, 1810 (Arthropoda). *Zootaxa*, 1668: 295–312.
- Banks, N. 1892a. A new genus of Phalangiidae. *Proceedings of the Entomological Society of Washington*, 2(2): 249–251.
- Banks, N. 1893. The Phalanginae of the United States. *The Canadian Entomologist*, 25: 205–211.
- Banks, N. 1895. Notes on the Pseudoscorpionida. *Journal of the New York Entomological Society*, 3: 1–13.
- Banks, N. 1896. New North American spiders and mites. *Transactions of the American Entomological Society*, 23: 57–77.
- Banks, N. 1905. Arachnids from the Cocos Island. *Proceedings of the Entomological Society of Washington*, 7: 20–23.
- Barbour, E. H. 1914. Carboniferous eurypterids of Nebraska. *American Journal of Science*, 4th Series, 38: 507–510.

- Beecher, C. E. 1902. Note on a new xiphosuran from the Upper Devonian of Pennsylvania. *American Geologist*, 29, 143–146.
- Beecher, C. E. 1904. Note on a new Permian xiphosuran from Kansas. *American Journal of Science*, 4th Series, 17: 23–24.
- Beier, M. 1932a. Pseudoscorpionidea I. Subord. Chthoniinea et Neobisiinea. *Tierreich*, 57: i–xx, 1–258.
- Beier, M. 1932b. Pseudoscorpionidea II. Subord. C. Cheliferina. *Tierreich*, 58: i–xxi, 1–294.
- Beier, M. 1937. Pseudoscorpione aus dem baltischen Bernstein. *Festschrift zum 60. Geburtstag von Professor Dr. Embrik Strand, Riga*, 2: 302–316.
- Beier, M. 1947a. Pseudoskorpione im Baltischen Bernstein und die Untersuchung von Bernstein-Einschlüssen. *Mikroskopie, Wien*, 1: 188–199.
- Beier, M. 1947b. Zur Kenntnis der Pseudoscorpionidenfauna des südlichen Afrika, insbesondere der südwest und südafrikanischen Trockengebiet. *Eos, Madrid*, 23: 285–339.
- Beier, M. 1955. Pseudoscorpione im baltischen Bernstein aus dem Geologischen Staatsinstitut in Hamburg. *Mitteilungen aus dem Mineralogisch-Geologischen Staatsinstitut in Hamburg*, 25: 48–54.
- Bell, W. A. 1922. A new genus of Characeae and new Merostomata from the Coal Measures of Nova Scotia. *Transactions of the Royal Society of Canada*, 4: 159–167.
- Bergström, J., Stürmer, W. & Winter, G. 1980. *Palaeoisopus*, *Palaeopantopus* and *Palaeothea*, pycnogonid arthropods from the Lower Devonian Hunsrück Slate, West Germany. *Paläontologische Zeitschrift*, 54: 7–54.
- Berland, L. 1913. Araignées. In *Mission du Service géographique de l'armée pour la mesure d'un arc du méridien équatorial en Amérique du Sud (1899-1906)*. Paris, 10: 78-119.
- Berland, L. 1939. Description de quelques Araignées fossils. *Revue Française d'Entomologie*, 6: 1–9.
- Berlese, A. 1885. Acarorum Systematis. *Bullettino della Società Entomologica Italiana*, 17: 121–135.
- Berlese, A. 1888. Acari Austro-Americani quos collegit Aloysius Balzan. Manipulus primus. Species novas circiter quinquaginta complectens. *Bollettino della Società Entomologica Italiana*, 20: 171–222.
- Berlese, A. 1896. Acari, Myriapoda et Scorpiones hucusque in Italia reperta. *Acari, Myriapoda et Scorpiones in Italia reperta*, Fasc. 79, 15 pp., 6 pls.
- Berlese, A. 1899. Gli acari agrarii. Puntat II. *Rivista di Patologia Vegetale, Padova*, 7: 312–344.
- Berlese, A. 1910. Lista di nuove specie e nuovi generi di Acari. *Redia*, 6: 242–271.
- Berlese, A. 1914. Acari nuovi. *Redia*, 10: 1–150.
- Berlese, A. 1923. Centuria sesta di Acari nuovi. *Redia*, 15: 237–262.
- Bernini, F. 1975. Notulae Oribatologicae XII. Una nuova specie di *Carabodes* affine a *C. minusculus* Berlese 1923 (Acarida, Oribatei). *Redia* 56: 455–471.
- Bertkau, P. 1872. Über die Respirationsorgane der Araneen. *Archiv für Naturgeschichte*, 38: 208–233.

- Bertkau, P. 1878a. Versuch einer natürlichen Anordnung der Spinnen, nebst Bemerkungen zu einzelnen Gattungen. *Archiv für Naturgeschichte*, 44: 351–410.
- Bertkau, P. 1878b. Einige Spinnen und ein Myriapode aus der Braunkohle von Rott. *Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westfalens, Bonn*, 35: 346–360.
- Bertkau, P. 1882. Ueber das Cribellum und Calamistrum. Ein Beitrag zur Histologie, Biologie und Systematik der Spinnen. *Archiv für Naturgeschichte*, 48: 316–362.
- Beyschlag, F. & Fritsch, K. von 1899. Das jüngere Steinkohlengebirge und das Rothliegende in der Provinz Sachsen und den angrenzenden Gebieten. *Abhandlungen der Königlich Preussischen geologischen Landesanstalt*, 10: 1–263.
- Blackwall, J. 1833. Characters of some undescribed genera and species of Araneidae. *London philosophical Magazine and Journal of Science*, 3: 104–112, 187–197, 344–352, 436–443.
- Blackwall, J. 1834a. Characters of some undescribed species of Araneidae. *London philosophical Magazine and Journal of Science*, 5: 50–53.
- Blackwall, J. 1834b. *Researches in Zoology*. London, pp. 229–433.
- Blackwall, J. 1841. The difference in the number of eyes with which spiders are provided proposed as the basis of their distribution into tribes; with descriptions of newly discovered species and the characters of a new family and three new genera of spiders. *Transactions of the Linnean Society, London*, 18: 601–670.
- Blackwall, J. 1853. Descriptions of some newly discovered species of Araneida. *Annals and Magazine of Natural History, series 2*, 11: 14–25.
- Blackwall, J. 1859. Descriptions of newly discovered spiders captured by James Yate Johnson Esq., in the island of Maderia. *Annals and Magazine of Natural History, series 3*, 4: 255–267.
- Blackwall, J. 1862. Descriptions of newly-discovered spiders from the island of Madeira. *Annals and Magazine of Natural History, series 3*, 9: 370–382.
- Blackwall, J. 1864. *A History of the Spiders of Great Britain and Ireland. Part II*. The Ray Society, London, 1864 pp. 175–384.
- Blackwall, J. 1870. Notes on a collection of spiders made in Sicily in the spring of 1868, by E. Perceval Wright, M.D., with a list of the species, and descriptions of some new species and of a new genus. *Annals and Magazine of Natural History, series 4*, 5: 392–405.
- Blanchard, E. 1852. Arachnides. In *L'organisation du règne animal, 2nd Edition, vol. 2*. E. Blanchard, Paris.
- Błaszak, J., Cokendolpher, J. C. & Polyak, V. J. 1995. *Paleozerccon cavernicolous*, a new genus and new species of fossil mite from a cave in the southwestern U.S.A. (Acari, Gamasida: Zerconidae). *International Journal of Acarology*, 21: 253–259.
- Bleicher, M. 1897. Sur la découverte d'une nouvelle espèce de limule dans les marnes irisées de Lorraine. *Bulletin de la Societe des Sciences de Nancy*, (2)14: 116-126.

- Bloch, M. [E.] 1776. Naturgeschichte des Kopals. *Beschäftigungen der Berlinischen Gesellschaft Naturforschender Freunde*, 2: 91–196.
- Bode, A. 1951. Ein Liassischer Skorpionide. *Palaeontologische Zeitschrift*, 24: 58–65.
- Bolland, H. R. & Magowski, W. Ł. 1990. *Neophyllobius succineus* n. sp. from Baltic amber (Acari: Raphignathoidea: Camerobiidae). *Entomologische Berichten*, 50: 17–21.
- Bosselaers, J. 2004. A new *Garacops* species from Madagascar copal (Araneae: Selenopidae). *Zootaxa*, 445: 1–7.
- Bosselaers, J., Dierick, M., Cnudde, V., Masschaele, B., Van Hoorebeke, L. & Jacobs, P. 2010. High resolution X-ray computed tomography of an extant new *Donuea* (Araneae: Liocranidae) species in Madagascan copal. *Zootaxa*, 2427: 25–35.
- Bottali, P. 1975. Note su due rari esemplari di Araneidi (Aracnidi) rinvenuti nei depositi diatomitici (facies lacustre) di Riano Flaminio (Roma). *Fragmenta entomologica*, 11: 169–174.
- Braddy, S. J., Aldridge, R. J. & Theron, J. N. 1995. A new eurypterid from the Late Ordovician Table Mountain Group, South Africa. *Palaeontology*, 38: 563–581.
- Braddy, S. J., Selden, P. A. & Doan Nhat T. 2002. A new carcosomatid eurypterid from the Upper Silurian of Northern Vietnam. *Palaeontology*, 45: 897–915.
- Brauckmann, C. 1982. Der Schwertschwanz *Euproops* (Xiphosuraidea, Limulina, Euproopacea) aus dem Ober-Karbon des Piesbergs bei Osnabrück. *Osnabücker naturwissenschaftliche Mitteilungen*, 9: 17–26.
- Brauckmann, C. 1984. Eine neue Arachniden-Art aus dem Westfalium des Saargebietes (West-Deutschland). *Dortmunder Beiträge zur Landeskunde, naturwissenschaftliche Mitteilungen*, 18: 95–103.
- Brauckmann, C. 1987. Neue Arachniden (Ricinuleida, Trigonotarbida) aus dem Namurium B von Hagen-Vorhalle (Ober-Karbon; West-Deutschland). *Dortmunder Beiträge der Landeskunde, naturwissenschaftliche Mitteilungen*, 21: 97–109.
- Brauckmann, C. & Koch, L. 1983. *Prothelyphonus naufragus* n. sp., ein neuer Geisselskorpion [Arachnida: Thelyphonida: Thelyphonidae] aus dem Namurium unteres Oberkarbon) von West-Deutschland. *Entomologica Germanica*, 9: 63–74.
- Brauckmann, C., Koch, L. & Kemper, M. 1985. Spinnentiere (Arachnida) und Insekten aus den Vorhalle-Schichten (Namurian B; Ober-Karbon) von Hagen-Vorhalle (West-Deutschland). *Geologie und Paläontologie in Westfalen*, 3: 1–131.
- Brauer, F., Redtenbacher, J. & Ganglbauer, L. 1889. Fossile Insekten aus der Juraformation Ost-Siberiens. *Mémoires de l'Académie Impériale des Sciences de St.-Petersbourg, VII serie*, 36(15): 1–22.
- Braun, C. F. W. 1860. Die Thiere in den Pflanzenschifern der Gegend von Bayreuth. Programm zum Jahresbericht der Königl. Kreis-Landwirtschafts- und Gewerbschule zu Bayreuth für das Schuljahr 1859/60. *Jahresbericht von der Königl. Kreis-Landwirtschafts- und Gewerbschule zu Bayreuth für das Schuljahr 1859/60*: 11 pp.

- Brescovit, A. D. 1997. Revisão de Anyphaeninae Bertkau a nível de gêneros na região neotropical (Araneae, Anyphaenidae). *Revista Brasileira de Zoologia*, 13: 1–187.
- Briggs, D. E. G., Siveter, D. J., Siveter, D. J., Sutton, M. D., Garwood, R. J. & Legg, D. 2012. Silurian horseshoe crab illuminates the evolution of arthropod limbs. *Proceedings of the National Academy of Sciences of the United States of America*, 109: 15702–15703.
- Briggs, T. A. 1971. Relict harvestmen from the Pacific northwest (Opiliones). *Pan-Pacific Entomologist*, 74: 165–178.
- Bristowe, W. S. 1938. The classification of spiders. *Proceedings of the Zoological Society of London*, 108: 285–322.
- Bristowe, W. S. 1939. *The comity of spiders. Volume 1*. London, 228 pp.
- Brongniart, C. 1877. Note sur une Aranéide fossile des terrains tertiaires. *Annales de la Société Entomologique de France*, (5) 7: 221–224.
- Bruce, W. A. & Johnston, D. E. 1976. *Gaudoglyphus* n. gen., based on *Analges minor* Nörner (Acari: Gaudoglyphidae n. fam.). *International Journal of Acarology*, 2: 29–33.
- Broili, F. 1928. Crustaceenfunde aus dem rheinischen Unterdevon. I. Über Extremitätenreste. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung*, 1928: 197–201.
- Broili, F. 1930. Über ein neues Exemplar von *Palaeopantopus*. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung*, 1930: 209–214.
- Bromley, R.G. 1981. Concepts in ichnotaxonomy illustrated by small round holes in shells. *Acta Geologica Hispànica*, 16: 55–64.
- Bronn, H. G. 1856. *Lethaea Geognostica oder Abbildung und Beschreibung für die Gebirgs-Formationen bezeichnendsten Versteinerungen. Dritter Band*. Schweizerbart'sche Verlagshandlung und Druckerei 1853–1856, pp. 622–639.
- Buckland, W. 1837. *The Bridgewater treatises on the power, wisdom and goodness of God as manifested in the creation. Treatise IV. Geology and mineralogy with reference to natural theology. 2nd Edition*. William Pickering, London.
- Bulanova-Zachvatkina, E. M. 1974. [New genera of oribatid mites from the Upper Cretaceous of Tajmyr.] *Paleontological Journal*, 1974: 141–144. [In Russian]
- Burmeister, H. 1843. *Die Organisation der Trilobiten, aus ihren lebenden Verwandten entwickelt; nebst systematischen Uebersicht aller zeither beschriebenen Arten*. G. Reimer, Berlin, 148 pp.
- Cambridge, F. O. P.- 1893. Handbook to the study of British spiders (Drassidae and Agalenidae). *British Nature Supplement*, 3: 117–170.
- Cambridge, F. O. P.- 1899. Arachnida. Araneida. Vol. 2. *Biologia Centrali-Americana*: pp. 41–88.

- Cambridge, O. P.- 1870. Descriptions and sketches of two new species of Araneida, with characters of a new genus. *Journal of the Linnean Society of London*, 10: 398–405.
- Cambridge, O. P.- 1871. Arachnida (1870). *The Zoological Record*, 7: 207–224.
- Cambridge, O. P.- 1873. On some new genera and species of Araneida. *Proceedings of the Zoological Society of London*, 1873: 112–129.
- Cambridge, O. P.- 1874. On some new genera and species of Araneida. *Annals and Magazine of Natural History*, series 4, 14: 169–183.
- Cambridge, O. P.- 1876. On a new order and some new genera of Arachnida from Kerguelen's Land. *Proceedings of the Zoological Society of London*, 1876: 258–265.
- Cambridge, O. P.- 1877. On some new species of Araneida, with characters of two new genera and remarks on the families Podophthalmides and Dinopides. *Proceedings of the Zoological Society of London*, 1877: 557–578.
- Cambridge, O. P.- 1879a. On some new and rare spiders from New Zealand, with characters of four new genera. *Proceedings of the Zoological Society of London*, 1879: 681–703.
- Cambridge, O. P.- 1879b. On some new and rare British spiders, with characters of a new genus. *Annals and Magazine of Natural History*, 4: 190–215.
- Cambridge, O. P.- 1881. On some new genera and species of Araneidea. *Proceedings of the Zoological Society of London*, 1881: 765–775.
- Cambridge, O. P.- 1882a. On new genera and species of Araneidea. *Proceedings of the Zoological Society of London*, 1882: 423–442.
- Cambridge, O. P.- 1882b. Arachnida (1881). *The Zoological Record*, 18: 1–32.
- Cambridge, O. P.- 1894. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americana*: pp. 121–144.
- Cambridge, O. P.- 1895. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americana*: pp. 145–160.
- Cambridge, O. P.- 1898. Arachnida. Araneida. Vol. 1. *Biologia Centrali-Americana*: pp. 233–288.
- Cambridge, O. P.- 1902. On new and rare British Arachnida. *Proceedings of the Dorset Natural History and Antiquarian Field Club*, 23: 16–40.
- Camin, J. H. & Gorirossi, F. E. 1955. A revision of the suborder Mesostigmata (Acarina), based on new interpretations of comparative morphological data. *Chicago Academy of Sciences Special Publication*, 11: 1–70.
- Camin J. H., Moss W. W. & Oliver J. H. 1967. Cloacaridae, a new family of cheyletoid mites from the cloaca of aquatic turtles. *Journal of Medical Entomology*, 4: 261–272.
- Campos, D. R. B. 1986. Primeiro registro fóssil de Scorpionoidea na Chapada do Araripe (Cretáceo Inferior), Brasil. *Anais do Academia Brasileira das Ciências*, 58: 135–137.
- Canestrini, G. & Fanzago, F. 1877. Intorno agli Acari italiani. - Atti del R. Istituto Veneto Scienze, Lettere ed Arti, Ser. 5 4: 69–208.

- Canestrini, G. & Pavesi, P. 1870. Catalogo sistematico degli Araneida italiani. *Archivio per la zoologia, l'anatomia e la fisiologia*, (2)2: 1–44.
- Caporiacco, L. di 1949. Aracnidi della colonia de Kenya raccolti da Toschi e Meneghetti negli anni 1944–1946. *Commentationes Pontificiae Academiae Scientiarum*, 13: 309–492.
- Carvalho, M. P. G. de & Lourenço, W. R. 2001. A new family of fossil scorpions from the Early Cretaceous of Brazil. *Comptes Rendus de l'Académie de Sciences de Paris, Earth and Planetary Sciences*, 332: 711–716.
- Caster, K. E. & Brooks, H. K. 1956. New fossils from the Canadian–Chazan (Ordovician) hiatus in Tennessee. *Bulletin of American Palaeontology*, 36: 157–199.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1953. *Melbournopterus*, a new Silurian eurypterid from Australia. *Journal of Paleontology*, 27: 153–156.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1955. *Marsupipterus*, an unusual eurypterid from the Downtonian of England. *Journal of Paleontology*, 29: 1040–1041.
- Caster, K. E. & Kjellesvig-Waering, E. N. 1956. Some notes on the genus *Dolichopterus* Hall. *Journal of Paleontology*, 30: 19–28.
- Caster K. E. & Kjellesvig-Waering, E. N. 1964. Upper Ordovician eurypterids of Ohio. *Palaeontographica Americana*, 4 (32): 297–358.
- Chamberlin, J. C. 1923a. The genus *Pseudogarypus* Ellingsen (Pseudoscorpionida – Feallidae). *Entomological News*, 34: 146–149, 161–166.
- Chamberlin, J. C. 1923b. New and little known pseudoscorpions, principally from the islands and the adjacent shores of the Gulf of California. *Proceedings of the California Academy of Science*, (4)12: 353–387.
- Chamberlin, J. C. 1929. A synoptic classification of the false scorpions or chela-spinners, with a report on a cosmopolitan collection of the same. Part I. The Heterosphyronida (Chthoniidae) (Arachnida-Chelonethida). *Annals and Magazine of Natural History, series 10*, 4: 50–80.
- Chamberlin, J. C. 1930. A synoptic classification of the false scorpions or chela-spinners, with a report on a cosmopolitan collection of the same. Part II. The Diplosphyronida (Arachnida-Chelonethida). *Annals and Magazine of Natural History, series 10*, 5: 1–48, 585–620.
- Chamberlin, J. C. 1931a. The arachnid order Chelonethida. *Stanford University Publications, Biological Sciences*, 7: 1–284.
- Chamberlin, J. C. 1931b. A synoptic revision of the generic classification of the chelonethid family Cheliferidae Simon (Arachnida). *Canadian Entomologist*, 64: 289–294.
- Chamberlin, J. C. 1947. The Vachoniidae – a new family of false scorpions represented by two new species from caves in Yucatan (Arachnida, Chelonethida, Neobisioidea). *Bulletin of the University of Utah, Biological Series*, 10(4): 1–15.

- Chamberlin, R. V. 1917. New spiders of the family Aviculariidae. *Bulletin of the Museum of Comparative Zoology*, 61: 25–75.
- Chamberlin, R. V. 1922. Two new American arachnids of the order Pedipalpida. *Proceedings of the Biological Society of Washington*, 235: 11–12.
- Chamberlin, R. V. & Ivie, W. 1943. New genera and species of North American linyphiid spiders. *Bulletin of the University of Utah*, 33(10): 1–39.
- Chamberlin, R. V. & Mulaik, S. 1942. On a new family in the Notostigmata. *Proceedings of the Biological Society of Washington*, 55: 125–132.
- Chang A.-c. 1957. On the discovery of the Wenlockian *Eurypterus*-fauna from south China. *Acta Palaeontologica Sinica*, 5: 446–450.
- Chang J.-p. 2004. Some new species of spider and Sacculinidae fossils in Jehol biota. *Global Geology*, 23: 313–320.
- Chapman, F. 1932. Two new Australian fossil king-crabs. *Proceedings of the Royal Society of Victoria, New Series*, 44: 100–102.
- Charbonnier, S., Vannier, J. & Riou, B. 2007. New sea spiders from the Jurassic La Voulte-sur-Rhône Lagerstätte. *Proceedings of the Royal Society B*, 274: 2555–2561.
- Cheng X.-d., Meng Q.-j., Wang X.-r. & Gao C.-l. 2008. [New discovery of Nephilidae in Jehol biota (Araneae, Nephilidae).] *Acta zootaxonomica Sinica*, 33: 330–334. [in Chinese with English summary]
- Cheng X.-d., Shen C.-z. & Gao C.-l. 2009. [A new fossil spider of the Philodromidae from the Yixian Formation of western Liaoning Province, China (Arachnida, Araneae).] *Acta Arachnologica Sinica*, 18: 23–27. [in Chinese with English summary]
- Chernyshev, B. I. 1928. Nouvelles donnees sur les Xiphosura du basin Donetz. *Bulletin du Comité Géologique*, 47: 519–531.
- Chernyshev, B. I. 1933. [Arthropoda from the Urals and other regions of the USSR.] *Materials of the Central Scientific and Prospecting Institute Paleontology and Stratigraphy, Magazine*, 1: 15–25. [in Russian with English summary]
- Chernyshev, B. I. 1948. New representative of Merostomata from the Lower Carboniferous. *State of Kiev, Geological Collections*, 2: 119–130.
- Chlupáč, I. 1994. Pterygotid eurypterids (Arthropoda, Chelicerata) in the Silurian and Devonian of Bohemia. *Journal of the Czech Geological Society*, 39: 147–162.
- Chlupáč, I. 1995. Lower Cambrian arthropods from the Paseky Shale (Barrandian area, Czech Republic). *Journal of the Czech Geological Society*, 40: 9–36.
- Chlupáč, I. & Havlíček, V. 1965. *Kodymirus* n. g., a new aglaspid merostome of the Cambrian of Bohemia. *Sborník Geologických Věd. Paleontologie*, 6: 7–20.

- Ciurca Jr., S. J. & Tetlie, O. E. 2007. Pterygotids (Chelicerata; Eurypterida) from the Silurian Vernon Formation of New York. *Journal of Paleontology*, 81: 725–736.
- Clarke, J. M. 1902. Notes on Paleozoic crustaceans. *New York State Museum Report*, 54: 83–110.
- Clarke, J. M. 1907. The *Eurypterus* shales of the Shawangunk Mountains in Eastern New York. *New York State Museum Bulletin* 107: p. 295.
- Clarke, J. N. & Ruedemann, R. 1912. The Eurypterida of New York. – *New York State Museum, Memoir*, 14, 1–439.
- Clarke, J. M. 1919. *Bunaia woodwardi*, a new merostome from the Silurian waterlimes of New York. *Geological Magazine, Decade* 6, 6: 531–532.
- Claypole, E. W. 1890a. Palaeontological notes from Indianapolis (A. A. A. S.) *Pterichthys* – *Castoroides* – *Eurysoma* g. n. *American Geologist*, 6: 255–260.
- Claypole, E. W. 1890b. *Carcinosoma newlini*. *American Geologist*, 6: 400.
- Clerck, C. 1757. *Araneae suecici, descriptionibus et figuris oeneis illustrati, ad genera subalterna redacti speciebus ultra LX determinati. Svenska Spindlar, uti sina hufvud-slagter indelte samt...* - Stockholm, 154 pp.
- Cockerell, T. D. A. 1905. Two Carboniferous genera of xiphosurans. *American Geologist*, 36: 330.
- Cockerell, T. D. A. 1907. Some fossil arthropods from Florissant, Colorado. *Bulletin of the American Museum of Natural History*, 23: 605–616.
- Cockerell, T. D. A. 1916. The uropods of *Acanthotelson stimpsoni*. *Journal of the Washington Academy of Science*, 6: 234–236.
- Cockerell, T. D. A. 1917a. Arthropods in Burmese amber. *American Journal of Science, series 4*, 44: 360–368.
- Cockerell, T. D. A. 1917b. Arthropods in Burmese amber. *Psyche*, 24: 40–45.
- Cockerell, T. D. A. 1920. Fossil arthropods in the British Museum. I. *Annals and Magazine of Natural History, series 9*, 5: 273–279.
- Cockerell, T. D. A. 1925. Fossil insects in the United States National Museum. *Proceedings of the U. S. National Museum*, 64: 1–15.
- Coineau, Y. 1974. Un type nouveau d'Acariens Prostigmates libres: les Saxidromoidea, nouvelle super-famille. *Comptes rendus de l'Académie des Sciences, Paris série D*, 278: 1059–1062.
- Coineau, Y. & Magowski, W. Ł. 1994. Caeculidae in amber. *Acarologia*, 35: 243–246.
- Coineau, Y. & Poinar Jr., G. O. 2001. Un Caeculidae de l'ambre de la République Dominicaine. *Acarologia*, 41: 141–144.
- Coineau, Y & Theron, P. 1983. Les Micropsammidae, n. fam. d'Acariens Endeostigmata des sables fin. *Acarologia*, 24: 275–280.
- Cokendolpher, J. C. 1987. A new species of fossil *Pellobunus* from Dominican Republic amber (Arachnida: Opiliones: Phalangodidae). *Caribbean Journal of Science*, 22: 205–211.

- Cokendolpher, J. C. & Poinar Jr., G. O. 1992. Tertiary harvestmen from Dominican Republic amber (Arachnida: Opiliones: Phalangodidae). *Bulletin of the British arachnological Society*, 9: 53–56.
- Cokendolpher, J. C. & Poinar Jr., G. O. 1998. A new fossil harvestman from Dominican Republic amber (Opiliones, Samoidae, *Hummelinckiolus*). *Journal of Arachnology*, 26: 9–13.
- Comstock, J. H. 1940. *The spider book, revised and edited by Willis J. Gertsch*. Ithaca, New York, 729 pp.
- Condé, B. 1996. Les Palpigrales, 1885–1995: acquisitions et lacunes. *Revue suisse de Zoologie*, hors série 1: 87–106.
- Cook, D.R. 1967. Water mites from India. *Memoirs of the American Entomological Institute*, 9: 1–411.
- Cooke, J. A. L. 1965. Spider genus *Dysdera* (Araneae, Dysderidae). *Nature*, 205: 1027–1028.
- Conda, A. J. C. 1835. Ueber den in der Steinkohlenformation bei Cholme gefundenen fossilen Scorpion. *Verhandlungen der Gesellschaft des vaterländischen Museums in Böhmen, Prag*: 36.
- Conda, A. J. C. 1839. Ueber eine fossile Gattung der Afterscorpione. *Verhandlungen der Gesellschaft des vaterländischen Museums in Böhmen, Prag*: 14–18.
- Corronca, J. A. 2003. New genus and species of Selenopidae (Arachnida, Araneae) from Madagascar and neighbouring islands. *African Zoology*, 38: 387–392.
- Crônier, C. & Courville, P. 2005. New xiphosuran merostomata from the Upper Carboniferous of the Graissessac Basin (Massif Central, France). *Comptes Rendus Palevol*, 4: 123–133.
- Crosby, C. R. & Bishop, S. C. 1925. A new genus and two new species of spiders collected by *Bufo quercicus* (Holbrook). *Florida Entomologist* 9: 33–36.
- Cross, E. A. 1965. The generic relationships of the family Pyemotidae (Acarina: Trombidiformes). *Kansas University Science Bulletin*, 45: 29–275.
- Cunliffe, F. 1958. *Pyroglyphus morlani*, a new genus and species of mite forming a new family, Pyroglyphidae, in the Acaridae. *Proceedings of the Entomological Society of Washington*, 60: 85–86.
- Currie, L. D. 1927. On *Cyamocephalus*, a new synziphosuran from the Upper Silurian of Lesmahagow, Lanarkshire. *Geological Magazine*, 64: 153–157.
- Cutler, B. 1970. A fossil crab spider from West-ventral Wyoming (Araneae: Thomisidae). *Entomological News*, 81: 38–40.
- Daber, R. 1990. Arachnidenrest aus dem Westfal D von Zwickau-Oelsnitz. *Zeitschrift für geologische Wissenschaft, Berlin*, 18: 679–682.
- Dabert, J. 1994. Kiwilichidae fam. nov. eine neue Federfamilie (Astigmata, Pterolichoidea). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 11: 101–110.
- Daday, E. 1888. A Magyar nemzeti Muzeum àlskorpionak áttekintése. *Természetrázi Füzetek*, 11: 111–136, 165–192.

- Dahl, F. 1908. Die Lycosiden oder Wolfsspinnen Deutschlands und ihre Stellung im Haushalt der Natur. Nach statistischen Untersuchungen dargestellt. *Nova Acta Academiae Caesareae Leopoldino-Carolinae*, 88: 175–678.
- Dahl, F. 1912. Arachnoidea. In Korschelt, E. et al. (eds). *Handwörterbuch der Naturwissenschaften*, 1: 485–514.
- Dahl, F. 1913. *Vergleichende Physiologie und Morphologie der Spinnentiere unter besonderer Berücksichtigung der Lebensweise. 1. Die Beziehungen des Körperbaues und der Farben zur Umgebung*. Jena, 1913: 113 pp.
- Dalla Vecchia, F. M. & Selden, P. A. 2013. A Triassic spider from Italy. *Acta Palaeontologica Polonica*, 58: 325–330.
- Dalman, J. W. 1826. Om Insekter inneslutne I Copal, jemte beskrifning på några deribland förekommande nya släkten och arter. *Kungliga Svenska Vetenskapsakademiens Handlingar*, 46: 375–410.
- Dalmas, R. de 1916. Révision du genre *Orchestina* E.S., suivie de la description de nouvelles espèces du genre *Oonops* et d'une étude sur les Dictynidae du genre *Scotolathys*. *Annales de la Société Entomologique de France*, 85: 203–258.
- Dalmas, R. de 1917. Araignées de Nouvelle Zélande. *Annales de la Société Entomologique de France*, 86: 317–430.
- Dana, J. D. 1853. Crustacea, pt. II, Arachnopaoda or Pycnogonida. In United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842. Under the command of Chales Wilkes, U.S.N.C. Sherman, Philadelphia, 1382–1391.
- Davies, V. T. 1978. A new family of spiders (Araneae: Teemanaaridae). *Symposium of the Zoological Society of London*, 42: 293–302.
- Davies, V. T. 1980. *Malkara loricata*, a new spider (Araneidae: Malkarinae) from Australia. *Verhandlungen des 8. Internationalen Arachnologen-Kongresses. Wien, 1980*: 377–382.
- Deeleman-Reinhold, C. L. 1995. The Ochyroceratidae of the Indo-Pacific region (Araneae). *Raffles Bulletin of Zoology Supplement*, 2: 1–103.
- Delle, N. 1937. Zemgales lidzenuma, Augszemes un Lietuvas devona nogulumi. *Acta Universitatis Latviensis, Matēmatikas un Dabas Zinātņu Fakultātes Serija* 2(5): 105–384.
- De Geer, C. 1778. *Mémoires pour Servir à l'Histoire des Insectes, vol. 7*. Stockholm.
- De Kay, J. E. 1825. Observations on a fossil crustaceous animal of the order Branchiopoda. *Annals of the New York Lyceum of Natural History*, 1: 375–377.
- Delfinado, M. D. & Baker, E. W. 1974. Varroidae, a new family of mites on honeybees (Mesostigmata: Acarina). *Journal of the Washington Academy of Science*, 64: 4–10.
- De Lima, W. 1890. Note sur un nouvel *Eurypterus* du Rothliegendes de Bussaco. *Comunicações da Comissão dos Trabalhos Geològicos da Portugal*, 2: 153–157.

- Desmarest, A.-G. 1822. Les crustacés proprement dits. 66–154. In *Histoire naturelle des crustacés fossiles, sous les rapports zoologiques et géologiques*. F.-G. Levrault, Paris, Strasbourg, xx pp.
- Diener, C. 1924. Eurypterida. In Diener, C. (ed.). *Fossilium Catalogus I : Animalia*. W. Junk, Berlin, pp. 1–26.
- Dix, E. & Pringle, J. 1929. On the fossil Xiphosura from the South Wales Coalfield with a note on the myriapod *Euphoberia*. *Summary of Progress, Geological Survey of Great Britain*, 1928: 90–113.
- Dix, E. & Pringle, J. 1930. Some Coal Measures arthropods from the South Wales Coalfield. *Annals and Magazine of Natural History*, 6: 136–144.
- Dohrn, A. 1881. Die Pantopoden des Golfes von Neapel und der angrenzenden Meeresabschnitte. *Monographie der Fauna und Flora des Golfes von Neapel*, 3: 1–252.
- Doleschall, L. 1852. Systematisches Verzeichnis der im Kaiserthum Österreich vorkommenden Spinnen. *Sitzungsberichte der Mathematisch-Naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaft*, 9: 622–651.
- Donnadieu, A. L. 1875. *Recherches pour servir à l'histoire des Tetranyques*. – These. Faculte des Sciences de Lyon: 134 pp. [Thesis also published in a regular journal in 1876.]
- Dresco, E. 1970. Recherches sur la variabilité et la phylogénie chez les Opiliones du genre *Ischyropsalis* C. L. Koch (Fam. Ischyropsalidae), avec la creation de la famille nouvelle des Sabaconidae. *Bulletin du Muséum National d'Histoire Naturelle*, 2^e Serie, 41: 1200–1213.
- Dubey, D. P. 1985. A preliminary note on the eurypterid and trilobitid remains from the Upper Vidhyan rocks around Rewa, Madhya Pradesh. *Current Trends in Geology (IV Indian Geological Congress)*, 6: 63–78.
- Dubinin, V. B. 1953. Feather mites (Analgesoidea). II Families Epidermoptidae and Freyanidae. *Fauna SSSR. Paukoobrazyne* 6 (6): 3–411. [In Russian].
- Dubinin, V. B. 1957. On the orientation of the cephalic end of the Devonian pycnogonids of the genus *Palaeoisopus* and their systematic position in the Arthropoda. *Doklady Akademii Nauk SSSR*, 117: 881–884. [In Russian].
- Dufour, L. 1820. Description de cinq Arachnides nouvelles. *Annales générales des sciences physiques*, 5: 198–209.
- Dugès, A. 1834. Recherches sur l'ordre des Acariens et la famille des Trombidés en particulier. *Annales des Sciences Naturelles, Zoologie, série 2*, 1: 5–46.
- Dujardin, F. 1851. Sur des acariens a quatre pieds, parasites des vegeteux et qui doivent former un genre particulier (*Phytoptus*). In *Observations Zoologiques. Annales des Sciences Naturelles, série 3*, 15: 158–175.
- Dunbar, C. O. 1923. Kansas Permian insects, Part 2. *Paleolimulus*, a new genus of Paleozoic Xiphosura, with notes on other genera. *American Journal of Science*, 5th series, 5: 443–454.
- Dunbar, C. O. 1924. Kansas Permian insects. Part 1. The geologic occurrence and the environment of the insects. *American Journal of Science*, 5th series, 7: 171–209.

- Dunlop, J. A. 1995. Redescription of the Pennsylvanian trigonotarbid arachnid *Lissomartus* Petrunkevitch 1949 from Mazon Creek, Illinois. *Journal of Arachnology*, 23: 118–124.
- Dunlop, J. A. 1996. A trigonotarbid arachnid from the Upper Silurian of Shropshire. *Palaeontology*, 39: 605–614.
- Dunlop, J. A. 1998. A fossil whipscorpion from the Lower Cretaceous of Brazil. *Journal of Arachnology*, 26: 291–295.
- Dunlop, J. A. 1999. A replacement name for the trigonotarbid arachnid *Eotarbus* Dunlop. *Palaeontology*, 42: 191.
- Dunlop, J. A. 2002. Arthropods from the Lower Devonian Severnya Zemlya Formation of October Revolution Island, Russia. *Geodiversitas*, 24: 349–379.
- Dunlop, J. A. 2004. A spiny harvestman (Arachnida: Opiliones) from the Upper Carboniferous of Missouri, USA. In Logunov, D. V. & Penney, D (eds). Proceedings of the 21st European Colloquium of Arachnology, St.-Petersburg, 4–9 August 2003. *Arthropoda Selecta*, Special Issue No. 1: 67–74.
- Dunlop, J. A. 2007. A large parasitengonid mite (Acari, Erythraeoidea) from the Early Cretaceous Crato Formation of Brazil. *Fossil Record*, 10: 91–98.
- Dunlop, J. A. & Anderson, L. I. 2005. A fossil harvestman (Arachnida, Opiliones) from the Mississippian of East Kirkton, Scotland. *Journal of Arachnology*, 33: 482–489.
- Dunlop, J. A. & Bertrand, M. 2011. Fossil labidostomatid mites (Prostigmata: Labidostommatidae) from Baltic amber. *Acarologia*, 51: 191–198.
- Dunlop, J. A. & Braddy, S. J. 2011. *Cteniza bavincourti* and the nomenclature of arachnid related trace fossils. *The Journal of Arachnology*, 39: 250–257.
- Dunlop, J. A. & Brauckmann, C. 2006. A new trigonotarbid from the Coal Measures of Hagen Vorhalle, Germany. *Fossil Record*, 9: 130–136.
- Dunlop, J. A. & Giribet, G. 2003. The first fossil cyphophthalmid (Arachnida, Opiliones) from Bitterfeld amber, Germany. *The Journal of Arachnology*, 31: 371–378.
- Dunlop, J. A. & Horrocks, C. A. 1996. A new Upper Carboniferous whip scorpion (Arachnida: Uropygi: Thelyphonida) with a revision of the British Carboniferous Uropygi. *Zoologischer Anzeiger*, 234: 293–306.
- Dunlop, J. A. & Horrocks, C. A. 1997. Phalangiotarbid arachnids from the Coal Measures of Lancashire, UK. *Geological Magazine*, 134: 369–381.
- Dunlop, J. A. & Jekel, D. 2009. Nomenclatural notes on fossil spiders. *Bulletin of the British arachnological Society*, 14: 357–360.
- Dunlop, J. A. & Martill, D. M. 2002. The first whipspider (Arachnida: Amblypygi) and three new whipscorpions (Arachnida: Thelyphonida) from the Lower Cretaceous Crato Formation of Brazil. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 92: 325–334.
- Dunlop, J. A. & Mammitzsch, L. 2010. A new genus and species of harvestman from Baltic amber. *Palaeodiversity*, 3: 23–32.

- Dunlop, J. A. & Mitov, P. G.. 2011. The first fossil cyphophthalmid harvestman from Baltic amber. *Arachnologische Mitteilungen*, 40: 47–54.
- Dunlop, J. A. & Penney, D. 2012. *Fossil arachnids*. Siri Scientific Press, Manchester, 192 pp.
- Dunlop, J. A. & Poschmann, M. 1997. On the Emsian (Lower Devonian) arthropods of the Rhenish Schiefergebirge: 1. *Xenarachne*, an enigmatic arachnid from Willwerath, Germany. *Paläontologische Zeitschrift*, 71: 231–236.
- Dunlop, J. A. & Rößler, R. 2003. An enigmatic, solifuge-like fossil arachnid from the Lower Carboniferous of Kamienna Góra (Intra-Sudetic Basin), Poland. *Paläontologische Zeitschrift*, 77: 389–400.
- Dunlop, J. A. & Selden, P. A. 2004. A trigonotarbid arachnid from the Lower Devonian of Tredomen, Wales. *Palaeontology*, 47: 1469–1476.
- Dunlop, J. A. & Selden, P. A. 2013. Scorpion fragments from the Silurian of Powys, Wales. *Arachnology*, 16: 27–32.
- Dunlop, J. A., Anderson, L. I. & Braddy, S. J. 1999. A new chasmataspid (Chelicerata: Chasmataspida) from the Lower Devonian of the Midland Valley of Scotland. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 89: 161–165.
- Dunlop, J. A., Anderson, L. I. & Braddy, S. J. 2004. A redescription of *Chasmataspis laurencii* Caster & Brooks (Chelicerata: Chasmataspidida) from the Middle Ordovician of Tennessee, USA, with remarks on chasmataspid phylogeny. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 94: 207–205.
- Dunlop, J. A., Bartel, C. & Mitov, P. G. 2012. An enigmatic spiny harvestman from Baltic amber. *Fossil record*, 15: 91–101.
- Dunlop, J. A., Harms, D., Penney, D. 2008. A fossil tarantula (Araneae: Theraphosidae) from Miocene Chiapas amber, Mexico. *Revista Ibérica de Aracnología*, 15: 9–17.
- Dunlop, J. A., Kontschán, J. & Zwanzig, M. 2013. Fossil mesostigmatid mites (Mesostigmata: Gamasina, Microgyniina, Uropodina), associated with longhorn beetles (Coleoptera: Cerambycidae) in Baltic amber. *Naturwissenschaften*, DOI 10.1007/s00114-013-1031-8.
- Dunlop, J. A., Sempf, C. & Wunderlich, J. 2010. A new opilioacarid mite in Baltic amber. In Nentwig, W., Entling, M. & Kropf, C. (eds). *European Arachnology 2008*, pp. 59–70.
- Dunlop, J. A., Wunderlich, J. & Poinar Jr., G. O. 2004. The first fossil opilioacariform mite (Acari: Opilioacariformes) and the first Baltic amber camel spider (Solifugae). *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 94: 261–273.
- Dunlop, J. A., Anderson, L. I., Kerp, H. & Hass, H. 2004. A harvestman (Arachnida: Opiliones) from the Early Devonian Rhynie cherts, Aberdeenshire, Scotland. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 94: 341–354.

- Dunlop, J. A., Fayers, S. F., Hass, H. & Kerp, H. 2006. A new arthropod from the early Devonian Rhyndie chert, Aberdeenshire (Scotland), with a remarkable feeding device in the mouthparts. *Paläontologische Zeitschrift*, 80: 296–306.
- Dunlop, J. A., Wirth, S., Penney, D., McNeil, A., Bradley, R. S., Withers, P. J. & Preziosi, R. F. 2012. A minute fossil phoretic mite recovered by phase-contrast X-ray computed tomography. *Biology Letters*, 8: 475–460.
- Ebert, T. 1892. *Prestwichia (Euproops) scheeleana*. – *Abhandlung und Jahrbuch Königlich Preussische Geologisches Landesanstalt*, 10: 215–220.
- Edgecombe, G. D. 1998. Early myriapodous arthropods from Australia: *Maldybulakia* from the Devonian of New South Wales. *Records of the Australian Museum*, 50: 293–314.
- Ehlers, G. M. 1935. A new eurypterid from the Upper Devonian of Pennsylvania. *Contributions from the Museum of Palaeontology, University of Michigan*, 4 (18): 291–295.
- Eichwald, E. 1854. Die Grauwackenschichten von Liv- und Esthland. *Bulletin de la Société Imperiale des Naturalistes de Moscou*, 27: 1–211.
- Eichwald, E. 1860. *Lethaea Rossica. Vol. 1. Seconde section de l'ancienne Période*. Librairie et Imprimerie de E. Schweizerbart, Stuttgart, 1657 pp.
- Eldredge, N. 1974. Revision of the suborder Synziphosurina (Chelicerata, Merostomata), with remarks on merostome phylogeny. *American Museum Novitates*, 2543: 1–41.
- Elias, M. K. 1936. Character and significance of the late Paleozoic flora of Garnett, Kansas. *Journal of Geology*, 44: 9–23.
- Eller, E. R. 1938a. A review of the xiphosuran genus *Belinurus* with the description of a new species, *B. alleganyensis*. *Annals of the Carnegie Museum*, 27: 129–150.
- Eller, E. R. 1938b. A new xiphosuran, *Euproops morani*, from the Upper Devonian of Pennsylvania. *Annals of the Carnegie Museum*, 27: 152–153.
- Eller, E. R. 1940. *Belinurus carteri* a new xiphosuran from the Upper Devonian of Pennsylvania. *Annals of the Carnegie Museum*, 28: 133–136.
- Ellingsen, E. 1906. Report on the pseudoscorpions of the Guinea Coast (Africa) collected by Leonardo Fae. *Annali del Museo Civico de Storia Naturale di Genova*, (3)2: 243–265.
- Ellingsen, E. 1909. On some North American pseudoscorpions collected by Dr. F. Silvestri. *Bollettino del Laboratorio di Zoologia Generale e Agraria della R. Scuola sup. d'Agricoltura, Portici*, 3: 216–221.
- Elzinga, R. J. 1993. Larvamimidae, a new family of mites (Acari: Dermanyssoidea) associated with army ants. *Acarologia*, 34: 95–103.
- Emerton, J. H. 1875 Notes on spiders from Caves in Kentucky, Virginia and Indiana. *American Naturalist*, 9: 278–281.

- Emerton, J. H. 1882. New England spiders of the family Theridiidae. *Transactions of the Connecticut Academy of Arts and Sciences*, 6: 1–86.
- Eskov, K. Y. 1984. A new fossil spider family from the Jurassic of Transbaikalia from (Araneae: Chelicerata). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1984: 645–653.
- Eskov, K. Y. 1987. A new archaeid spider (Chelicerata: Araneae) from the Jurassic of Kazakhstan, with notes on the so-called “Gondwanan” ranges of recent taxa. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 175: 81–106.
- Eskov, K. Y. 1992. Archaeid spiders from Eocene Baltic amber (Chelicerata: Araneida: Arachaeidae) with remarks on the so-called “Gondwanan” ranges of Recent taxa. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 185: 311–328.
- Eskov, K. Y. & Marusik, Y. M. 1992. [Fossil spiders of the family Nesticidae.] *Palaeontologicheskii Zhurnal*, 2: 87–95. [In Russian]
- Eskov, K. Y. & Selden, P. A. 2005. First record of spiders from the Permian period (Araneae: Mesothelae). *Bulletin of the British arachnological Society*, 13: 111–116.
- Eskov, K. Y. & Wunderlich, J. 1995 (for 1994). On the spiders of the Taimyr ambers, Siberia, with the description of a new family and with general notes on the spiders from the Cretaceous resins. *Beiträge zur Araneologie*, 4: 95–107.
- Eskov, K. Y. & Zonstein, S. 1990. First Mesozoic mygalomorph spiders from the Lower Cretaceous of Siberia and Mongolia, with notes on the system and evolution of the infraorder Mygalomorphae (Chelicerata: Araneae). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 178: 325–368.
- Eskov, K. Y. & Zonstein, S. L. 2000. The first Ctenizoid Mygalomorph Spiders from Eocene Baltic amber (Araneida: Mygalomorphae: Ctenizidae). *Paleontological Journal*, 34: S268–S274. [Translated into English; original in Russian]
- Etheridge Jr., R. 1877. On the remains of a large crustacean, probably indicative of a new species of *Eurypterus*, or allied genus (*Eurypterus? Stevensoni*), from the Lower Carboniferous Series (Cementstone Group) of Berwickshire. *Quarterly Journal of the Geological Society*, 33: 223–228.
- Ewing, H. E. 1922. Studies on the taxonomy and biology of the tarsnemid mites, together with a note on the transformation of *Acarapis (Tarsonemus) woodi* Rennie (Acarina). *Canadian Entomologist*, 54: 104–113.
- Ewing, H. E. 1929. A synopsis of the American arachnids of the primitive order Ricinulei. *Annals of the Entomological Society of America*, 22: 583–600.
- Ewing, H. E. 1930. A fossil arachnid from the Lower Carboniferous shales (Pococno formation) of Virginia. *Annals*
- Fage, L. 1912. Etudes sur les araignées cavernicoles. I. Revision des Ochyroceratidae (n. fam.). *In Biospologica*, XXV. *Archives de Zoologie expérimentale et generale*, 10: 97–162.
- Fage, L. 1913. Etudes sur les Araignées cavernicoles. II. Revision des Leptonetidae. *In Biospologica*, XXIX. *Archives de Zoologie expérimentale et generale*, 10: 479–576.

- Fain, A. 1956. Une nouvelle famille d'acariens endoparasites des chauves-souris: Gastronyssidae fam. nov. *Annales de la Société Belge de Médecine Tropicale*, 36: 87–98.
- Fain, A. 1961. Une nouvelle famille d'acariens, parasites de serpents du genre *Mehelya* au Congo: Omentolaelaptidae Fam. nov. (Mesostigmata). *Revue de Zoologie et de Botanique Africaine*, 64: 283–296.
- Fain, A. 1967a. Nouveaux hypopes vivant dans les follicules pileux de Rongeurs américains. *Revue de Zoologie et de Botanique Africaine*, 76: 157–162.
- Fain, A. 1967b. Un acarien remarquable récolté sur un Tarsier (Heterocoptidae f.n. : Sarcoptiformes). – *Zoologischer Anzeiger*, 178: 90–94.
- Fain, A. 1968. Deux nouveaux Acariens Cavernicoles du Gabon (Sarcoptiformes). *Revue Biologia Gabonica*, 4: 195–205.
- Fain, A. 1974. Acariens récoltés par le Dr. J. Travé aux îles subantarctiques. I. Familles Saprogllyphidae et Hyadesidae (Astigmata). *Acarologia*, 16: 684–708.
- Fayers, S. R., Dunlop, J. A. & Trewin, N. H. 2005. A new early Devonian trigonotarbid arachnid from the Windyfield chert, Rhynie, Scotland. *Journal of Systematic Palaeontology*, 2: 269–284.
- Feider, Z. 1955. Arachnida, Acarina Trombidioidea. *Fauna RPR*, 5: 1–187.
- Feider, Z. & Vasiliu, N. 1969. Révision critique de la famille des Nicoletiellidae. In Proc. 2nd International Congress of Acarology, Sutton Bonington, England 1967. Acad. Kiado, Budapest: pp. 202–207.
- Feldmann, R. M., Schweitzer, C. E., Dattilo, B. & Farlow, J. O. 2011. Remarkable preservation of a new genus and species of limuline horseshoe crab from the Cretaceous of Texas, USA. *Palaeontology*, 54: 1337–1346.
- Feldmann, R. M., Vega, F. J., Applegate, S. P., & Bishop, G. A. 1998. Early Cretaceous arthropods from the Tlayua Formation at Tepexi de Rodriguez, Puebla, México. *Journal of Paleontology*, 72: 79–90.
- Fet, V. & Bechly, G. 2001. Case 3120a. Liochelidae, fam. nov. (Scorpiones): proposed introduction as a substitute name for Ischnuridae Simon, 1879, as an alternative to the suggested emendment of Ischnurinae Fraser, 1957 (Insecta, Odonata) to Ischnurinae in order to remove homonymy. *Bulletin of Zoological Nomenclature*, 58: 280–281.
- Fischer de Waldheim, G. 1839. Notes sur un crustacé fossile du genre *Eurypterus* de Podolie. *Bulletin de la Societe Imperiale des Naturalistes de Moscou*, 11: 125–128.
- Flower, R. H. 1945. A new Deepkill eurypterid. *American Midland Naturalist*, 34: 717–719.
- Flower, R. [H.] 1969. Merostomes from a Cotter horizon of the El Paso Group. *New Mexico Bureau of Mines and Mineral Resources Memoir*, 22: 35–44.
- Fraipont, J. 1889. Euryptérides nouveaux du Dévonien Supérieur de Belgique (Psammites du Condroz). *Annales de la Société Géologique de Belgique*, 17: 53–62.

- Forsslund, K.-H. 1941. Schwedische Arten der Gattung *Suctobelba* Paoli (Acari, Oribatei). *Zoologiska bidrag fran Uppsala*, 20: 381–396.
- Forsslund, K.-H. 1947. Über die Gattung *Autogneta* Hull (Acari, Oribatei). *Zoologiska bidrag fran Uppsala*, 25: 111–117.
- Forsslund, K.-H. 1956. Schwedische Oribatei (Acari). III. *Entomologisk Tidskrift*, 77: 210–218.
- Forster, R. R. 1948. A new sub-family and species of New Zealand Opiliones. *Records of the Auckland Institute and Museum*, 3: 313–318.
- Forster, R. R. 1954. The New Zealand harvestmen (sub-order Laniatores). *Canterbury Museum Bulletin*, 2: 1–329.
- Forster, R. R. 1955. A new family of spiders of the sub-order Hypochilomorphae. *Pacific Science*, 9: 277–285.
- Forster, R. R. & Forster, L. 1999. *Spiders of New Zealand and their worldwide kin*. University of Otago Press, Dunedin, vi + 270 pp.
- Forster, R. R. & Platnick, N. I. 1984. A review of archaeid spiders and their relatives, with notes on the superfamily Palpimanoidea (Arachnida: Araneae). *Bulletin of the American Museum of Natural History*, 178: 1–106.
- Forster, R. R. & Wilton, C. L. 1973. The spiders of New Zealand. Part IV. *Otago Museum Bulletin*, 4: 1–309.
- Frič, A. 1873. Fauna der Steinkohlenformation Böhmens. *Archiv für Naturwissenschaftliche Landesdurchforschung von Böhmen*, 2(2): 1–16.
- Frič, A. 1899a. On *Prolimulus woodwardi*. *Geological Magazine*, 6: 57–58.
- Frič, A. 1899b. *Fauna der Gaskohle und der Kalksteine der Permformation Böhmens*. Vol. IV: pp. 33–64.
- Frič, A. 1901. *Fauna der Gaskohle und der Kalksteine der Permformation Böhmens*. Vol. IV, part 2. Myriopoda pars II. Arachnoidea, pp. 56–63, pls 153, 154, Prague.
- Frič, A. 1904. *Palaeozoische Arachniden*. A Frič, Prague, 85 pp.
- Fritsch, K. von 1906. Beitrag zur Kenntnis der Tierwelt der deutschen Trias. *Abhandlungen der naturforschender Gesellschaft Halle*, 24: 220–285.
- Fry, W. G. 1978. A classification within the pycnogonids. *Zoological Journal of the Linnean Society*, 63: 35–58.
- García-Villafuerte, M. Á. 2006a. A new fossil *Episinus* (Araneae, Theridiidae) from Tertiary Chiapas amber, Mexico. *Revista Ibérica de Aracnología*, 13: 120–125.
- García-Villafuerte, M. Á. 2006b. Selenopidae y Thomisidae (Arachnida: Araneae) en ámbar de Chiapas, México. *Boletín Sociedad Entomológica Aragonesa*, 38: 209–212.
- García-Villafuerte, M. Á. 2008. Primer registro fósil del género *Hemirraghus* (Araneae, Theraphosidae) en ámbar del Terciario, Chiapas, México. *Revista Ibérica de Aracnología*, 16: 43–47.
- Garwood, R. J., Dunlop, J. A., Giribet, G. & Sutton, M. D. 2011. Anatomically modern Carboniferous harvestmen demonstrate early cladogenesis and stasis in Opiliones. *Nature Communications*, 2:444: 1–7.

- Gaud, J. & Ateyo, W. T. 1975. Gabuciniidae, famille nouvelle de Sarcoptiformes plumicoles. *Acarologia*, 16: 522–561.
- Gaud, J. & Ateyo, W. T. 1976. Ascouracarinae, n. sub-fam. des Sringobiidae, Sarcoptiformes plumicoles. *Acarologia*, 18: 143–162.
- Gaud, J. & Ateyo, W. T. 1977. A new name for *Ovacarus* and Ovacaridae (Acarina: Analgoidea). *Acarologia*, 18: 568–569.
- Gaud, J. & Ateyo, W. T. 1978. Nouvelles superfamilles pour les Acariens astigmatés parasites d'oiseaux. *Acarologia*, 19: 678–685.
- Gaud, J., Ateyo, W.T. & Berla, H.F. 1972. Acariens Sarcoptiformes plumicoles parasites des Tinamous. *Acarologia*, 14: 393–453.
- Gaud, J., Ateyo, W. T. & Klompen, J. S. H. 1989. Oconnoriidae, a new family of feather mites (Acarina, Pterolichoidea). *Journal of Entomological Science*, 24: 417–421.
- Geinitz, H. B. 1882. *Kreischeria wiedeii*, ein Pseudoskorpion aus der Steinkohlenformation von Zwickau. *Zeitschrift der Deutschen geologischen Gesellschaft*, 34: 238–242.
- Gerecke, R., Smith, I. M. & Cook, D. R. 1999. Three new species of *Apheviderulix* gen. nov. and proposal of Apheviderulicidae fam. nov. (Acari: Hydrachnidia: Eylaoidea). *Hydrobiologia*, 397: 133–147.
- Gerson, U. & Walter, D. E. 1998. Transfer of *Mecognatha* Wood from Stigmaeidae to Mecognathidae, fam. nov., a new synonymy, and a key to families of Raphignathoidea (Acari: Prostigmata). *Systematic and Applied Acarology*, 3: 145–147.
- Gerstaecker, C. E. A. 1863. Pantopoda. 248–350. In Carus, J. V. & Gerstaecker, C. E. A. (eds). *Handbuch der Zoologie*, 2. W. Engelmann, Leipzig, 642 pp.
- Gertsch, W. J. 1941. Report on some arachnids from Barro Colorado Island, Canal Zone. *American Museum Novitates*, 1146: 1–14.
- Gertsch, W. J. & Davis, L. I. 1946. Report on a collection of spiders from Mexico. V. *American Museum Novitates*, 1313: 1–11.
- Gervais, P. M. 1844. Remarques sur la famille des Scorpiones et descriptions des plusieurs espèces nouvelles de la collection du Muséum. *Archives du Muséum d'Histoire Naturelle, Paris*, 4: 201–240.
- Giebel, C. G. 1856. *Die Insekten und Spinnen der Vorwelt mit steter Berücksichtigung der lebenden Insekten und Spinnen; monographisch dargestellt*. Leipzig, 511 pp.
- Gill, E. L. 1909. An arachnid from the Coal Measures of the Tyne Valley. *Transactions of the Natural History Society of Northumberland, Durham and Newcastle-upon-Tyne, new series*, 3(2): 3–16.
- Gill, E. L. 1911. A Carboniferous arachnid from Lancashire. *Geological Magazine*, 48: 395–398.
- Gill, E. L. 1924. Fossil arthropods from the Tyne Coalfield. *Geological Magazine*, 61: 445–471.
- Giribet, G. & Dunlop, J. A. 2005. First identifiable Mesozoic harvestman (Opiliones: Dyspnoi) from Cretaceous Burmese amber. *Proceedings of the Royal Society B*, 272: 1007–1013.

- Giribet, G., Tourhino, A. L., Shih, C.-k. & Ren, D. 2012. An exquisitely preserved harvestman (Arthropoda, Arachnida, Opiliones) from the Middle Jurassic of China. *Organisms, Diversity & Evolution*, 12: 51–56.
- Giribet, G., Sharma, P. P., Benavides, L. R., Boyer, S. L., Clouse, R. M., De Bivort, B. L., Dimitrov, D., Kawauchi, G. Y., Murienne, J., Schwendinger, P. J. 2012. Evolutionary and biogeographical history of an ancient and global group of arachnids (Arachnida: Opiliones: Cyphophthalmi) with a new taxonomic arrangement. *Biological Journal of the Linnean Society*, 105: 92–130.
- Gistel, J. 1848. *Naturgeschichte des Thierreichs für höhere Schulen*. Stuttgart, pp. 155–156.
- Gjelstrup, P. & Solhøy, T. 1994. Oribatid mites (Acari). In *The Zoology of Iceland*. *Steenstrupia*, (3) 57: 1–78.
- Glushenko, N. V. & Ivanov, V. K. 1961. [*Paleolimulus* from the Lower Permian of the Donetsk Basin.] *Paleontologiceskij Žurnal*, 1861: 128–130. [in Russian]
- Goldenberg, F. 1873. *Fauna Saraepontana Fossilis. Die fossilen Thiere aus der Steinkohlenformation von Saarbrücken. Erstes Heft*. Chr. Möllinger Verlag, Saarbrücken, 26 pp.
- Goodnight, J. C. & Goodnight, M. L. 1942. Phalangids from Central America and the West Indies. *American Museum Novitates*, 1184: 1–23.
- Gonzalez, R. H. 1978. A new species of xenocaligonellid mite from the Galapagos Islands (Acari). *Proceedings of the Entomological Society of Washington*, 80: 191–196.
- González-Sponga, M. A. 1997. Arácnidos de Venezuela. Una nueva familia, dos nuevos géneros y dos nuevas especies de Opiliones Laniatores. *Acta Biologica Venezuelica*, 17: 51–58.
- Gourret, P. 1887. Recherches sur les Arachnides tertiaires d'Aix en Provence. *Recueil Zoologique Suisse*, 4: 431–496.
- Grabau, A. W. 1920. A new species of *Eurypterus* from the Permian of China. *Bulletin of the Geological Survey of China*, 2: 61–68.
- Grandjean, F. 1931. Observations sur les Oribates (1^{re} Série). *Bulletin du Muséum National d'Histoire Naturelle*, 3: 131–144.
- Grandjean, F. 1932a. Observations sur les Oribates (3^e série). *Bulletin du Muséum National d'Histoire Naturelle*, 4: 292–306.
- Grandjean, F. 1932b. Au sujet des *Palaeacariformes* Trägårdh. *Bulletin du Muséum National d'Histoire Naturelle*, 4: 411–426.
- Grandjean, F. 1933. Études sur les Développement des Oribates. *Bulletin de la Société zoologique de France*, 58: 30–61.
- Grandjean, F. 1934. La notation des poils gastronomiques et des poils dorsaux du propodosoma chez les Oribates (Acariens). *Bulletin de la Société zoologique de France*, 59: 12–44.
- Grandjean, F. 1936a. Les Microzetidae n. fam. (Oribates). *Bulletin de la Société zoologique de France*, 61: 60–93.

- Grandjean, F. 1936b. Les Oribates de Jean Frédéric Hermann et de son père [Arachn. Acar.]. *Annales Société Entomologique de France*, 105: 27–110.
- Grandjean, F. 1936c. Observations sur les Oribates (10^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 8: 246–253.
- Grandjean, F. 1937. Le Genre *Pachygnathus* Dugès (*Alycus* Koch) (Acariens). Cinquième et dernière partie. *Bulletin du Muséum National d'Histoire Naturelle*, 9: 262–269.
- Grandjean, F. 1939. Quelques genres d'Acariens appartenant au groupe des Endeostigmata. *Annales des Sciences Naturelles – Zoologie et Biologie Animale, Série 11*, 2: 1–122.
- Grandjean, F. 1947a. Études sur les Smarisidae et quelques autres Êrythroïdes (Acariens). *Archives de Zoologie Expérimental et Générale*, 85: 1–126.
- Grandjean, F. 1947b. Les Enarthronota (Acariens). Première série. *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 8: 213–248.
- Grandjean, F. 1948. Les Enarthronota (Acariens). (2^e série). *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 10: 29–58.
- Grandjean, F. 1950. Les Enarthronota (Acariens). (3^e série). *Annales des Sciences Naturelles – Zoologie et Biologie Animale*, 12: 85–107.
- Grandjean, F. 1951. Observations sur les Oribates (22^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 23: 91–98.
- Grandjean, F. 1953. Observations sur les Oribates (25^e Série). *Bulletin du Muséum National d'Histoire Naturelle*, 25: 155–162.
- Grandjean, F. 1954a. Observations sur les Oribates (28^e série). *Bulletin du Muséum National d'Histoire Naturelle*, 26: 204–211.
- Grandjean, F. 1954b. Essai de classification des Oribates (Acariens). *Bulletin de la Société zoologique de France*, 78: 421–446.
- Grandjean, F. 1954c. Etude sur les Palaeacaroides (Acariens, Oribates). *Mémoires du Muséum National d'Histoire Naturelle*, 7: 179–274.
- Grandjean, F. 1956a. Sur deux espèces nouvelles d'oribates (Acariens) apparantées a *Oripoda elongata* Banks 1904. *Archives de Zoologie Expérimentale et Générale*, 93: 185–218.
- Grandjean, F. 1956b. Galumnidae sans carènes lamellaires (Acariens, Oribates), 1^{re} série. *Bulletin de la Société zoologique de France*, 81: 134–150.
- Grandjean, F. 1958a. *Perlohmannia dissimilis* (Hewitt) (Acarien, Oribate). *Mémoires du Muséum National d'Histoire Naturelle*, 16: 57–120.
- Grandjean, F. 1958b. *Charassobates cavernosus* Grandj. 1929 (Acarien, Oribate). *Mémoires du Muséum National d'Histoire Naturelle*, 16: 121–140.
- Grandjean, F. 1959. *Polypterozetes cherubin* Berl. 1916 (Oribate). *Acarologia*, 1: 147–180.

- Grandjean, F. 1960a. Les Mochlozetidae n. fam. (Oribates). *Acarologia*, 2: 101–148.
- Grandjean, F. 1960b. Les Autognetidae n. fam. (Oribates). *Acarologia*, 2: 575–609.
- Grandjean, F. 1961a. Les Plasmobatidae n. fam. (Oribates). *Acarologia*, 3: 96–129.
- Grandjean, F. 1961b. Les Amerobelbidae (Oribates). (1^{re} partie). *Acarologia*, 3: 303–343.
- Grandjean, F. 1963. Les Autognetidae (Oribates). Deuxième partie. *Acarologia*, 4: 632–689.
- Grandjean, F. 1965a. Nouvelles observations sur les Oribates (4^e série). *Acarologia*, 7: 91–112.
- Grandjean, F. 1965b. Oribates mexicains (2^e série). *Stelechobates megalotrichus* n.g., n.sp. *Acarologia*, 7: 532–563.
- Grandjean, F. 1965c. Complément à mon travail de 1953 sur la classification des Oribates. *Acarologia*, 7: 713–734.
- Grandjean, F. 1966. Les Staurobatidae n. fam. (Oribates). *Acarologia*, 8: 696–727.
- Grandjean, F. 1967. Nouvelles observations sur les Oribates (5^e série). *Acarologia*, 9: 242–272.
- Grandjean, F. 1969. Considérations sur le classement des Oribates. Leur division en 6 groupes majeurs. *Acarologia*, 11: 127–153.
- Grandjean, F. 1970. Nouvelles observations sur les Oribates (8^e série). *Acarologia*, 12: 849–876.
- Grassi, B. & Calandruccio, S. 1885. Intorno ad un nuovo Aracnide Artrogastro (*Koenenia mirabilis* [sic]) che crediamo rappresentante d'un nuovo ordine (Microteliphonida). *Naturalista Siciliano*, 4: 127–133, 162–168.
- Griffiths, D. A. 1977. A new family of astigmatid mites from the Iles Crozet, sub-Antarctica, introducing a new concept relating to ontogenetic development of idiosomal setae. *Journal of Zoology, London*, 182: 291–308.
- Griswold, C., Audisio, T. & Ledford, J. 2012. An extraordinary new family of spiders from caves in the Pacific Northwest (Araneae, Trogloraptoridae, new family). *ZooKeys*, 215: 77–102.
- Gromov, A.V. 1998. [A new family, genus and species of scorpions (Arachnida, Scorpiones) from southern Central Asia.] – *Zoologicheskyy Zhurnal*, 77: 1003–1009. [In Russian.]
- Grote, A. R. & Pitt, W. H. 1875. I. Description of a new Crustacean from the Water Lime Group at Buffalo. *Bulletin of the Buffalo Society of Natural Sciences*, 3: 1–2.
- Gross, W. 1933. Die unterdevonischen Fische und Gigantostraken von Overath. *Abhandlungen der Preußischen Geologischen Landesanstalt (N. F.)*, 145: 41–77.
- Guérin-Méneville, F. E. 1839. Gastéracanthes scuptée et de Feisthamel, nouvelles espèces d'aranéides. *Revue zoologique*. 1839: 109–111.
- Guthörl, P. 1934. Die Arthropoden aus dem Carbon und Perms des Saar-Nahe-Pfalz-Gebietes. *Abhandlungen der Preußischen Geologischen Landesanstalt (N.F.)*, 164: 1–219.
- Guthörl, P. 1938. *Eophrynus waechteri* n. sp. (Arac., Anthracom.) aus der Tiefbohrung Stangenmühle, Saar-Karbon. *Senckenbergiana*, 20: 465–470.

- Guthörl, P. 1964. Zur Arthropoden-Fauna des Karbons und Perms. 20. Neue Arachniden-Funde (Anthracom.) aus dem Westfal A des Aachener Karbons. *Paläontologische Zeitschrift*, 38: 98–103.
- Guthörl, P. 1965. Zur Arthropoden-Fauna des Karbons und Perms. 19. Weiteres über die Arachniden aus dem Westfal und Stefan des saar-lothringischen und pfälzischen Karbons. *Annales Universitatis Saraviensis*, 4: 10–24.
- Haase, E. 1890. Beitrag zur Kenntniss der fossilen Arachniden. *Zeitschrift der Deutsche geologische Gesellschaft*, 1890: 629–657.
- Haeckel, E. 1866. *Generale Morphologie der Organismen. Band 2*. Berlin, 574 pp.
- Hadži, J. 1931. Skorpionreste aus dem tertiären Sprudelsinter von Böttingen (Schwäbische Alb). *Paläontologische Zeitschrift*, 13: 134–148.
- Hadži, J. 1935. Ein eigentümlicher neuer Höhlen-Opilionid aus Nord-Amerika, *Cladonychium corii* g.n. sp. n. *Biologia Generalis*, 11: 49–72.
- Halbert, J. N. 1915. Clare Island Survey, 39. Acarinida. Section II. Terrestrial and marine Acarina. *Proceedings of the Royal Irish Academy*, 31: 45–136.
- Hall, J. 1859. *Natural History of New York: Palaeontology, III*. New York State Museum, 532 pp.
- Hall, C. E. 1877. Contributions to Palaeontology from the Museum of the Second Geological Survey. *Proceedings of the American Philosophical Society*, 16: 621??.
- Hall, J. 1884a. Description of a New Species of *Stylonurus* from the Catskill Group. *New York State Museum (36th Annual Report)*: 76–77.
- Hall, J. 1884b. Note on Eurypteridae of the Devonian and Carboniferous Formations of Pennsylvania, with a supplementary note on the *Stylonurus excelsior*. *Proceedings of the American Association for the Advancement of Science*, 33: 420–422.
- Hall, J. 1884c. Eurypteridae from the Lower Productive Coal Measures in Beaver County, and the Lower Carboniferous Pithole Shale in Venango County. 2nd *Geological Survey of Pennsylvania. Report of Progress PPP*: 23–39.
- Halliday, R. B. 2006. New taxa of mites associated with Australian termites (Acari: Mesostigmata). *International Journal of Acarology*, 32: 27–38.
- Hall, J. & Clarke, J. M. 1888. *Paleontology of New York*. New York, 236 pp.
- Hall, J. & Clarke, J. M. 1888. Trilobites and other Crustacea of the Oriskany, Upper Helderberg, Hamilton, Portage, Chemung, and Catskill Groups. *Geological Survey of the State of New York, Palaeontology*, 7: xxx.
- Hammer, M. 1966. Investigations on the Oribatid Fauna of New Zealand, Part 1. *Biologiske Skrifter udgivet af Det Kongelige Danske Videnskabernes Selskab*, 15(2): 1–108.
- Hammer, M. 1967. Some oribatids from Kodiak Island near Alaska. *Acta Arctica*, 14: 5–25.

- Hammer, M. 1973. Oribatids from Tongatapu and Eua, the Tonga Islands, and from Upolu, Western Samoa. *Biologiske Skrifter udgivet af Det Kongelige Danske Videnskabernes Selskab*, 20(3): 1–70.
- Hansen, H. J. 1894. Arthrogastra Danica: en monographisk fremstilling af de i Danmark levende Meiere og Mosskorpioner med bidrag til sidstnaevnte underordens systematic. *Naturhistorisk Tidsskrift*, (3) 14: 491–554.
- Hansen, H. J. & Sørensen, W. 1904. *On two orders of Archanida*. Cambridge University Press, Cambridge, xi + 178 pp.
- Harger, O. 1874. Notice of a new spider from the Coal Measures of Illinois. *American Journal of Science*, 7: 219–223.
- Harlan, R. 1834. Critical notices of various organic remains hitherto discovered in North America. *Transactions of the Geological Society of Pennsylvania*, 1: 46–112.
- Harvey, M. S. 1990. Pezidae, a new freshwater mite family from Australia (Acarina: Halacaroidea). *Invertebrate Taxonomy*, 3: 771–781.
- Harvey, M. S. 1991. *Catalogue of the Pseudoscorpionida*. Manchester University Press, Manchester, vi + 726.
- Harvey, M. S. 1992. The phylogeny and classification of the Pseudoscorpionida (Chelicerata: Arachnida). *Invertebrate Taxonomy*, 6: 1373–1435.
- Harvey, M. S. 2002. Nomenclatural notes on Solifugae, Amblypygi, Uropygi and Araneae (Arachnida). *Records of the Western Australian Museum*, 20: 449–459.
- Harvey, M. S. 2003. *Catalogue of the smaller arachnid orders of the world*. CSIRO Publishing, Collingwood VC, xi + 385 pp.
- Harvey, M. A. & Selden, P. A. 1995. *Nyranytarbus*, replacement name for *Hemiphrynus* Frič, 1901 (Trigonotarvida: Eophrynidae). *Bulletin of the British arachnological Society*, 10: 74.
- Haupt, H. 1956. Beitrag zu Kenntnis der eözanen Arthropodenfauna des Geiselthals. *Nova Acta Leopoldina n.s.*, 128: 1–90.
- Haupt, H. 1957. Eine spinnenartige Arthropode aus dem Rotliegenden: *Rhabdotarchooides simoni* n. gen. n. sp. *Hallesches Jahrbuch für Mitteldeutsche Erdgeschichte*, 2(4): 246–247.
- Haupt, J. 1983. Vergleichende Morphologie der Genitalorgane und Phylogenie der liphistomorphen Webspinnen (Araneae: Mesothelae). I. Revision der bisher bekannten Arten. *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 21: 275–293.
- Hauschke, N. & Wilde, V. 1987. *Paleolimulus fuchsbergensis* n. sp. (Xiphosura, Merostomata) aus der oberen Trias von Nordwestdeutschland, mit einer Übersicht zur Systematik und Verbreitung rezenter Limuliden. *Paläontologische Zeitschrift*, 61: 87–108.
- Hauschke, N. & Wilde, V. 1989. Ein Limulide aus dem Zechstein (Oberes Perm) der Korbacher Bucht (Hessen, Bundesrepublik Deutschland). *Geologisches Jahrbuch Hessen*, 117: 17–21.

- Hauschke, N. & Wilde, V. 2000. Limulidenreste aus dem Unteren Buntsandstein (Benberg-Formation) von Beesenlaublingen (Sachsen-Anhalt). *Hallesches Jahrbuch für Geowissenschaften, Reihe B*, 22: 87–90.
- Hauschke, N. & Wilde, V. 2004. Palaeogene limulids (Xiphosura) from Saxony-Anhalt (Germany) – systematics and palaeobiogeography. *Hallesches Jahrbuch für Geowissenschaften, Reihe B*, 18: 161–168.
- Hauschke, N. & Wilde, V. 2008. Limuliden aus dem Oberen Buntsandstein von Süddeutschland. *Hallesches Jahrbuch für Geowissenschaften*, 30: 21–26.
- Hauschke, N., Osterink, H. W. & Wilde, V. 2009. Erster Nachweis eines Limuliden (Xiphosura, Limulacea) im Muschelkalk von Winterswijk (Niederlande). *Der Aufschluss*, 60: 13–23.
- Hauschke, N., Wilde, V. & Brauckmann, C. 2004. Triassic limulids from Madagascar – missing links in the distribution of Mesozoic Limulacea. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 2004(2): 87–94.
- Hauschke, N., Wilde, V. & Pietrzeniuk, E. 1992. Ein Limulide aus dem Muschelkalk (mittlere Trias) von Rüdersdorf bei Berlin. *Zeitschrift für geologische Wissenschaft*, 20: 461–466.
- Hedgpeth, J. W. 1978. A reappraisal of the Palaeopantopoda with description of a species from the Jurassic. *Zoological Journal of the Linnean Society*, 63: 23–34.
- Heer, O. 1865. *Die Urwelt der Schweiz*. Friedrich Schultheß, Zürich, xxix + 622 pp.
- Heetoff, M., Helfen, L. & Norton, R. A. 2009. Description of *Neoliodes dominicus* n. sp. (Acari, Oribatida) from Dominican Amber, aided by synchrotron X-ray microtomography. *Journal of Paleontology*, 83: 153–159.
- Heide, S. van der 1951. Les arthropodes du terrain houiller du Limbourg meridionale (excepte les scorpions et les insects). *Mededeelingen van de Geologische Stichting Serie C-IV-3 5*: 1–84.
- Heineken C. & Lowe R. T. 1832. Descriptions of two species of Araneidae, natives of Madeira. *Zool. Journ.*, 5: 320–323.
- Henderickx, H. 2005. A new *Geogarypus* from Baltic amber (Pseudoscorpiones: Geogarypidae). *Phegea*, 33: 87–92.
- Henderickx, H., Tafforeau, P. & Soriano, C. 2012. Phase contrast synchrotron microtomography reveals the morphology of a partially visible new *Pseudogarypus* in Baltic amber (Pseudoscorpiones: Pseudogarypidae). *Palaeontologia Electronica*, 15: 2;17A, 11 p.
- Henderickx, H., Cnudde, V., Masschaele, B., Dierick, M., Vlassenbroeck, J. & Hoorebeke, L. van 2006. Description of a new fossil *Pseudogarypus* (Pseudoscorpiones: Pseudogarypidae) with the use of X-ray micro-CT to penetrate opaque amber. *Zootaxa*, 1305: 41–50.
- Hentz, N. M. 1832. On North American spiders. *American Journal of Science*, 21: 99–109.
- Hentz, N. M. 1845. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History*, 5: 189–202.
- Hentz, N. M. 1847. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History* 5: 443–478.

- Hentz, N. M. 1850. Descriptions and figures of the Araneides of the United States. *Boston Journal of Natural History*, 6: 18–35, 271–295.
- Herbst, J. F. W. 1798. *Naturgeschichte der Ungeflügelten Insekten. Zweytes Heft*. Berlin, xx pp.
- Hermann, J. F. 1804. *Mémoire Apterologique*. F. G. Levrault, Strasbourg, 144 pp.
- Heyden, C. H. G. von 1826. Versuch einer systematischen Eintheilung der Acariden. *Isis von Oken*, 18: 609–613.
- Heyden, C. H. G. von 1859. Fossile Insekten aus der Rheinischen Braunkohle. *Palaeontographica*, 8: 1–15.
- Hibbert, S. 1836. On the fresh-water limestone of Burdiehouse in the neighbourhood of Edinburgh belonging to the Carboniferous Group of rocks. With supplementary notes on other fresh-water limestones. *Transactions of the Royal Society of Edinburgh*, 13: 169–282.
- Hickman, V. V. 1931. A new family of spiders. *Proceedings of the Zoological Society of London (B)*, 1931: 1321–1328.
- Hickman, V. V. 1944. On some new Australian Apneumonomorphae with notes on their respiratory system. *Papers and Proceedings of the Royal Society of Tasmania*, 1943: 179–195.
- Hickmann, V. V. 1945. A new group of apneumone spiders. *Transactions of the Connecticut academy of Arts and Sciences*, 36: 135–148.
- Hickman, V. V. 1949. Tasmanian littoral spiders with notes on their respiratory systems, habits and taxonomy. *Papers and Proceedings of the Royal Society of Tasmania*, 1948: 31–43.
- Hickman, V. V. 1957. A fossil spider from Tertiary resin from Allendale Victoria. *Proceedings of the Royal Society of Victoria, N.S.*, 69: 25–27.
- Hilton, W. A. 1942. Pantopoda (continued) II. Family Callipallenidae. *Journal of Entomology and Zoology, Pomona College, Claremont*, 34: 38–41.
- Hirschmann, W. 1971. A fossil mite of the genus *Dendrolaelaps* (Acarina, Mesostigmata, Digamasellidae) found in amber from Chiapas, Mexico. *University of California Publications in Entomology*, 63: 69–70.
- Hirst, S. 1923. On some arachnid remains from the Old Red Sandstone (Rhynie Chert bed, Aberdeenshire). *Annals and Magazine of Natural History, Series 9*, 12: 455–474.
- Hoek, P. C. C. 1881. Report on the Pycnogonida dredged by HMS Challenger 1873–76. *Reports of the Scientific Results of the Exploring Vessel HMS Challenger*, 3(10): 1–167.
- Hoff, C. C. 1963. Sternophorid pseudoscorpions, chiefly from Florida. *American Museum Novitates*, 1875: 1–36.
- Holl, F. 1829. *Handbuch der Peterefactenkunde*. Hilscher, Dresden, 489 pp.
- Holland F. D., Jr., Erickson, J. M. & O'Brien, D. E. 1975. *Casterolimulus*: a new Late Cretaceous generic link in Limulid lineage. Studies in Paleontology and Stratigraphy. *Bulletin of American Paleontology*, 62: 235–249.
- Holmberg, E. L. 1882. Observations à propos du sous-ordre des araignées territoriales (Territelariae), spécialement du genre nordaméricain *Catadysas* Hentz et de la sous-famille Mecicobothrioidae, Holmberg. *Boletín de la Academia Nacional de Ciencias en Cordoba (Argentina)*, 4: 153–174.

- Holmberg, E. L. 1883. *Neothereutes darwini* Holmb., representante de una nueva familia de Citrigradas. *Boletín de la Academia Nacional de Ciencias en Cordoba (Argentina)*, 5: 35–48.
- Hong Y.-c. 1982. [Study on new spider genus in amber.] *Science in China*, 24(12): 1500–1515. [In Chinese]
- Hong Y.-c. 1983a. [Discovery of a Miocene scorpion from the diatoms of Shanwang in Shandong Province.] *Bulletin of the Tianjin Institute of Geology and Mineral Resources*, 8, 17–21. [In Chinese]
- Hong Y.-c. 1983b. [Discovery of new fossil pseudoscorpiononods in amber.] *Bulletin of the Tianjin Institute of Geology and Mineral Resources*, 8: 24–29. [In Chinese]
- Hong Y.-c. 1984. Arachnida. 185–187 In Tianjin Institute of Geology and Mineral Resources (eds). *Palaeontological Atlas of North China II. Mesozoic Volume*. Geological Publishing House, Beijing. [In Chinese with English summary]
- Hong Y.-c. 1985. *Fossil Insects, scorpionids and araneids in the diatoms of Shanwang*. Geological Publishing House, Beijing, 80 pp.
- Hopkins, D. M., Gitterman, R. E. & Matthews, J. V. 1976. Interstadial mammoth remains and associated pollen and insect fossils, Kotzebue Sound area, northwestern Alaska. *Geology*, 4: 169–173.
- Huang D.-y., Selden, P. A. & Dunlop, J. A. 2009. Harvestmen (Arachnida: Opiliones) from the Middle Jurassic of China. *Naturwissenschaften*, 96: 955–962.
- Huber, B. A. 2003. Southern African pholcid spiders revision and cladistic analysis of *Quamtana* gen. nov. and *Spermophora* Hentz (Araneae: Pholcidae), with notes on male–female covariation. *Zoological Journal of the Linnean Society*, 139: 477–527.
- Huber, B. A. & Wunderlich, J. 2006. Fossil and extant species of the genus *Leptopholcus* in the Dominican Republic, with the first cases of egg-parasitism in pholcid spiders (Araneae: Pholcidae). *Journal of Natural History*, 40: 2341–2360.
- Hull, J. E. 1920. The spider family Linyphiidae: an Essay in Taxonomy. *Vasculum*, 6: 7–11.
- Hünicken, M. A. 1980. A giant fossil spider (*Megarachne servinei*) from Bajo de Véliz, Upper Carboniferous, Argentina. *Boletín de la Academia Nacional de Ciencias, Córdoba*, 53: 317–341.
- Hunter, J. R. S. 1886. Notes on the discovery of a fossil scorpion (*Paleophonus caledonicus*) in the Silurian strata of Logan water. *Transactions of the Geological Society of Glasgow*, 8: 169–170.
- Jackson, R. T. 1906. A new species of fossil *Limulus* from the Jurassic of Sweden. *Arkiv för Zoologi*, 3(11): 1–7.
- Jaekel, O. 1914. Ein großer *Pterygotus* aus dem rheinischen Unterdevon. *Palaeontologische Zeitschrift*, 1: 379–382.
- Jävi, T. H. 1912/14. Das Vaginalsystem der Sparassiden. *Annales Academiae Scientiarum Fennicae*, A4: 1–248.
- Jell, P. A. & Duncan, P. M. 1986. Invertebrates, mainly insects, from the freshwater Lower Cretaceous Koonwarra fossil bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association of Australian Palaeontology*, 3: 111–205.

- Jeram, A.J. 1994a. Scorpions from the Viséan of East Kirkton, West Lothian, Scotland, with a revision of the infraorder Mesoscorpionina. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 84: 283–299.
- Jeram, A.J. 1994b. Carboniferous Orthosterni and their relationship to living scorpions. *Palaeontology*, 37: 513–550.
- Jocqué, R. 1994. Halidae, a new spider family from Madagascar (Araneae). *Bulletin of the British arachnological Society*, 9: 281–289.
- Jocqué, R. 2001. Chummidae, a new spider family (Arachnida, Araneae) from South Africa. *Journal of Zoology, London*, 254: 481–493.
- Jones, T. R. & Woodward, H. 1888. On some Scandanavian Phyllocarida. *Geological Magazine, New Series, Decade 3*, 5: 145–150.
- Jones, T. R. & Woodward, H. 1899. Contributions to fossil Crustacea. *Geological Magazine, New Series, Decade 4*, 6: 388–395.
- Jordan, H. & Meyer, H. von 1854. Ueber die Crustaceen der Steinkohlenformation von Saarbrücken. *Palaeontographica*, 4: 1–15.
- Judson, M. [L. I.] 2003. Baltic amber pseudoscorpions (Arachnida: Chelonethi): a new species of *Neobisium* (Neobisiidae) and the status of *Obisium rathkii* Koch and Berendt. *Geodiversitas*, 25: 445–450.
- Judson, M. L. I. 2007. First fossil record of the pseudoscorpion family Pseudochiridiidae (Arachnida, Chelonethi, Cheirioidea) from Dominican amber. *Zootaxa*, 1393: 45–51.
- Judson, M. L. I. 2009. Cheliferoid pseudoscorpions (Arachnida, Chelonethi) from the Lower Cretaceous of France. *Geodiversitas*, 31: 61–71.
- Judson, M. L. 2010. Redescription of *Chelifer eucarpus* Dalman (Arachnida, Chelonethi, Withiidae) and first records of pseudoscorpions in copal from Madagascar and Colombia. *Palaeodiversity*, 3: 33–42.
- Judson, M. L. I. & Mağol, J. 2009. A mite of the family Tanaupodidae (Arachnida, Acari, Parasitengona) from the Lower Cretaceous of France. *Geodiversitas*, 31: 41–47.
- Judson, M. [L. I.] & Wunderlich, J. 2003. Rhagidiidae (Acari, Eupodoidea) from Baltic amber. *Acta zoologica cracoviensa*, 46 (suppl.–Fossil Insects): 147–152.
- Jux, U. 1982. *Somaspidion hammapheron* n.gen. n.sp. – ein Arachnide aus dem Oberkarbon der subvaristischen Saumsenke NW Deutschlands. *Paläontologische Zeitschrift*, 56: 77–86.
- Kaddumi, H. F. 2007. *Amber of Jordan: the oldest prehistoric insects in fossilized resin. Second edition*. Eternal River Museum of Natural History, Amman, Jordan, 224 pp.
- Karpinen, E. & Koponen, M. 1973. The subfossil oribatid fauna of Piilonsuo, a bog in southern Finland. *Annales entomologici Fennici*, 39: 22–32.
- Karpinen, E. & Koponen, M. 1974. Further observations on subfossil remains of oribatids (Acar., Oribatei) and insects in Piilonsuo, a bog in southern Finland. *Annales entomologici Fennici*, 40: 172–175.

- Karpinen, E., Krivolutsky, D. A., Koponen, M., Kozlovskaja, L. S., Laskova, L. M. & Viitasaari, M. 1979. List of subfossil oribatid mites (Acarina, Oribatei) of northern Europe and Greenland. *Annales entomologici Fennici*, 45: 103–108.
- Karsch, F. 1879. Arachnologische Beiträge. *Zeitschrift für die gesammten Naturwissenschaften*, 52: 534–562.
- Karsch, F. 1880a. Arachnologische Blätter. I. Ueber *Corinna* (C. L. Koch) und ihre Verwandtschaften. *Zeitschrift für die gesammten Naturwissenschaften*, 53: 373–378.
- Karsch, F. 1880b. Arachnologische Blätter. X. Scorpionologische Fragmente. *Zeitschrift für die gesammten Naturwissenschaften*, 53: 404–409.
- Karsch, F. 1882. Ueber ein neues Spinnenthier aus der Schlesischen Steinkohle und die Arachnoiden überhaupt. *Zeitschrift der Deutschen geologischen Gesellschaft*, 34: 556–561.
- Karsch, F. 1884. Neue Milben in Bernstein. *Berliner Entomologische Zeitschrift*, 28: 175–176.
- Keegan, H. L., Yunker, C. E. & Baker, E. W. 1960. Malaysian parasites. XLVI. *Hystrichonyssus turneri*, n.sp. n.g. representing a new subfamily of Dermasyddidae (Acarina) from a Malayan porcupine. *Studies from the Institute for Medical Research Federation of Malaya*, 107: 455–473.
- Keferstein, C. 1834. *Die Naturgeschichte des Erdkörpers in ihren ersten Grundzügen*, Vol. 2. F. Fleischer, Leipzig, 896 pp.
- Keifer, H. H. 1966. [untitled.] *Californian Department of Agriculture. Eriophyid Series*, B-21: 1–20.
- Keirans, J. E., Lane, R. S. & Cauble, R. 2002. A series of larval *Amblyomma* species (Acari : Ixodidae) from amber deposits in the Dominican Republic. *International Journal of Acarology*, 28: 61–66.
- Kethley, J. B. 1974. Developmental chaetotaxy of a paedomorphic celaenopsoid, *Neotenogynium malkini* n.g., sp. (Acari: Parasitiformes: Neotenogyniidae, n. fam.) associated with millipedes. *Annals of the Entomological Society of America*, 67: 571–579.
- Kethley, J. B. 1977a. The Status of *Hybolicus* Berlese, 1913 and *Oehserchestes* Jacot, 1939 (Acari: Acariformes: Endeostigmata). *Fieldiana Zoology*, 72: 59–64.
- Kethley, J. B. 1977b. An unusual Parantennuloid, *Philodana johnstoni* n.g., n.sp. (Acari: Parasitiformes: Philodanidae, n. fam.) associated with *Neatus tenebrioides* (Coleoptera: Tenebrionidae) in North America. *Annals of the Entomological Society of America*, 70: 487–494.
- Kethley, J. B. 1979. A cladistic analysis of the Trignyaspida (Acari: Parasitiformes) with a review of the higher categories and nominate taxa. In Piffel, E. (ed). *Proceedings of the 4th International Congress of Acarology – Saalfelden (Austria)*. Akadémiai Kiadó, Budapest, pp. 459–466.
- Kethley, J. B. 1989. Proteonematolycidae (Acari: Acariformes), a new mite family from fore-dune sand of Lake Michigan. *International Journal of Acarology*, 15: 209–217.
- Kethley, J. B., Norton, R. A., Bonamo, P. M. & Shear, W. A. 1989. A terrestrial alicorhagiid mite (Acari: Acariformes) from the Devonian of New York. *Micropaleontology*, 35: 367–373.

- Kew, H. W. 1911. A synopsis of the false scorpions of Britain and Ireland. *Proceedings of the Royal Irish Academy (B)*, 29: 38–64.
- Keyserling, E. 1877. Ueber amerikanische Spinnenarten der Unterordnung Citigradae. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 26: 609–708.
- Keyserling, E. 1880a. *Die Spinnen Amerikas, I. Laterigradae*. Nürnberg, 1, 283 pp.
- Keyserling, E. 1880b. Neue Spinnen aus Amerika. I. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 29: 293–349.
- Keyserling, E. 1882. Neue Spinnen aus Amerika. III. *Verhandlungen der Zoologisch-Biologischen Gesellschaft in Wien*, 31: 269–314.
- Keyserling, E. 1884. *Die Spinnen Amerikas. Theridiidae*. Nürnberg, 2, 222 pp.
- Khaustov A. A. 2000. Bembidiacaridae, a new family of mites (Acari: Heterostigmata) associated with carabid beetles of the genus *Bembidion* (Coleoptera: Carabidae). *Acarina*, 8: 3–8.
- Khaustov, A. A. & Perkovsky, E. E. 2010. The first fossil record of mites of the family Pyemotidae (Acari: Heterostigmata), with description of a new species from Rovno Amber. *Palaeontological Journal*, 44: 418–421.
- Khaustov, A. A. & Poinar jr., G. O. 2010. *Protoresinacarus brevipedis* gen. n., sp. n. from Early Cretaceous Burmese amber: the first fossil record of mites of the family Resinacaridae (Acari: Heterostigmata: Pyemotoidea). *Historical Biology*: 23: 219–222.
- Kirchner, H. 1923. *Limulus Sandbergi* n. sp. aus dem fränkischen oberen Buntsandstein (Plattensandstein). *Centralblatt für Mineralogie, Geologie und Paläontologie*, 20: 634–639.
- Kim, C. M. 2008. Euphysalozirconidae, a new mesostigmatid mite family (Acari: Mesostigmata: Trigynaspida: Aenictequoidea). *Acarologia*, 48: 33–38.
- Kim, J.-p. & Nam, K.-s. 2008. [Mesozoic spider (Araneae: Pisauridae) from Korea.] *Korean Arachnology*, 24: 119–125. [in Korean with English summary]
- Kim, J.-p. & Nam, K.-s. 2008. [Mesozoic spider (Araneae: Lycosidae) from China.] *Korean Arachnology*, 28: 35–45. [in Korean with English summary]
- Kishida, K. 1930. A new scheme of classification of spider families and genera. *Lansania*, 2: 33–43.
- Kjellesvig-Waering, E. N. 1934. Note on a new eurypterid from the Moscow Shales of New York. *American Journal of Science, 5th Series*, 27: 386–387.
- Kjellesvig-Waering, E. N. 1948a. Two new eurypterids from the Silurian of Indiana. *Journal of Paleontology*, 22: 465–472.
- Kjellesvig-Waering, E. N. 1948b. The Mazon Creek Eurypterid: A revision of the genus *Lepidoderma*. *Scientific Papers, Illinois*, 3(4): 1–48.
- Kjellesvig-Waering, 1950a. A new Silurian *Hughmilleria* from West Virginia. *Journal of Paleontology*, 24: 226–228.
- Kjellesvig-Waering, 1950b. A new Silurian Eurypterid from Florida. *Journal of Paleontology*, 24: 229–231.

- Kjellesvig-Waering, E. N. 1951. Downtonian (Silurian) Eurypterida from Perton, near Stoke Edith, Herefordshire. *Geological Magazine*, 88: 1–24.
- Kjellesvig-Waering, E. N. 1954. Note on a new Silurian (Downtonian) scorpion from Shropshire, England. *Journal of Palaeontology*, 28: 485–486.
- Kjellesvig-Waering, E. N. 1955. A new phyllocarid and eurypterid from the Silurian of Florida. *Journal of Paleontology*, 29: 295–297.
- Kjellesvig-Waering, E. N. 1958. The genera, species and subspecies of the family Eurypteridae Burmeister, 1845. *Journal of Paleontology*, 32: 1107–1148.
- Kjellesvig-Waering, E. N. 1959. A taxonomic review of some late Paleozoic Eurypterida. *Journal of Palaeontology*, 33: 251–256.
- Kjellesvig-Waering, E. N. 1961a. Eurypterida of the Devonian Holland Quarry Shale of Ohio. *Fieldiana, Geology*, 14(5): 79–98.
- Kjellesvig-Waering, E. N. 1961b. The Silurian Eurypterida of the Welsh Boderland. *Journal of Paleontology*, 35: 251–256.
- Kjellesvig-Waering, E. N. 1963a. Revision of some Upper Devonian Stylonuridae (Eurypterida) from New York and Pennsylvania. *Journal of Paleontology*, 37: 490–495.
- Kjellesvig-Waering, E. N. 1963b. Pennsylvanian invertebrates of the Mazon Creek area, Illinois, Eurypterida. *Fieldiana, Geology*, 14(9): 169–197.
- Kjellesvig-Waering, E. N. 1964a. A synopsis of the Family Pterygotidae Clarke and Ruedemann 1912 (Eurypterida). *Journal of Paleontology*, 38: 331–361.
- Kjellesvig-Waering, E. N. 1964b. Eurypterida: Notes on the subgenus *Hughmilleria* (*Nanahughmilleria*) from the Silurian of New York. *Journal of Paleontology*, 38: 410–412.
- Kjellesvig-Waering, E. N. 1966b. Silurian scorpions of New York. *Journal of Paleontology*, 40: 359–375.
- Kjellesvig-Waering, E. N. 1966c. The scorpions of Trinidad and Tobago. *Caribbean Science*, 6: 123–135.
- Kjellesvig-Waering, E. N. 1969. A new phalangiotarbid (Arachnida) from the Pennsylvanian of Oklahoma. *Journal of Paleontology*, 43: 1280–1282.
- Kjellesvig-Waering, E. N. 1971. A new Downtonian stylonurid from Central England (Silurian, Eurypterida). *Journal of Paleontology*, 45: 538–539.
- Kjellesvig-Waering, E. N. 1972. *Brontoscorpius anglicus*: a giant Lower Palaeozoic scorpion from central England. *Journal of Paleontology*, 46: 39–42.
- Kjellesvig-Waering, E. N. 1973. A new Silurian *Slimonia* (Eurypterida) from Bolivia. *Journal of Paleontology*, 47: 549–550.
- Kjellesvig-Waering, E. N. 1979. Eurypterids. In Jaanusson, V., Laufeld, S. & Skoglund, R. (eds). Lower Wenlock faunal and floral dynamics – Vattenfallet section, Gotland. *Sveriges Geologiska Undersökning, Serie C*, NR 762, Årsbok 73 NR, 3: 121–136.

- Kjellesvig-Waering, E. N. 1986. A restudy of the fossil Scorpionida of the world. *Palaeontographica Americana*, 55: 1–287.
- Kjellesvig-Waering, E. N. & Caster, K. E. 1955. The Pterygotidae of the Silurian Vernon Shales of New York. *Journal of Paleontology*, 29: 1041–1047.
- Kjellesvig-Waering, E. N. & Heubusch, C. A. 1962. Some Eurypterida from the Ordovician and Silurian of New York. *Journal of Paleontology*, 36: 211–221.
- Kjellesvig-Waering, E. N. & Leutze, W. P. 1966. Eurypterida from the Silurian of West Virginia. *Journal of Paleontology*, 40: 1109–1122.
- Kjellesvig-Waering, E. N. & Størmer, L. 1952. The *Dolichopterus*–*Strobilopterus* group in the Eurypterida. *Journal of Paleontology*, 26: 659–661.
- Klompen, H. & Grimaldi, D. 2001. First Mesozoic record of a parasitiform mite: a larval argasid tick in Cretaceous amber (Acari: Ixodida: Argasidae). *Annals of the Entomological Society of America*, 94: 10–15.
- Kobayashi, T. 1933. On the occurrence of Xiphosuran remains in Chosen (Korea). *Japanese Journal of Geology and Geography*, 10: 175–182.
- Koçak, A. Ö. & Kemal, M. 2008. New synonyms and replacement names in the genus group taxa of Araneida. *Centre for entomological Studies, Miscellaneous Papers*, 139-140: 1–4.
- Koch, C. L. 1829–1844. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Fortgesetzt von Herrich-Schäffer, Hefte 111-190*. Regensburg. [1833, Hefte 119–121]
- Koch, C. L. 1834. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Hefte 122-125, 127*. Regensburg.
- Koch, C. L. 1835. Arachniden. In Panzer (ed). *Faunae Insectorum Germaniae initia. Hefte 128-131*. Regensburg.
- Koch, C. L. 1837. *Uebersicht des Arachnidensystems 1*. C. H. Zeh'sche Buchhandlung, Nürnberg, 39 pp.
- Koch, C. L. 1839a. *Uebersicht des Arachnidensystems 2*. C. H. Zeh'sche Buchhandlung, Nürnberg, 38 pp.
- Koch, C. L. 1839b. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Sechster Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 156 pp.
- Koch, C. L. 1839c. *Deutschlands Crustaceen, Myriapoden und Arachniden*. Hefte 23–30.
- Koch, C. L. 1842a. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Neunter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 108 pp.
- Koch, C. L. 1842b. *Uebersicht des Arachnidensystems 3*. C. H. Zeh'sche Buchhandlung, Nürnberg, 130 pp.
- Koch, C. L. 1843a. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben. Zehnter Band*. C. H. Zeh'sche Buchhandlung, Nürnberg, 142 pp.
- Koch, C. L. 1843b. *Uebersicht des Arachnidensystems 3*. C. H. Zeh'sche Buchhandlung, Nürnberg, 130 pp [continuation of 1842b; see above].
- Koch, C. L. 1844. Systematische Übersicht über die Ordnung der Zecken. *Archiv für Naturgeschichte*, 1: 217–239.

- Koch, C. L. 1846. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben*. Dreizehnter Band. C. H. Zeh'sche Buchhandlung, Nürnberg, 234 pp.
- Koch, C. L. 1847. *Die Arachniden. Getreu nach der Natur abgebildet und beschrieben*. Vierzehnter Band. C. H. Zeh'sche Buchhandlung, Nürnberg, 210 pp.
- Koch, C. L. 1851. *Übersicht des Arachnidensystems* 5. C. H. Zeh'sche Buchhandlung, Nürnberg, 104 pp.
- Koch, C. L. & Berendt, G. C. 1854. Die im Bernstein befindlichen Myriapoden, Arachniden und Apteren der Vorwelt. In Berendt, G. C. *Die in Bernstein befindlichen organischen Reste der Vorwelt gesammelt in Verbindung mit mehreren bearbeitet und herausgegeben* 1. Berlin, Nicolai, 124 pp.
- Koch, L. 1866. *Die Arachniden-Familie der Drassiden*. 1–6. J. L. Lotzbeck, Nürnberg, 352 pp.
- Koch, L. 1871–1883. *Die Arachniden Australiens nach der Natur beschrieben und abgebildet*. Bauer & Raspe, Nürnberg, 1489 pp.
- Koch, L. 1873. *Uebersichtliche Darstellung der europäischen Chernetiden (Pseudoscorpione)*. Bauer und Raspe, Nürnberg, xx pp.
- Kraepelin, K. 1899. Zur Systematik der Solifugen. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg*, 16: 195–258.
- Kraepelin, K. 1901. Palpigradi und Solifugae. *Tierreich*, 12: i–x, 1–159.
- Kraepelin, K. 1905. Die geographische Verbreitung der Skorpione. - *Zoologische Jahrbücher, Abtheilung für Systematik*, 22: 321–364.
- Kramer, P. 1885. Ueber Halarachne Halichoeri, Allm. *Zeitschrift für Naturwissenschaften*, 58: 1–31.
- Krause, T., Hauschke, N. & Wilde, V. 2009. Ein Limulide aus den Gelben Basisschichten des Oberen Muschelkalks von Ohrdruf bei Gotha (Thüringen). *Geowissenschaftliche Mitteilungen von Thüringen*, 13: 163–168.
- Kratochvíl, J. 1958. Höhlenweberknechte Bulgariens (Palpatores – Nemastomatidae). *Acta Academiae Scientiarum Čechoslovenicae Basis Brunensis*, 30: 523–576.
- Krivolutsky, D. A. & Krasilov, B. A. 1977. Oribatid mites from Upper Jura deposits of USSR. 16–24. In Skarlato, O. A. & Balashov, Y. S. (eds) *Morphology and Diagnostics of Mites*. Zoological Institute, Leningrad, 85 pp. [in Russian]
- Krüger, J. & Dunlop, J. A. 2010. Schizomids (Arachnida: Schizomida) from Dominican Republic amber. *Alavesia*, 3: 43–53.
- Kues, B. S. & Kietzke, K. K. 1981. A large assemblage of a new eurypterids from the Red Tanks Member, Madera Formation (Late Pennsylvania - Early Permian) of New Mexico. *Journal of Paleontology*, 55: 709–729.
- Kühl, G., Poschmann, M. & Rust, J. 2013. A ten-legged sea spider (Arthropoda: Pycnogonida) from the Lower Devonian Hunsrück Slate (Germany). *Geological Magazine*, 150: 556–564.

- Kühl, G., Bergamnn, A., Dunlop, J. A., Garwood, R. J. & Rust, J. 2012. Redescription and palaeobiology of *Palaeoscorpius devonicus* Lehmann, 1944 from the Lower Devonian Hunsrück Slate of Germany. *Palaeontology*, 55: 775–787.
- Kulczynski, L. 1902. Species Oribatarum (Oudms.) (Damaeinarum Michael) in Galicia collectae. *Dissertationum mathematicarum et physicarum Academiae Litterarum Cracoviensis*, 42: 1–50.
- Kulicka, R. 1990. The list of animal inclusions in Baltic amber from collection of the Museum of Earth in Warsaw. *Prace Muzeum Ziemi*, 41: 144–146.
- Kury, A. B. 2003. Annotated catalogue of the Laniatores of the New World (Arachnida, Opiliones). *Revista Ibérica de Aracnología*, Volumen especial monográfico 1: 1–337.
- Kury, A. B. & Pérez González, A. 2002. A new family of Laniatores from northwestern South America (Arachnida, Opiliones). *Revista Ibérica de Aracnología*, 6: 3–11.
- Kušta, J. 1883. *Anthracomartus krejci*, eine neue Arachnide aus dem Böhmischem Karbon. *Sitzungsberichte der Königlich Böhmischem Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1883: 7.
- Kušta, J. 1884a. Neue Arachniden aus der Steinkohlenformation von Rakonitz. *Sitzungsberichte der Königlich Böhmischem Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1884: 398–401.
- Kušta, J. 1884b. *Thelyphonus bohemicus* n. sp., ein fossiler Geisselscorpion aus der Steinkohlenformation von Rakonitz. *Sitzungsberichte der Königlich Böhmischem Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1884: 186–191.
- Kušta, J. 1885. Neue fossile Arthropoden aus dem Noeggarathienschiefer von Rakonitz. *Sitzungsberichte der Königlich Böhmischem Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1885: 1–7.
- Kušta, J. 1888. O nových arachnidech z karbonu Rakovnického. (Neue Arachniden aus der Steinkohlenformation bei Rakonitz). *Sitzungsberichte der Königlich Böhmischem Gesellschaft der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 1888: 194–208.
- Kutorga, S. 1838. *Beitrag zur Kenntnis der organischen Überreste des Kupfersandsteins am westlichen Abhange des Urals*. St. Petersburg, 38 pp.
- Kuznetsov, N. N., Khaustov, A. A. & Perkovsky, E. E. 2010. First record of mites of the family Stigmaeidae (Acari, Raphignathoidea) from Rovno amber with description of a new species of the genus *Mediolata*. *Vestnik zoologii*, 44: 545–547.
- Lamarck, J. B. P. A. 1801. *Système des animaux sans vertèbres*. Lamarck and Deterville, Paris, xx pp.
- Lamont, A. 1955. Scottish Silurian Chelicerata. *Transactions of the Edinburgh Geological Society*, 16: 200–216.
- Lamsdell, J. C. 2011. The eurypterid *Stoermeropterus conicus* from the Lower Silurian of the Pentland Hills, Scotland. *Monographs of the Palaeontographical Society*, 165: 1–84.

- Lamsdell, J. C. 2013a. Revised systematics of the Palaeozoic 'horseshoe crabs' and the myth of the monophyletic Xiphosura. *Zoological Journal of the Linnaen Society*, 167: 1–27.
- Lamsdell, J. C. 2013b. Redescription of *Drepanopterus pentlandicus* Laurie, 1892, the earliest known mycteropoid (Chelicerata: Eurypterida) from the early Silurian (Llandovery) of the Pentland Hills, Scotland. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 103: 77–103.
- Lamsdell, J. C. & Selden, P. A. 2013. Babes in the wood – a unique window into sea scorpion ontogeny. *BMC Evolutionary Biology* 13: 98.
- Lamsdell, J. C., Braddy, S. J. & Tetlie, O. E. 2010. The systematics and phylogeny of the Stylonurina (Arthropoda: Chelicerata: Eurypterida). *Journal of Systematic Palaeontology*, 8: 49–61.
- Lamsdell, J. C., Hoşgör, İ & Selden, P. A. 2013. A new Ordovician eurypterid (Arthropoda: Chelicerata) from southeast Turkey: evidence for a cryptic Ordovician record of Eurypterida. *Gondwana Research*, 23: 354–366.
- Lamsdell, J. C., Xue, J.-h. & Selden, P. A. 2013. A horseshoe crab (Arthropoda: Chelicerata: Xiphosura) from the Lower Devonian (Lochkovian) of Yunnan, China. *Geological Magazine*, 150: 367–370.
- Lamsdell, J. C., Braddy, S. J., Loeffler, E. J. & Dineley, D. L. 2010. Early Devonian stylonurine eurypterids from Arctic Canada. *Canadian Journal of Earth Sciences*, 47: 1405–1415.
- Lane, R. S. & Poinar jr., G. O. 1986. First fossil tick (Acari: Ixodidae) in new world amber. *International Journal of Acarology*, 12: 75–78.
- Latreille, P. A. 1795. Observations sur la variété des organes de la bouche des tiques, et distribution méthodique des insectes de cette famille d'après les caractères établis sur la conformation de ces organes. *Magasin Encyclopédique, ou Journal des Sciences, des Lettres et des Arts*, 4: 15–20.
- Latreille, P. A. 1796. *Précis de caractères génériques des insectes, disposés dans un ordre naturel*. Prévot, Paris, xx pp.
- Latreille, P. A. 1802. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*. Dufart, Paris, xx pp.
- Latreille, P. A. 1804a. Tableau méthodique des Insectes. *Nouveau Dictionnaire d'histoire naturelle*, 24: 129–200.
- Latreille, P. A. 1804b. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes, Vol. 7*. F. Dufart, Paris, pp. 144–305.
- Latreille, P. A. 1806. *Genera Crustaceorum et Insectorum. Vol. 1*. A. Koenig, Paris, pp. 82–127.
- Latreille, P. A. 1809. *Genera Crustaceorum et Insectorum. Vol. 4*. Paris, pp. 73–371.
- Latreille, P. A. 1810. *Considérations générales sur l'Ordre Naturel des Animaux composant les Classes des Crustacés, des Arachnides et des Insectes*. Paris, 446 pp.
- Latreille, P. A. 1819. [Articles sur les Araignées]. *Nouveau Dictionnaire d'histoire naturelle* 30-35 : ?? pp.
- Latreille, P. A. 1829. Les Arachnides. In Cuvier, G (ed.) *Le règne animal, nouv. ed.* Paris, pp. 206–291.

- Laurentiaux-Viera, F. & Laurentiaux, D. 1961. *Prothelyphonus neerlandicus*, nov. sp., Uropyge du Westphalien du Limbourg Hollandais. *Mededelingen van de Geologische Stichting, N.S.*, 13: 29–34.
- Laurentiaux-Viera, F. & Laurentiaux, D. 1963. Sur quelques restes nouveaux d'Arachnides du terrain houiller. *Annales de la Société Géologique du Nord*, 83: 23–29.
- Laurie, M. 1892. On some eurypterid remains from the Upper Silurian rocks of the Pentland Hills. *Transactions of the Royal Society of Edinburgh*, 37: 151–162.
- Laurie, M. 1896. Further notes on the anatomy and development of scorpions, and their bearing on the classification of the order. *Annals and Magazine of Natural History, series 6*, 17: 185–193.
- Laurie, M. 1899. On a Silurian scorpion and some additional eurypterid remain from the Pentland Hills. *Transactions of the Royal Society of Edinburgh*, 39: 575–590.
- Leach, W. E. 1815. A tabular view of the external characters of four classes of animals which Linné arranged under Insecta; with the distribution of the genera composing three of these classes into orders, andc. And descriptions of several new genera and species. *Transactions of the Linnean Society of London*, 11: 306–400.
- Leach, W. E. 1819. *Dictionnaire des Sciences Naturelles, Vol. 14*. Paris, pp. 537–538.
- Leary, R.L. 1980. *Labriscorpio alliedensis*, a new Carboniferous scorpion from Rock Island County, Illinois. *Journal of Paleontology*, 54: 1255–1257.
- Leech, R. & Matthews Jr., J. V. 1971. *Xysticus archaeopalpus* (Arachnida: Thomisidae), a new species of crab spider from Pliocene sediments in western Alaska. *Canadian Entomologist*, 103: 1337–1340.
- Lehmann, W.M. 1944. *Palaeoscorpius devonicus* n. g., n. sp., ein Skorpion aus dem rheinischen Unterdevon. *Neues Jahrbuch für Paläontologie, Monatshefte, B*: 177–185.
- Lehtinen, P. T. 1967. Classification of the cribellate spiders and some allied families, with notes on the evolution of the suborder Araneomorpha. *Annales Zoologici Fennici*, 4: 199–468.
- Lehtinen, P. T. 1981. New Holothyrina (Arachnida, Anactinotrichida) from New Guinea and South America. *Acarologia*, 22: 3–13.
- Lenz, H. 1886. Beiträge zur Kenntniss der Spinnenfauna Madagascars. *Zoologische Jahrbücher, Systematik*, 1: 379–408.
- Leutze, W. P. 1958. Eurypterids from the Silurian Tymochtee dolomite of Ohio. *Journal of Paleontology*, 32: 937–942.
- Leutze, W. P. 1961. Arthropods from the Syracuse Formation, Silurian of New York. *Journal of Paleontology*, 35: 49–64.
- Levy, G. 2007. The first troglobite scorpion from Israel and a new chactoid family (Arachnida: Scorpiones). *Zoology in the Middle East*, 40: 91–96.
- Li S.-q. & Wunderlich, J. 2008. Sinopimoidae, a new spider family from China (Arachnida, Araneae). *Acta zootaxonomica sinica*, 33: 1–6.

- Lin Q.-b., Zhang, Z.-f. & Wang, B.-z. 1989. New evidences for Miocene climatic optimum event—review on the Miocene spider fossils from Shanwang collection. *Proceedings of International Symposium on Pacific Neogene and Marine Events*. Nanjing University Press, pp. 137–147.
- Lin Q.-b., Yao Y.-m., Xiang W.-d. & Xia Y.-r. 1988. An Oligocene micropalaeontofauna from Gubei district of Shandong and its ecological environment. *Acta Micropalaeontologica Sinica*, 5: 331–345.
- Lindquist, E. E. & Moraza, M. L. 1993. Pyrosejidae, a new family of trigynaspid mites (Acari: Mesostigmata: Cercomegistina) from Middle America. *Acarologia*, 34: 283–307.
- Lindquist, E. E. & Palacios-Vargas, J. G. 1991. Proterorhagiidae (Acari: Endeostigmata), a new family of rhagidiid-like mites from Mexico. *Acarologia*, 32: 341–363.
- Linnaeus, C. 1758. *Systema naturae*, 10th edition. Vol 1. L. Salvii, Holmiae.
- Loman, J. C. C. 1900. Ueber die geographische Verbreitung der Opilioniden. *Zoologische Jahrbücher, Systematik*, 16: 71–104.
- Lourenço, W. R. 1995. Description de trois nouveaux genres et quatre nouvelles espèces de scorpions Buthidae de Madagascar. *Bulletin du Muséum National d'Histoire Naturelle (4)*, 17A: 95–106.
- Lourenço, W. R. 1996a. *Faune de Madagascar. 87. Scorpions (Chelicerata, Scorpiones)*. Muséum National d'Histoire Naturelle, Paris, 102 pp.
- Lourenço, W. R. 1996b. Premier cas connu d'un sub-fossile de scorpion dans le copal de Madagascar. *Compte Rendus de l'Académie des Sciences, Paris, Sér. Ila*, 323: 889–891.
- Lourenço, W. R. 1998. Panbiogeographie, les distributions disjointes et le concept de famille relictuelle chez les Scorpions. *Biogeographica*, 74: 133–144.
- Lourenço, W. R. 2000a. More about the Buthoidea of Madagascar, with special references to the genus *Tityobuthus* Pocock (Scorpiones, Buthidae). *Revue suisse de Zoologie*, 107: 721–736.
- Lourenço, W. R. 2000b. Premier cas d'un sub-fossile d'araignee appartenant au genre *Archaea* Koch and Berendt (Archaeidae) dans le copal de Madagascar. *Comptes rendus de l'Académie des Sciences Paris, Sciences de la Terre et des planets*, 330: 509–512.
- Lourenço, W. R. 2001. A remarkable scorpion fossil from the amber of Lebanon. Implications for the phylogeny of Buthoidea. *Comptes rendus de l'Académie des Sciences Paris, Sciences de la Terre et des planets*, 332: 641–646.
- Lourenço, W. R. 2002. The first scorpion fossil from the Cretaceous amber of Myanmar (Burma). New implications for the phylogeny of Buthoidea. *Comptes Rendus Palevol*, 1: 97–101.
- Lourenço, W. R. 2003. The first scorpion fossil from the Cretaceous amber of France. New implications for the phylogeny of Chactoidea. *Comptes Rendus Palevol*, 2: 213–219.
- Lourenço, W. R. 2004. Description of a further species of fossil scorpion in Baltic amber. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 3: 1886–1889.

- Lourenço, W. R. 2009a. A new sub-fossil scorpion of the genus *Microcharmus* Lourenço from Malagasy copal (Scorpiones, Microcharmidae). *Boletín Sociedad Entomológica Aragonesa*, 44: 135–137.
- Lourenço, W. R. 2009b. A new species of *Tityus* C. L. Koch, 1836 (subgenus *Brazilotityus* Lourenço, 2006) from the Dominican amber (Scorpiones: Buthidae). *Euscorpius*, 83: 1–5.
- Lourenço, W. R. 2012. Further considerations on scorpions found in Baltic amber, with a description of a new species (Scorpiones: Buthidae). *Euscorpius*, 146: 1–7.
- Lourenço, W. R. 2013. A new species of *Tityus* C. L. Koch, 1836 (Scorpiones: Buthidae) from Dominican amber. *Euscorpius*, 156: 1–5.
- Lourenço, W.R. & Gall, J.-C. 2004. Fossil scorpions from the Buntsandstein (Early Triassic) of France. *Comptes Rendus Palevol*, 3: 369–378.
- Lourenço, W. R. & Henderickx, H. 2012. Another new sub-fossil species of scorpion of the genus *Palaeogrosphus* Lourenço, 2000 from Malagasy copal (Scorpiones: Buthidae). *Euscorpius*, 137: 1–4.
- Lourenço, W. R. & Weitschat, W. 1996. More than 120 years after its description, the enigmatic status of the genus of the Baltic amber scorpion “*Tityus eogenus*” Menge, 1869 can finally be clarified. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 79: 183–188.
- Lourenço, W. R. & Weitschat, W. 2000. New fossil scorpions from the Baltic amber – implications for Cenozoic biodiversity. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 84: 247–260.
- Lourenço, W. R. & Weitschat, W. 2001. Description of another fossil scorpion from Baltic amber with considerations on the evolutionary levels of Cenozoic Buthoidea. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 85: 277–283.
- Lourenço, W. R. & Weitschat, W. 2005a. A new genus and species of fossil scorpion from a different kind of Baltic amber (Scorpiones, Buthidae). – *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89, 183–188.
- Lourenço, W. R. & Weitschat, W. 2005b. First sub-fossil scorpion of genus *Chactas* Gervais from Colombian copal (Scorpiones, Chactidae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89: 179–182.
- Lourenço, W. R. & Weitschat, W. 2009. A new species of *Palaeoananteris* Lourenço & Weitschat, 2001, fossil scorpion from Ukrainian amber (Scorpiones, Buthidae). *Boletín Sociedad Entomológica Aragonesa*, 45: 231–235.
- Lourenço, W. R., Henderickx, H. & Weitschat, W. 2005. A new genus and species of fossil scorpion from Baltic amber (Scorpiones, Buthidae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 89: 159–166.
- Lucas, H. 1835. Sur une monographie du genre Thélyphone. *Magasin de Zoologie*, 5: Classe VIII, pls. 8–10.

- Lucas, H. 1846. Histoire naturelle des Animaux articulés. *In Exploration scientifique de l'Algérie pendant les années 1840, 1841, 1842, publiée par ordre du Gouvernement et avec le concours d'une commission académique. Sciences physiques, Zoologie, 5 tomes, Paris, 1846–1850.* Vol. 1: 89–271.
- Luxton, M. 1985. Cryptostigmata (Arachnida: Acari): a concise review. *Fauna of New Zealand*, 7: 1–112.
- Luxton, M. 1988. A new mite superfamily (Acari: Cryptostigmata). *Zoological Journal of the Linnean Society*, 93: 71–91.
- Lyubarsky, G. Y. & Perkovsky, E. E. 2012. The first Eocene species of the genus *Cryptophagus* (Coleoptera, Clavicornia, Cryptophagidae). *Vestnik zoologii*, 46: 36–40.
- MacLeay, W. S. 1839. On some new forms of Arachnida. *Annals and Magazine of Natural History*, 2: 1–14.
- Magowski, W. Ł. 1994. Discovery of the first representative of the mite subcohort Heterostigmata (Arachnida: Acari) in the Mesozoic Siberian amber. *Acarologia*, 35: 229–241.
- Magowski, W. Ł. 1995. Fossil heterostigmatid mites in amber – 85 million year-old an arthropod mite Relationships 53–58. *in* Kropczynska, D., Boczek, J. & Tomczyk, A. (eds) *The Acari: Physiological and Ecological Aspects of Acari – Host Relationships* Dabor, Warsaw, 698 pp.
- Malz, H. & Poschmann, M. 1993. Erste Süßwasser-Limuliden (Arthropoda, Chelicerata) aus dem Rotliegenden der Saar-Nahe-Senke. *Osnabrücker naturwissenschaftliche Mitteilungen*, 19: 21–24.
- Mahnert, V. 1979. Pseudoskopione (Arachnida) aus dem Amazonas-Gebiet (Brasilien). *Revue suisse de Zoologie*, 86: 719–810.
- Mahunka, S. 1986. A survey of the family Carabodidae C. L. Koch, 1836 (Acari: Oribatida). *Acta Zoologica Hungarica*, 32: 73–135.
- Mahunka, S. 1990. A survey of the superfamily Euphthiracaroidea Jacot, 1930 (Acari: Oribatida). *Folia Entomologica Hungarica*, 51: 37–80.
- Martens, J. 1976. Genitalmorphologie, System und Phylogenie der Weberknechte (Arachnida: Opiliones). *Entomologica Germanica*, 3: 51–68.
- Martens, J. 1988. Fissiphalliidae, a new family of South American laniatorean harvestmen (Arachnida: Opiliones). *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 26: 114–127.
- Martin, W. 1809. *Petrificata Derbiensia* 1, Wigan, xx pp.
- Marusik, Y. M. & Penney, D. 2004. A survey of Baltic amber Theridiidae (Araneae) inclusions, with descriptions of six new species. *In* Logunov, D. V. & Penney, D (eds). *European Arachnology 2003 (Proceedings of the 21st European Colloquium of Arachnology, St.-Petersburg, 4–9 August 2003)*. *Arthropoda Selecta*, Special Issue No. 1: 201–208.
- Marx, G. 1888. On a new and interesting spider. *Entomologica Americana*, 4: 160–162.
- Marx, G. 1890a. Arachnida. *In* Howard, L. O. (ed.) *Scientific results of the explorations by the U. S. Fish Commission Steamer Albatross. No. V. – Annotated catalogue of the insects collected in 1887–'88. – Proceedings of the United States National Museum*, 12: 207–211.

- Marx, G. 1890b. Catalogue of the described Araneae of temperate North America. *Proceedings of the United States National Museum*, 12: 497–594.
- Matthew, G. F. 1888. On some remarkable organisms of the Silurian and Devonian rocks in Southern New Brunswick. *Transactions of the Royal Society of Canada*, 1888: 49–61.
- Matthew, G. F. 1895. Organic remains of the Little River Group, No. IV. *Transactions of the Royal Society of Canada*, 2nd Ser., 1: 273–279.
- McAlpine, J. F. & Martin, J. E. H. 1969. Canadian amber – a paleontological treasure chest. *Canadian Entomologist*, 101: 819–838.
- McCook, H. C. 1888. A new fossil spider, *Eoatypus woodwardii*. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1888: 200–202.
- Meek, F. B. 1867. Notes on a new genus of fossil Crustacea. *Geological Magazine, Decade 4*, xx: 320–321.
- Meek, F. B. & Worthen, A. H. 1865. Notice of some new types of organic remains from the Coal Measures of Illinois. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 17: 41–45.
- Meek, F.B & Worthen, A.H. 1868a. Preliminary notice of a scorpion, a *Eurypterus?* and other fossils from the Coal Measures of Illinois and Iowa. *American Journal of Science and Arts, series 2*, 45: 25.
- Meek, F.B. & Worthen, A.H. 1868b. Palaeontology of Illinois. In *Geological Survey of Illinois*, 3: 289–565.
- Melander, A. L. 1903. Some additions to the Carboniferous terrestrial fauna of Illinois. *Journal of Geology*, 11: 178–198.
- Melendez, B. 1971. Un novel Eurypteride du Westphalien des Asturies (NW Espagne). In Krefeld (ed.) *Septieme Congres de Stratigraphie et de Geologie du Carbonifere*, 3: 415–417.
- Mello-Leitão, C. F. de 1932. Notas sobre as Micratheneas do Brasil. *Anais do Academia Brasileira dos Ciências*, 4: 73–97.
- Mello-Leitão, C. F. de 1937. Distribution et Phylogénie des Faucheurs Sud-Américains. *Proceedings of the 12th International Congress of Zoology, Lisbon*, 2(5): 1217–1228.
- Mello-Leitão, C. F. de 1940. Arañas de las islas Juan Fernandez, recogidas por el Señor R. Wagenknecht. *Revista Chilena de Historia Natural*, 44: 236–239.
- Menge, A. 1854. Footnotes in Koch, C. L. & Berendt, G. C. Die im Bernstein befindlichen Myriapoden, Arachniden und Apteren der Vorwelt. In Berendt, G. C. *Die in Bernstein befindlichen organischen Reste der Vorwelt gesammelt in verbindung mit mehreren bearbeitet und herausgegeben 1*. Berlin, Nicolai, 124 pp.
- Menge, A. 1855. Ueber die Scheerenspinnen, Chernetidae. *Neueste Schriften der Naturforschenden Gesellschaft*, 5: 1–43.
- Menge, A. 1856. Lebenszeichen vorweltlicher, im Bernstein eingeschlossener Thiere. *Programm der Petrischule zu Danzig*, 8: 32 pp.
- Menge, A. 1866. Preussische Spinnen. Erste Abtheilung. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 1–152.

- Menge, A. 1868. Preussische Spinnen. II. Abtheilung. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 153–218.
- Menge, A. 1869. Ueber einen Scorpion und zwei Spinnen im Bernstein. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2: 1–9.
- Mesquita, M. V. 1996. *Cretaraneus matensnetoi* n.sp. (Araneoidea) da Formação Santana, Cretáceo Inferior da Bacia do Araripe. *Revista Universidade Guarulhos, Série Geociências*, 1(3): 24–31.
- Miller, S. A. 1874. Notes and descriptions of Cincinnatian Group fossils. *Cincinnati Quarterly Journal of Science*, 1: 343–351.
- Miller, S. A. & Gurley, W. F. E. 1896. New species of Echinodermata and a new crustacean from the Palaeozoic rocks. *Illinois State Museum Natural History Bulletin*, 10: 1–91.
- Millot, J. 1947. Une araignée malgache énigmatique, *Gallieniella mygaloides* n. g., n. sp. *Bulletin du Muséum National d'Histoire Naturelle, 2^e Série*, 19: 158–160.
- Millot, J. 1948. Faits nouveaux concernant les *Archaea* [Aranéides]. *Mémoires de l'Institut Scientifique de Madagascar*, 1(A1): 3–14.
- Moberg, J. C. 1892. Om en nyupptäckt fauna i block af kambrisk sandsten, insamlade af Dr N.O. Holst. *Geologiska Föreningens i Stockholm Förhandlingar*, 14: 103–120.
- Moore, J. I. 1923. A review of the present knowledge of fossil scorpions, with the description of a new species from the Pottsville Formation of Clay County, Indiana. *Proceedings of the Indiana Academy of Science*, 38: 125–134.
- Moore, R. A., McKenzie, S. C. & Lieberman, B. S. 2007. A Carboniferous synziphosurine (Xiphosura) from the Bear Gulch Limestone, Montana, USA. *Palaeontology*, 50: 1013–1019.
- Moore, R. A., Briggs, D. E. G., Braddy, S. J. & Shultz, J. W. 2011. Synziphosurines (Xiphosura: Chelicerata) from the Silurian of Iowa. *Journal of Paleontology*, 85: 83–91.
- Moore, R. A., McKenzie, S. C., Braddy, S. J., Anderson, L. I., Mikulic, D. G. & Kluesendorf, J. 2005. A new synziphosurine (Chelicerata: Xiphosura) from the Late Llandovery (Silurian) Waukesha Lagerstätte, Wisconsin, USA. *Journal of Paleontology*, 79: 242–250.
- Moran, R. J. 1986. The Sternodidae (Araneae, Araneomorpha), a new family of spiders from eastern Australia. *Bulletin of the British Arachnological Society*, 7: 87–96.
- Moraza, M. L. & Lindquist, E. E. 1999. Coprozerconidae, a new family of zerconoid mites from North America (Acari: Mesostigmata: Zerconoidea). *Acarologia*, 39: 291–313.
- Müller, O. F. 1785. *Entomotraca, seu, Insecta testacea quae in aquis Daniae et Norvegiae reperit, descripsit et iconibus illustravit*. Hauniae, Thiele, xx pp.
- Müller, A. H. 1957. Ein Arachnidenrest (*Brachylycosa* ? *manebachensis* n. sp.) aus dem Unteren Rotliegenden (Manebacher Schichten) von Thüringen. *Geologie*, 6: 95–98.

- Münster, G. Graf zu 1839. Die Rhyncholiten des Muschelkalks mit ihrem Fortsätzen. In Münster, G. Graf zu (ed.) *Beiträge zur Petrefacten-Kunde 1*: 48–51.
- Münster, G. Graf zu 1840. Über die fossilen Arten *Limulus* in den lithographischen Schiefern von Bayern. In Münster, G. Graf zu (ed.) *Beiträge zur Petrefacten-Kunde 3*: 26–27.
- Murray, A. 1877. *Economic Entomology, Aptera*. South Kensington Museum Handbooks, 433 pp.
- Nalepa, A. 1898. Eriophyidae (Phytoptidae). In *Das Tierreich. Eine Zusammenstellung und Kennzeichnung der rezenten Tierformen. 4. Lieferung. Acarina*. Deutsche Zoologische Gesellschaft, 4: 72 pp.
- Nicolet, H. 1855. Histoire naturelle des Acariens qui se trouvent aux environs de Paris. *Archives de Museum Nationale d'Histoire Naturelle de Paris*, 7: 381–482.
- Nieszkowski, J. 1859. Zusätze zur Monographie der Trilobiten der Ostseeprovinzen, nebst der Beschreibung einiger neuen obersilurischen Crustaceen. *Archiv für die Naturkunde Liv-, Ehst.- und Kurlands (Ser. 1)*, 1: 345–384.
- Nindel, F. 1955. Die tierischen Reste aus dem Karbon von Karl-Marx-Stadt und Hainichen i.S. *Geologie*, 4: 673–694.
- Nishikawa, Y. 1974. [Amber spiders from Mizunami, Japan.] *Bulletin of the Mizunami Fossil Museum*, 1: 401–406.
[in Japanese with English summary]
- Norton, R. A. 2006. First record of *Collohmanna* (*C. schusteri* n. sp.) and *Hermannia* (*H. sellnicki* n. sp.) from Baltic amber, with notes on Sellnick's genera of fossil oribatid mites (Acari: Oribatida). *Acarologia*, 46: 111–125.
- Norton R.A. & Metz, L. 1980. Nehypochthoniidae (Acari: Oribatei), a new family from the southeastern United States. *Annals of the Entomological Society of America*, 73: 54–62.
- Norton, R. A., Bonamo, P. N., Grierson, J. D. & Shear, W. A. 1988. Oribatid mite fossils from a terrestrial Devonian deposit near Gilboa, New York. *Journal of Paleontology*, 62: 259–269.
- Novojilov, N. J. 1959. Mérostomes du Dévonien inférieur et moyen de Sibérie. *Annales de la Société Géologique du Nord*, 78: 241–258.
- Novojilov, N. & Størmer, L. 1963. A new scorpion from the Upper Carboniferous of Siberia. *Norsk Geologisk Tidsskrift* 43: 83–87.
- O'Connell, M. 1916. The habitat of the Eurypterida. *Bulletin of the Buffalo Society of Natural Sciences*, 11: 1–278.
- Olivier, P. A. S. & Theron, P. D.. 2000. Pentapalpidae, a new family of eupodoid mites (Prostigmata:Eupodoidea) from South Africa. *Acarologia*, 40: 385–392.
- Ono, H. 1981. First record of a crab spider (Thomisidae) from Dominican amber (amber collection Stuttgart : Arachnida, Araneae). *Stuttgarter Beitrage zur Naturkunde (B)*, 73: 1–13.
- Opluštil, S. 1985. New findings of Arachnida from the Bohemian Upper Carboniferous. *Věstník Ústředního ústavu geologického*, 60: 35–42.

- Opluštil, S. 1986. *Promygaie janae* sp. n., the new anthracomartid (Arachnida) from the Upper Carboniferous of central Bohemia. *Věstník Ústředního ústavu geologického*, 61: 287–292.
- Oppenheim, P. 1887–1888. Die Insectenwelt des lithographischen Schiefers in Bayern. *Palaeontographica*, 34: 215–247.
- Orr, P. J., Siveter, D. J., Briggs, D. E. G., Siveter, D. J. & Sutton, M. D. 2000. A new arthropod from the Silurian Konservat-Lagerstätte of Herefordshire, UK. *Proceedings of the Royal Society B*, 267: 1497–1504.
- Oudemans, A. C. 1902. Classificatie der Acari. *Tijdschrift voor Entomologie*, 45: 50–64.
- Oudemans, A. C. 1909. Über die bis jetzt genauer bekannten Thrombidium-larven und über eine neue Klassifikation der Prostigmata. *Tijdschrift voor Entomologie*, 52: 19–61.
- Oudemans, A. C. 1916. Acarologische Aanteekeningen LX. *Entomologische berichten*, 4: 308–316.
- Oudemans, A. C. 1923. Studie over de sedert 1977 ontworpen system der Acari; nieuwe classificatie; phylogenerische beschouwingen. *Tijdschrift voor Entomologie*, 66: 49–85.
- Özdikmen, H. 2007. Nomenclatural changes for seven preoccupied spider genera (Arachnida: Araneae). *Munis Entomology & Zoology*, 2: 137–142.
- Packard, A. S. 1885. Types of Carboniferous Xiphosura new to North America. *American Naturalist*, 1885: 291–294.
- Packard, A. S. 1886. On the Carboniferous xiphosurous fauna of North America. *Memoirs of the National Academy of Sciences*, 3: 143–157.
- Page, D. 1856. *Advanced textbook of geology*. William Blackwood and Sons, Edinburgh, 326 pp.
- Page, D. 1859. *Advanced textbook of geology, 2nd edn*. William Blackwood and Sons, London, xx pp.
- Palmer, A. R. 1957. Miocene arthropods from the Mojave Desert California. *Geological Survey Professional Paper*, 294-G: 237–280.
- Pampaloni, L. 1902. I resti organici nel disodile di Melilli in Sicilia. *Palaeontographica Italica*, 8: 121–130.
- Patrick, R. R. 1989. A new phalangiotarbid (Arachnida) from the McLeansboro Group (Pennsylvanian) of Indiana. *Journal of Paleontology*, 63: 327–331.
- Peach, R. N. 1882. Further researches among Crustacea and Arachnida. *Transactions of the Royal Society of Edinburgh*, 30: 511–529.
- Peach, R. N. 1883. A new species of fossil scorpions from the Carboniferous rocks of Scotland and the English borders, with a review of the genera *Eoscorpius* and *Mazonia* of Messrs. Meek and Worthen. *Transactions of the Royal Society of Edinburgh*, 30: 397–412.
- Peach, R. N. 1888. On a new eurypterid from the Upper Coal-measures of Radstock, Somersetshire. *Proceedings of the Royal Physical Society, Edinburgh*, 9: 438–445.
- Peckham, G. W. & Peckham, E. G. 1892. Ant-like spiders of the Family Attidae. *Occasional Papers of the Natural History Society of Wisconsin*, 2(1): 1-83.

- Peckham, G. W. & Wheeler, W. H. 1889. Spiders of the subfamily Lyssomanae. *Transactions of the Wisconsin Academy of Science, Arts and Letters*, 7: 222–256.
- Penney, D. 2000. Miocene spiders in Dominican amber (Oonopidae, Mysmenidae). *Palaeontology*, 43: 343–357.
- Penney, D. 2001. Advances in the taxonomy of spiders in Miocene amber from the Dominican Republic (Arthropoda: Araneae). *Palaeontology*, 44: 987–1009.
- Penney, D. 2002. Spiders in Upper Cretaceous amber from New Jersey (Arthropoda: Araneae). *Palaeontology*, 45: 709–724.
- Penney, D. 2003a. *Afrarchaea grimaldii*, a new species of Archaeidae (Araneae) in Cretaceous Burmese amber. *The Journal of Arachnology*, 31: 122–130.
- Penney, D. 2003b. A new deinopid spider from Cretaceous Lebanese amber. *Acta Palaeontologica Polonica*, 48: 569–574.
- Penney, D. 2004a. New spiders in Upper Cretaceous amber from New Jersey in the American Museum of Natural History (Arthropoda: Araneae). *Palaeontology*, 47: 367–375.
- Penney, D. 2004b. Cretaceous Canadian amber spider and the palpimanoidean nature of lagonomegopids. *Acta Palaeontologica Polonica*, 49: 579–584.
- Penney, D. 2004c. A new genus and species of Pisauridae (Araneae) in Cretaceous Burmese amber. *Journal of Systematic Palaeontology*, 2: 141–145.
- Penney, D. 2005a. First fossil Filistatidae: a new species of *Misionella* in Miocene amber from the Dominican Republic. *The Journal of Arachnology*, 33: 93–100.
- Penney, D. 2005b. The fossil spider family Lagonomegopidae in Cretaceous ambers with descriptions of a new genus and species from Myanmar. *The Journal of Arachnology*, 33: 439–444.
- Penney, D. 2005c. First Caribbean *Floricomus* (Araneae: Linyphiidae), a new fossil species in Miocene Dominican Republic amber. A new synonymy for the extant North American fauna. *Geologica Acta*, 3: 59–64.
- Penney, D. 2005d. An annotated systematic catalogue, including synonymies and transfers, of Miocene Dominican Republic amber spiders described up until 2005. *Revista Ibérica de Aracnología*, 12: 25–52.
- Penney, D. 2006a. Fossil oonopid spiders in Cretaceous ambers from Canada and Myanmar. *Palaeontology*, 49: 229–235.
- Penney, D. 2006b. The oldest lagonomegopid spider, a new species in Lower Cretaceous amber from Álava, Spain. *Geologica Acta*, 4: 377–382.
- Penney, D. 2007a. The oldest fossil pholcid and selenopid spiders (Araneae) in lowermost Eocene amber from the Paris Basin France. *The Journal of Arachnology*, 34: 592–598.
- Penney, D. 2007b. A new fossil oonopid spider in lowermost Eocene amber from the Paris Basin, with comments on the fossil spider assemblage. *African Invertebrates*, 48: 71–75.
- Penney, D. 2009. A new spider family record for Hispaniola – a new species of *Plectreurus* (Araneae: Plectreuridae) in Miocene Dominican amber. *Zootaxa*, 2144: 65–68.

- Penney, D. 2011. Grandoculidae: a new fossil spider family from the Upper Cretaceous of Canada. *Bulletin of the British arachnological Society*, 15: 179–180.
- Penney, D. & Ortuño, V. N. 2006. Oldest true orb-weaving spider (Araneae: Araneidae). *Biology Letters*, 2: 447–450.
- Penney, D. & Selden, P. A. 2002. The oldest linyphiid spider in Lower Cretaceous Lebanese amber /Araneae, Linyphiidae, Linyphiinae). *The Journal of Arachnology*, 30: 487–493.
- Penney, D. & Selden, P. A. 2006. First fossil Huttoniidae (Arthropoda: Chelicerata: Araneae) in late Cretaceous Canadian amber. *Cretaceous Research*, 27: 442–446.
- Penney, D., Dierick, M., Cnudde, V., Masschaele, B., Vlassenbroeck, J., Hoorebeke, L. van & Jacobs, P. 2007. First fossil Micropholcommatidae (Araneae), imaged in Eocene Paris amber using X-Ray Computed Tomography. *Zootaxa*, 1623: 47–53.
- Penney, D., Green, D. I., Tichner, S. B., Titchner, B. G., Brown, T. A., Preziosi, R. F. 2012c. An unusual palaeobiocoenosis of subfossil spiders in Colombian copal. *Bulletin of the British Arachnological Society*, 15: 241–244.
- Penney, D., McNeil, A., Green D. I., Bradley, R., Marusik, Y. M., Withers, P. J. & Preziosi, R. F. 2011. A new species of anapid spider (Araneae: Araneoidea, Anapidae) in Eocene Baltic amber, imaged using phase contrast X-ray computed micro-tomography. *Zootaxa*, 2742: 60–66.
- Penney, D., McNeil, A., Green D. I., Bradley, R., Withers, P. J. & Preziosi, R. F. 2012a. The oldest fossil pirate spider (Araneae: Mimetidae), in uppermost Eocene Indian amber, imaged using X-ray computed tomography. *Bulletin of the British Arachnological Society*, 15: 299–302.
- Penney, D., Green D. I., McNeil, A., Bradley, R., Marusik, Y. M., Withers, P. J. & Preziosi, R. F. 2012b. A new species of *Craspedisia* (Araneae: Theridiidae) in Miocene Dominican amber, imaged using X-ray computed tomography. *Paleontological Journal* 46: 583–588. [Translation of Russian original]
- Pérez, d’A.V. 1988. Un oribatido del Eoceno (Terciario). Primar acaro fosil de Chile (Arachnida: Acari: Oribatida). *Revista Chilena de Entomologia*, 16: 23–24.
- Pérez González, A. & Kury A. 2007. Kimulidae. In Pinto da Rocha, R., Machado, G. & Giribet, G. (eds). *Harvestmen. The Biology of Opiliones*. Harvard University Press, Cambridge MA, pp. 207–209.
- Perkovsky, E. E., Rasnitsyn, A. P., Vlaskin, A. P., Taraschuk, M. V. 2007. A comparative analysis of the Baltic and Rovno amber arthropod faunas: representative samples. *African Invertebrates*, 48:229–245
- Perry, M. L. 1995. Preliminary description of a new fossil scorpion from the middle Eocene Green River Formation, Rio Blanco County, Colorado. In Dayvault, R. D. & Averett, W. R. (eds). *The Green River Formation in Piceance Creek and Estern Unita Basins Field Trip*. Grand Junction Geological Society, Grand Junction Colorado, pp. 131–133.

- Peters, W. 1861. (Ueber eine neue Eintheilung der Skorpione und ueber die von ihm in Mossambique gesammelten Arten von Skorpionen). *Monatsberichte der Königlich Preussischen Akademie der Wissenschaft zu Berlin*, 1861: 507–516.
- Petrunkevitch, A. I. 1913. A monograph of the terrestrial Palaeozoic Arachnida of North America. *Transactions of the Connecticut Academy of Arts and Sciences*, 18: 1–137.
- Petrunkevitch, A. I. 1922. Tertiary spiders and opilionids of North America. *Transactions of the Connecticut Academy of Arts and Sciences*, 25: 211–279.
- Petrunkevitch, A. I. 1923. On families of spiders. *Annals of the New York Academy of Science*, 29: 145–180.
- Petrunkevitch, A. I. 1928. Systema Araneorum. *Transactions of the Connecticut Academy of Arts and Sciences*, 29: 1–270.
- Petrunkevitch, A. I. 1942. A study of amber spiders. *Transactions of the Connecticut Academy of Arts and Sciences*, 34: 119–464.
- Petrunkevitch, A. I. 1945a. Palaeozoic Arachnida. An inquiry into their evolutionary trends. *Scientific Papers, Illinois State Museum*, 3(2): 1–76.
- Petrunkevitch, A. I. 1945b. *Calcitro fisheri*. A new fossil arachnid. *American Journal of Science*, 243: 320–329.
- Petrunkevitch, A. I. 1946. Fossil spiders in the collection of the American Museum of Natural History. *American Museum Novitates*, 1328: 1–36.
- Petrunkevitch, A. I. 1949. A study of Palaeozoic Arachnida. *Transactions of the Connecticut Academy of Arts and Sciences*, 37: 69–315.
- Petrunkevitch, A. I. 1950. Baltic amber spiders in the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology*, 103: 257–337.
- Petrunkevitch, A. I. 1953. Palaeozoic and Mesozoic Arachnida of Europe. *Memoirs of the Geological Society of America*, 53: 1–128.
- Petrunkevitch, A. I. 1955a. Arachnida. 42–162. In Moore, R. C. (ed.) *Treatise on invertebrate paleontology, Part P, Arthropoda 2*. Geological Society of America, Boulder, and University of Kansas Press, Lawrence, xvii + 181 pp.
- Petrunkevitch, A. I. 1955b. *Trigonotarbus arnoldi*, a new species of fossil arachnid from Southern France. *Journal of Paleontology*, 29: 475–477.
- Petrunkevitch, A. I. 1958. Amber spiders in European collections. *Transactions of the Connecticut Academy of Arts and Sciences*, 41: 97–400.
- Petrunkevitch, A. I. 1963. Chiapas amber spiders. *University of California Publications in Entomology*, 31: 1–40.
- Petrunkevitch, A. I. 1971. Chiapas amber spiders, II. *University of California Publications in Entomology*, 63: 1–44.
- Pickett, J. W. 1984. A new freshwater limuloid from the middle Triassic of New South Wales. *Palaeontology*, 27: 609–621.

- Pickett, J. W. 1993. A Late Devonian xiphosuran from near Parkes, New South Wales. *Memoirs of the Association of Australian Palaeontologists*, 15: 279–287.
- Pickford, M. 2000. Fossil spider's webs from the Namib Desert and the antiquity of *Seothyra* (Araneae, Eresidae). *Annales de Paléontologie*, 86: 147–155.
- Pictet, F. J. 1846. *Traite élémentaire de paléontologie. Vol. 4*. Paris, 458 pp.
- Pierce, W. D. 1945. A fossil whiptail scorpion from Cabrillo Beach. *Bulletin of the Southern California Academy of Sciences*, 44: 7–8.
- Pierce, W. D. 1950. Fossil arthropods from onyx-marble. *Bulletin of the Southern Californian Academy of Sciences*, 49: 101–104.
- Pierce, W. D. 1951. Fossil arthropods from onyx-marble. *Bulletin of the Southern Californian Academy of Sciences*, 50: 34–49.
- Pinto, I. D. & Hünicken, M. A. 1980. *Gondwanarachne* a new genus of the order Trigonotarbida (Arachnida) from Argentina. *Boletín de la Academia Nacional de Ciencias Córdoba*, 53: 307–315.
- Pirozhnikov, L. P. 1957. [Remains of Gigantostroma from the the series of Matakara (Devonian of North Minusinsk Depression).] *Annuaire de la Société paléontologique de Russie*, 16: 207–213. [in Russian]
- Platnick, N. I. 1977. The hypochiloid spiders: a cladistic analysis, with notes on the Atypoidea (Arachnida, Araneae). *American Museum Novitates*, 2627, 1–23.
- Platnick, N. I. 1989. *Advances in Spider Taxonomy 1981-1987: A Supplement to Brignoli's A Catalogue of the Araneae described between 1940 and 1981*. Manchester University Press, 673 pp.
- Pocock, R. I. 1892. *Liphistius* and its bearing upon the classification of spiders. *Annals and Magazine of Natural History, series 6*, 10: 306–314.
- Pocock, R. I. 1893. Notes on the classification of scorpions, followed by some observations on synonymy, with descriptions of new genera and species. *Annals and Magazine of Natural History, series 6*, 12: 303–330.
- Pocock, R. I. 1895. Description of two new spiders obtained by Messrs J. J. Quelch and F. MacConnel on the summit of Mount Roraima, in Demerara; with a note upon the systematic position of the genus *Desis*. *Annals and Magazine of Natural History, series 6*, 16: 139–143.
- Pocock, R. I. 1897. On the genera and species of tropical African Arachnida of the order Solifugae, with notes upon the taxonomy and habits of the group. *Annals and Magazine of Natural History, series 6*, 20: 249–272.
- Pocock, R. I. 1898. The Arachnida from the Province of Natal, South Africa, contained in the collection of the British Museum. *Annals and Magazine of Natural History, series 7*, 2: 197–226.
- Pocock, R. I. 1901. The Scottish Silurian scorpions. *Quarterly Journal of Microscopical Science*, (2) 44: 291–311.
- Pocock, R. I. 1902. *Eophrynus* and allied Carboniferous Arachnida. *Geological Magazine, Decade 4*, 9: 439–448, 487–493.
- Pocock, R. I. 1903a. A new Carboniferous arachnid. *Geological Magazine, Decade 4*, 10: 247–251.

- Pocock, R. I. 1903b. Further remarks upon the Carboniferous arachnid *Anthracosiro*, with the description of a second species of the genus. *Geological Magazine, Decade 4*, 10: 405–408.
- Pocock, R. I. 1903c. On the geographical distribution of spiders of the order Mygalomorphae. *Proceedings of the Zoological Society of London*, 1903: 340–368.
- Pocock, R. I. 1911. A monograph of the terrestrial Carboniferous Arachnida of Great Britain. *Monographs of the Palaeontographical Society*, 64: 1–84.
- Pohlman, J. 1882. Additional Notes on the Fauna of the Water-Lime Group near Buffalo. *Bulletin of the Buffalo Society of Natural Sciences*, 4(2): 41–47.
- Poinar Jr., J. O. 1985. Fossil evidence of insect parasitism by mites. *International Journal of Acarology*, 11: 37–38.
- Poinar Jr., G.O. 1988. Hair in Dominican amber: evidence for Tertiary land mammals in the Antilles. *Experientia*, 44: 88–89.
- Poinar Jr., G. O. 1995. First fossil soft tick, *Ornithodoros antiquus* n. sp. (Acari: Argasidae) in Dominican amber with evidence of their mammalian host. *Experimentia Basel*, 51: 584–587.
- Poinar Jr., G. [O.] 2008. *Palaeosiro burmanicum* n. gen., n. sp., a fossil Cyphophthalmi (Arachnida: Opiliones: Sironidae) in Early Cretaceous Burmese amber. In Makarov, S. E. & Dimitriević, R. N. (eds) *Advances in Arachnology and Developmental Biology. Papers dedicated to Prof. Dr. Božidar Čurčić*. Inst. Zool., Belgrade; BAS, Sofia; Fac. Life Sci., Vienna; SASA, Belgrade & UNESCO MAB Serbia. Vienna — Belgrade — Sofia, Monographs, 12: 267–274 .
- Poinar Jr., G. O. & Brown, A. E. 2003. A new genus of hard ticks in Cretaceous Burmese amber (Acari: Ixodida: Ixodidae). *Systematic Parasitology*, 54: 199–205.
- Poinar Jr., G. O. & Brown, A. E. 2004. A new whip spider (Arachnida: Amblypygi), *Phrynus mexicana*, is described from Mexican amber. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 3: 1881–1885.
- Poinar Jr., G. O. & Buckley, R. 2008. *Compluriscutula vetulum* (Acari: Ixodida: Ixodidae), a new genus and species of hard tick from Lower Cretaceous Burmese amber. *Proceedings of the Entomological Society of Washington*, 110: 445–450.
- Poinar Jr., G. O. & Buckley, R. 2012. Predatory behaviour of the social orb-weaver spider, *Geratonephila burmanica* n. gen., n. sp. (Araneae: Nephilidae) with its wasp prey, *Cascoscelio incassus* n. gen., n. sp. (Hymenoptera: Platygasteridae) in Early Cretaceous Burmese amber. *Historical Biology*, 24: 519–525.
- Poinar Jr., G. O. & Santiago-Blay, J. A. 1989. A fossil solpugid, *Haplodontus proterus*, new genus, new species (Arachnida: Solpugida) from Dominican amber. *Journal of the New York Entomological Society*, 97: 125–132.
- Ponomarenko, A. G. 1985. King crabs and eurypterids from the Permian and Mesozoic of the USSR. *Paleontological Journal*, 19: 100–104. [Translation of *Paleontologiceskij Žurnal*, 1985: 115–117.]

- Poschmann, M. 2009. Ein fossiler Skorpion aus der Oberkarbon (Westfalium C) des Saar-Nahe-Beckens (SW Deutschland). *Mitteilungen der Pollichia*, 94: 5–10.
- Poschmann, M. & Dunlop, J. A. 2006. A new sea spider (Arthropoda: Pycnogonida) with a flagelliform telson from the Lower Devonian Hunsrück Slate, Germany. *Palaeontology*, 49: 983–989.
- Poschmann, M. & Dunlop, J. A. 2010. Trigonotarbid arachnids from the Lower Devonian (Lower Emsian) of Alken an der Mosel (Rhineland-Palatinate, SW Germany). *Paläontologische Zeitschrift*, 84: 467–484.
- Poschmann, M. & Dunlop, J. A. 2011. Trigonotarbid arachnids from the Lower Devonian (Siegenian) of Bürdenbach (Lahrbach Valley, Westerwald area, Rhenish Slate Mountains, Germany). *Paläontologische Zeitschrift*, 85: 433–447.
- Poschmann, M. & Dunlop, J. A. 2012. Reassessing *Devonotarbus*, a phalangiotarbid arachnid from the Lower Devonian (Siegenian and Emsian) of the Rheinisches Schiefergebirge (SW Germany). *Paläontologisches Zeitschrift*, 86: 377–387.
- Poschmann, M. & Tetlie, O. E. 2004. On the Emsian (Early Devonian) arthropods of the Rhenish Slate Mountains: 4. The eurypterids *Alkenopterus* and *Vinetopterus* n. gen. (Arthropoda: Chelicerata). *Senckenbergiana lethaea*, 84: 173–193.
- Poschmann, M., Anderson, L. I. & Dunlop, J. A. 2005. Chelicerate arthropods, including the oldest phalangiotarbid arachnid, from the Early Devonian (Siegenian) of the Rhenish Massif, Germany. *Journal of Paleontology*, 79: 110–124.
- Poschmann, M., Dunlop, J. A., Kamenz, C. & Scholtz, G. 2008. The Lower Devonian scorpion *Waeringoscorpio* and the respiratory nature of its filamentous structures, with a description of a new species from the Westerwald area, Germany. *Paläontologische Zeitschrift*, 82: 418–436.
- Prach, F. K. 1860. Život Pavouků pravých či předuouých (Araneae). *Živa*, 8: 80–93.
- Presl, J. S. 1822. Additamenta ad faunam protogaeam, sistens descriptiones aliquot animalium in succino inclusorum. In Presl, J. S. & Presl, C. B. (eds). *Deliciae Pragenses Historiam Naturalem Spectantes. Tome I. Calvae, Pragae*, viii + 244 pp.
- Prestwich, J. 1840. Memoir on the geology of Coalbrook Dale. *Transactions of the Geological Society of London* 5: 413–495.
- Příbyl, A. 1952. On the genus *Adelophthalmus* Jordan and Meyer, 1854 (Euryperida) and its representatives in the Upper Carboniferous of Czechoslovakia. *Bulletin International de l'Académie tchéquie des Sciences*, 53: 63–70.
- Příbyl, A. 1958. Some new Carboniferous arachnids from the Ostrava-Karviná coal district. *Časopis pro Mineralogii a Geologii*, 3: 425–434.
- Příbyl, A. 1967. *Moravurus* gen.n. eine neue Xiphosurida Gattung aus dem mährisch-schlesischen Oberkarbon. *Časopis pro Mineralogii a Geologii*, 12: 457–460.

- Pritchard A. E. 1956. A new superfamily of trombidiform mites with the description of a new family, genus and species (Acarina: Iolinioidea: Iolinidae: *Iolina nana*). *Annals of the Entomological Society of America*, 49: 204–206.
- Protescu, O. 1937. Etude géologique et paléobiologique de l'ambre roumain. *Bulletin de la Société române Géologique*, 3: 65–110.
- Prószyński, J. & Żabka, M. 1980. Remarks on Oligocene amber spiders of the family Salticidae. *Acta Palaeontologica Polonica*, 25: 213–223.
- Pruvost, P. 1912. Note sur les Araignées du terrain houiller du Nord de la France. *Annales de la Société Géologique du Nord*, 41: 85–100.
- Pruvost, P. 1919. *Introduction à l'étude du terrain houiller du Nord et du Pas-de-Calais: La faune continentale du terrain houiller de la France*. pp. 339–364. *Classe des Arachnides*. Thèse Université de Lille, Lille.
- Pruvost, P. 1922. Les arachnides fossiles du Houiller de Belgique. *Annales de la Société Scientifique de Bruxelles*, 41: 349–355.
- Pruvost, P. 1926. Description de deux fossiles du terrain houiller de Noeux (*Anthracosiro corsini*, nov. sp. et *Fayolia stertzeli* Weiss). *Annales de la Société Géologique du Nord*, 51: 144–149.
- Pruvost, P. 1930. La Faune continentale du terrain houiller de la Belgique. Arachnides. *Mémoires du Musée royal d'Histoire naturelle de Belgique*, 44: 206–217.
- Pruvost, P. 1939. *Euypteris (Anthraconectes) corneti* du Westphalien A du couchant de Mons. *Annales de la Société Scientifique de Bruxelles*, 59: 56–59.
- Qin, T. K. & Halliday, R. B. 1997. Eriorhynchidae, a new family of Prostigmata (Acarina), with a cladistic analysis of eupodoid species of Australia and New Zealand. *Systematic Entomology*, 22: 151–171.
- Quintero Jr., D. 1996. Revision de la clasificación de Amblypygidos pulvanados: creación de subordenes, una nueva familia y un nuevo género con tres nuevas especies (Arachnida: Amblypygi). 203–212. In Eberhardt, W. G., Lubin, Y. D. & Robinson, B. C. (eds). *Proceedings of the Ninth International Congress of Arachnology, Panama 1983*. Smithsonian Institution Press, Washington, DC, xx pp.
- Racheboeuf, P. R. 1992. *Valloisella lievinensis* n. g. n. sp.: nouveau Xiphosure carbonifère du nord de la France. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1992(6): 336–342.
- Racheboeuf, P. R., Vannier, J. & Anderson, L. I. 2002. A new three-dimensionally preserved xiphosuran chelicerate from the Montceau-les-Mines Lagerstätte (Carboniferous, France). *Palaeontology*, 45: 125–147.
- Ramírez, M. J. & Grismado, C. J. 1997. A review of the spider family Filistatidae in Argentina (Arachnida: Araneae), with a cladistic reanalysis of filistatid genera. *Entomologica Scandinavica*, 28: 319–349.
- Ramsay, G.W. 1960. Sub-fossil mites from the Hutt Valley. *Transactions of the Royal Society of New Zealand*, 88: 575–576.

- Raven, R. J. 1985. The spider infraorder Mygalomorphae (Araneae): cladistics and systematics. *Bulletin of the American Museum of Natural History*, 182: 1–180.
- Raymond, P. E. 1944. Late Paleozoic xiphosurans. *Bulletin of the Museum of Comparative Zoology*, 94: 475–508.
- Redell, J. R. & Cokendolpher, J. C. 1995. Catalogue, bibliography and generic revision of the order Schizomida (Arachnida). *Texas Memorial Museum, Speleological Monographs*, 4: 1–170.
- Reeside, J. B. & Harris, D. V. 1952. A Cretaceous horseshoe crab from Colorado. *Journal of the Washington Academy of Science*, 42: 174–178.
- Reiskind, J. 1986. A new *Lyssomanes* from the Dominican amber and the possible use of insular fossils in building phylogenies. 423. In Barrientos, J. A. (ed.) *Actas X Congreso Internacional de Aracnología, Jaca. España, Volume 1*. Barcelona.
- Reiskind, J. 1989. The potential use of amber fossils in the study of the biogeography of spiders in the Caribbean with the description of a new species of *Lyssomanes* from Dominican amber (Araneae: Salticidae). 217–228. In Woods, C. A. (ed.) *Biogeography of the West Indies, past, present and future*. Sandhill Crane Press, Gainesville, Florida.
- Remy, W. & Remy, R. 1959. Arthropodenfunde im Stefan der Halleschen Mulde. *Monograph-Bericht der Deutschen Akademie Wissenschaft Berlin*, 1: 299–312.
- Reuss, A. E. 1855. Palaeontologische Miscellen. III. Über eine neue Krusterspecies aus der Böhmischen Steinkohlenformation. *Denkschrift der königlich-kaiserlichen Akademie der Wissenschaft in Wien*, 10: 81–83.
- Richter, R. & Richter, E. 1929. *Weinbergina opitzi* n. g., n. sp., ein Schwerträger (Merost. Xiphos.) aus dem Devon (Rheinland). *Senckenbergiana*, 11: 193–209.
- Ribera, C. 2003. El arácanido del Plesiotoceno inferior de Incaral V (Girona, NE de la Península Ibérica). *Paleontologia i Evolució*, 34: 51–53.
- Riek, E. F. 1955. A new xiphosuran from the Triassic sediments at Brookvale, New South Wales. *Records of the Australian Museum*, 23: 281–282.
- Riek, E. F. & Gill, E. D. 1971. A new xiphosuran genus from Lower Cretaceous Freshwater sediments at Koonwarra, Victoria, Australia. *Palaeontology*, 14: 206–210.
- Riquelme, F. & Hill, D. E. 2013. Insights into amber salticids from the Neogene of Middle America, with the first report of Marpissinae (Araneae: Salticidae) from the Chiapas amber. *Peckhamia*, 106.1: 1–5.
- Risso, A. 1826. Animaux articulés: description de quelques Myriapodes, Scorpionides, Arachnides et Acarides, habitant les Alpes Maritimes. In Risso, A. (ed.). *Histoire Naturelle des Principales Productions de l'Europe Méridionale et Principalement de Celles des Environs de Nice et des Alpes Maritimes*. Levrault, Paris, xx pp.

- Ritchie, A. 1968. *Lanarkopterus dolichoshelus* (Størmer) gen. nov., a mixopterid eurypterid from the Upper Silurian of the Lesmahagow and Hagshaw Hills inliers, Scotland. *Scottish Journal of Geology*, 4: 317–338.
- Robineau-Desvoidy, J. B. 1828. *Recherches sur l'organisation vertébrale des Crustacés, Arachnides et Insectes*. Composé Jeune, Paris, 228 pp.
- Roemer, F. 1866. *Protolycosa anthracophila*, eine fossile Spinne aus dem Steinkohlengebirge Oberschlesiens. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*: 136–143.
- Roemer, F. 1878. Auffindung und Vorlegung eines neuen Gliderthieres in dem Steinkohlengebiete der Ferdinandsgrube bei Glatz. *Jahresbericht der Schlesischen Gesellschaft für Vaterländische-Kultur.*, pp. 54–55.
- Roewer, C. F. 1912. Die Familien der Assamiden und Phalangodiden der Opiliones-Laniatores. (=Assamiden, Dampetriden, Phalangodiden, Epedaniden, Biantiden, Zalmoxiden, Samoiden, Palpipediden anderer Autoren.) *Archiv für Naturgeschichte* 78A (3): 1–242.
- Roewer, C.-F. 1913. Die Familie der Gonyleptiden der Opiliones-Laniatores. *Archiv für Naturgeschichte*, 79A (4, 5): 1–256, 257–473.
- Roewer, C.-F. 1923. *Die Weberknechte der Erde. Systematische Bearbeitung der bisher bekannten Opiliones*. Gustav Fischer, Jena, 1116 pp.
- Roewer, C.-F. 1933. Solifugae, Palpigradi. 161–480. In Bronn, H. G. (ed.). *Klassen und Ordnung des Tierreichs*. 5: *Arthropoda IV: Arachnoidea, vol. 5(IV) (4) (2–3)*. Akademische Verlagsgesellschaft M.B.H, Leipzig.
- Roewer, C.-F. 1934. Solifugae, Palpigradi. 481–723. In Bronn, H. G. (ed.). *Klassen und Ordnung des Tierreichs*. 5: *Arthropoda IV: Arachnoidea, vol. 5(IV) (4) (4–5)*. Akademische Verlagsgesellschaft M.B.H, Leipzig.
- Roewer, C.-F. 1935. Zwei myrmecophile Spinnen-Arten Brasiliens. *Veröffentlichungen aus dem Deutschen Kolonial- und Übersee-Museum in Bremen*, 1: 193–197.
- Roewer, C.-F. 1939. Opilioniden im Bernstein. *Palaeobiologica*, 7(1): 1–4.
- Roewer, C.-F. 1942. *Katalog der Araneae von 1758 bis 1940. 1. Band*. Kommissions-Verlag von „NATURA“: 1040 pp.
- Roewer, C.-F. 1943. Über Gonyleptiden. Weitere Weberknechte (Arachn., Opil.) XI. *Senckenbergiana*, 26: 12–68.
- Roewer, C.-F. 1951. Über Nemastomatiden. Weitere Weberknechte XVI. *Senckenbergiana*, 32: 95–153.
- Roivainen, H. 1953. Subfamilies of European eriophyid mites. *Annales entomologici Fennici*, 19: 83–87.
- Romero, A. & Via Boada, L. 1977. *Tarracolimulus rieki*, nov. gen., nov. sp., nuevo limulido del Triásico de Montral-Alcover (Tarragona). *Cuadernos de Geología Ibérica*, 4: 239–246.
- Ross, A. J. & Vannier, J. 2002. Crustacea (excluding Ostracoda) and Chelicerata of the Purbeck Limestone Group, southern England: a review. *Special Papers in Palaeontology*, 68: 71–82.

- Rößler, R. & Schneider, J. 1997. Eine bemerkenswerte Paläobiocoenose im Unterkarbon Mitteleuropas – Fossilführung und Paläoenvironment der Hainichen-Subgruppe (Erzgebirge-Becken). *Veröffentlichungen des Museums für Naturkunde Chemnitz*, 20: 5–44.
- Rowland, J. M. 1975. A partial revision of Schizomida (Arachnida) with descriptions of new species, genus, and family. *Occasional Papers of the Museum, Texas Tech University*, 31: 1–21.
- Rowland, J. M. & Sissom, W. D. 1980. Report on a fossil palpigrade from the Tertiary of Arizona, and a review of the morphology and systematics of the order (Arachnida: Palpigradida). *The Journal of Arachnology*, 8: 69–86.
- Rudkin, D. M., Young, G. A. & Nowlan, G. S. 2008. The oldest horseshoe crab: a new xiphosurid from late Ordovician Konservat-Lagerstätten deposits, Manitoba, Canada. *Palaeontology*, 51: 1–9.
- Rudkin, D. M., Cuggy, M. B., Young, G. A. & Thompson, D. P. 2013. An Ordovician pycnogonid (sea spider) with serially subdivided 'head' region. *Journal of Paleontology*, 87: 395–405.
- Ruedemann, R. 1916. Account of some new or little known species of fossils, mostly from the Palaeozoic rocks of New York. *New York State Museum Bulletin*, 189: 7–112.
- Ruedemann, R. 1921. A recurrent Pittsford (Salina) fauna. *New York State Museum Bulletin*, 219–20: 205–215.
- Ruedemann, R. 1926. The Utica and Lorraine Formations of New York, Part 2, Systematic Paleontology, no. 2, Mollusks, Crustacea and Eurypterids. *New York State Museum Bulletin*, 189: 98–112.
- Ruedemann, R. 1942. Some new eurypterids from New York. *New York State Museum Bulletin*, 327: 24–29.
- Russell, L. S. 1953. A new species of eurypterid from the Devonian of Gaspé. *Annual Report of the National Museum for the Fiscal Year 1952–1953, Bulletin*, 132: 83–91.
- Ryke, P. A. J. 1962. The subfamily Rhodacarinae with notes on a new subfamily Ologamasinae (Acarina: Rhodacaridae). *Entomologische Berichte Amsterdam*, 22: 155–162.
- Salter, J. W. 1856. On some new Crustacea from the uppermost Silurian Rocks. *Quarterly Journal of the Geological Society of London*, 12: 26–34.
- Sanchez, J. P., Nava, S., Lareschi, M., Ortiz, P. E. & Guglielmone, A. A. 2010. Finding of an ixodid tick inside a late Holocene owl pellet from northwestern Argentina. *Journal of Parasitology*, 96: 820–822.
- Santiago-Blay, J. A. & Poinar Jr., G. O. 1988. A fossil scorpion *Tityus geratus* new species (Scorpiones: Buthidae) from Dominican amber. *Historical Biology*, 1: 345–354.
- Santiago-Blay, J. A., Fet, V., Soleglad, M. E. & Anderson, S. R. 2004. A new genus and subfamily of scorpions from Lower Cretaceous Burmese amber (Scorpiones: Chaerilidae). *Revista Ibérica de Aracnología*, 9: 3–14.
- Sarle, C. J. 1903. A new eurypterid fauna from the base of the Salina in western New York. *New York State Museum Bulletin*, 69: 1080–1108.
- Sars, G. O. 1891. Pycnogonidea. *Norwegian North-Atlantic Expedition, 1876–1878*, 6 (Zool. 20): 1–163.

- Saupe, E. E. & Selden, P. A. 2009. First fossil Mecysmaucheniidae (Arachnida, Chelicerata, Araneae), from Lower Cretaceous (uppermost Albian) amber of Charente-Maritime, France. *Geodiversitas*, 31: 49–60.
- Saupe, E. E., Selden, P. A. & Penney, D. 2010. First fossil *Molinaranea* Mello-Leitão, 1940 (Araneae: Araneidae), from middle Miocene Dominican amber, with a phylogenetic and palaeobiogeographical analysis of the genus. *Zoological Journal of the Linnean Society*, 158: 711–725.
- Saupe, E. E., Pérez-de la Fuente, R., Selden, P. A., Delclòs, X., Tafforeau, P. & Soriano, C. 2012. New *Orchestina* Simon, 1882 (Araneae: Oonopidae) from Cretaceous ambers of Spain and France: First spider described using phase-contrast x-ray synchrotron microtomography. *Palaeontology*, 55: 127–143.
- Savage, T. E. 1916. Alexandrian rocks of northeastern Illinois and eastern Wisconsin. *Bulletin of the Geological Society of America*, 27: 305–324.
- Sayre, R. M., Smiley, R. L. & Walter, D. E. 1992. Report of a teneriffiid mite (Acari) in Baltic amber and notes on recent discoveries. *International Journal of Acarology*, 18: 303–305.
- Scharf, W. 1924. Beitrag zur Geologie des Steinkohlengebietes im Südharz. *Jahrbuch des Halleschen Verbands für die Erforschung der Mitteldeutschen Bodenschätze und ihrer Verwaltung*, 4: 404–437.
- Schawaller, W. 1978. Neue Pseudoskorpione aus dem Baltischen Bernstein der Stuttgarter Bernsteinsammlung (Arachnida: Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 42: 1–21.
- Schawaller, W. 1979a. Erstnachweis eines Skorpions in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Scorpionida). *Stuttgarter Beiträge zur Naturkunde (B)*, 45: 1–15.
- Schawaller, W. 1979b. Erstnachweis der Ordnung Geisselspinnen in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Amblypygi). *Stuttgarter Beiträge zur Naturkunde (B)*, 50: 1–12.
- Schawaller, W. 1980a. Fossile Chthoniidae in Dominikanischem Bernstein, mit phylogenetischen Anmerkungen (Stuttgarter Bernsteinsammlung: Arachnida, Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 63: 1–19.
- Schawaller, W. 1980b. Erstnachweis tertiärer Pseudoskorpione (Chernetidae) in Dominikanischen Bernstein. *Stuttgarter Beitrag zur Naturkunde (B)*, 57: 1–20.
- Schawaller, W. 1981. Cheiridiidae in Dominikanischem Bernstein, mit Anmerkungen zur morphologischen Variabilität (Stuttgarter Bernsteinsammlung: Arachnida, Pseudoscorpionidea). *Stuttgarter Beiträge zur Naturkunde (B)*, 75: 1–14.
- Schawaller, W. 1982a. Zwei weitere Skorpione in Dominikanischem Bernstein (Stuttgarter Bernsteinsammlung: Arachnida, Scorpionida). *Stuttgarter Beiträge zur Naturkunde (B)*, 82: 1–14.
- Schawaller, W. 1982b. Der erste Pseudoskorpion (Chernetidae) aus Mexikanischem Bernstein. *Stuttgarter Beiträge zur Naturkunde (B)*, 85: 1–9.
- Schawaller, W. 1982c. Spinnen der Familien Tetragnathidae, Uloboridae und Dipluridae in Dominikanischem Bernstein und allgemeine Gesichtspunkte (Arachnida, Araneae). *Stuttgarter Beiträge zur Naturkunde (B)*, 89: 1–19.

- Schawaller, W. 1982d. Zur fossilen Spinnenfauna des Pliozäns von Willershäusen in Norddeutschland (Arachnida, Araneae). *Berichte der Naturhistorischen Gesellschaft zu Hannover*, 125: 89–95.
- Schawaller, W. 1984. The family Selenopidae in Dominican amber (Arachnida: Araneae). *Stuttgarter Beiträge zur Naturkunde (B)*, 103: 1–8.
- Schawaller, W., 1991. The first Mesozoic pseudoscorpion, from Cretaceous Canadian amber. *Palaeontology*, 34: 971–976.
- Schawaller, W. & Ono H. 1979. Fossile Spinnen aus miozänen Sedimenten des Randecker Maars in SW-Deutschland (Arachnida: Araneae). *Jahreshefte der Gesellschaft für Naturkunde in Württemberg*, 134: 131–141.
- Schawaller, W., Shear, W. A. & Bonamo, P. M. 1991. The first Paleozoic pseudoscorpions (Arachnida, Pseudoscorpionida). *American Museum Novitates*, 3009: 1–17.
- Schille, F. 1916. Entomologie aus der Mammut- und Rhinoceros-Zeit Galiziens. *Entomologische Zeitschrift*, 30: 42–43.
- Schimkewitsch, W. 1913. Ein Beitrag zur Klassifikation der Pantopoden. *Zoologischen Anzeiger*, 41: 597–615.
- Schimper, W. P. 1853. Paleontologica alsatica ou fragments paléontologiques des différents terrains stratifiés qui se rencontrent en Alsace. *Mémoires de la Société du Muséum d'Histoire Naturelle de Strasbourg*, 4: 1–10.
- Schmidt, A. R., Jancke, S., Lindquist, E. E., Ragazzi, E., Roghi, G., Nascimbene, P. C., Schmidt, K., Wappler, T. & Grimaldi, D. A. 2012. Arthropods in amber from the Triassic period. *Proceedings of the National Academy of Science, USA*, doi/10.1073/pnas.1208464109.
- Schmidt, A. R., Perrichot, V., Svojtka, M., Anderson, K. B., Belete, K. H., Bussert, R., Dörfelt, H., Jancke, S., Mohr, B., Mohrmann, E., Nascimbene, P. C., Nel, A., Nel, P., Ragazzi, E., Roghi, G., Saupe, E. E., Schmidt, K., Schneider, H., Selden, P. A., Vávra, N. 2010. Cretaceous life captured in amber. *Proceedings of the National Academy of Sciences, USA*: doi/10.1073/pnas.1000948107.
- Schmidt, F. 1883. Nachtrag zur Monographie der Russischen Leperditen II. Die Crustaceenfauna der Euryptereenschichten von Rootziküll auf Oesel. *Miscellanea silurica III. Memoirs of the Academy of Science de St. Petersburg*, 31: 28–85.
- Schram, F. R. 1979. Limulines of the Mississippian Bear Gulch Limestone of Central Montana, USA. *Transactions of the San Diego Society of Natural History*, 19: 67–74.
- Schultka, S. 1991. *Trigonotarbus stoermeri* n. sp. – ein Spinnentier aus den Bensberger Schichten (Ems/Unter-Devon) des Rheinischen Schiefergebirge. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 183: 375–390.
- Schuster, R. 1963. *Thalassozetes riparius* n. gen., n. sp., eine litoralbewohnende Oribatide von bemerkenswerter morphologischer Variabilität (Oribatei-Acari). *Zoologischer Anzeiger*, 171: 391–403.
- Scopoli, J. A. 1763. *Entomologia Carniolica, exhibens Insecta Carniolae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana. Vindobonae*, 1763: 420 pp.

- Scott, A. G. 2003. Sub-fossil spiders from Holocene peat cores. *Journal of Arachnology*, 31: 1–7.
- Scudder, S. H. 1868. Supplement to descriptions of Articulatcs. Description of fossil insects found on Mazon Creek and near Morris, Grundy Co., Ill. *Geological Survey of Illinois*, 3: 566–572.
- Scudder, S. H. 1876. New and interesting insects from the Carboniferous of Cape Breton. *Canadian Naturalist and Quarterly Journal of Science*, 8: 88–90.
- Scudder, S. H. 1878. Additions to the Insect-Fauna of the Tertiary Beds at Quesnel, British Columbia. *Geological Survey of Canada. Report of Progress, 1876–1877*: 457–464.
- Scudder, S. H. 1884. A contribution to our knowledge of Paleozoic Arachnida. *Proceedings of the American Academy of Arts and Sciences*, 20: 13–22.
- Scudder, S. H. 1885. 3. Classe. Arachnoidea. Spinnen. Skorpione. 732–746. In Zittel, K. A. (ed), *Handbuch der Palaeontologie. I. Abtheilung. Palaeozoologie* 2. R. Oldenbourg, München & Leipzig.
- Scudder, S. H. 1890a. The Tertiary Insects of North America. *Report of the United States Geological Survey*, 13: 734 pp.
- Scudder, S. H. 1890b. Illustrations of the Carboniferous Arachnida of North America, of the orders Anthracomarti and Pedipalpi. *Memoirs of the Boston Society of Natural History*, 4: 443–456.
- Scudder, S. H. 1891. Index to the known fossil insects of the world including myriapods and arachnids. *Bulletin of the United States Geological Survey* 71: 1–744.
- Seemann, F. 1906. Beiträge zur Gigantotrakenfauna Böhmens. *Beiträge zur Paläontologie Österreich-Ungarns und des Orients*, 19: 49–57.
- Selden, P. A. 1990. Lower Cretaceous spiders from the Sierra de Montsech, north-east Spain. *Palaeontology*, 33: 257–285.
- Selden, P. A. 1992. Revision of the fossil ricinuleids. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 83: 595–634.
- Selden, P. A. 1996. First fossil mesothele spider from the Carboniferous of France. *Revue suisse de Zoologie*, hors série: 585–596.
- Selden, P. A. 2000. *Palaeothele*, replacement name for the fossil mesothele spider *Eothele* non Rowell. *Bulletin of the British arachnological Society*, 11: 292.
- Selden, P. A. 2001. Eocene spiders from the Isle of Wight with preserved respiratory structures. *Palaeontology*, 44: 695–729.
- Selden, P. A. 2002. First British Mesozoic spider, from Cretaceous amber of the Isle of Wight, southern England. *Palaeontology*, 45: 973–983.
- Selden, P. A. 2010. A theridiosomatid spider from the Early Cretaceous of Russia. *Bulletin of the British arachnological Society*, 15: 69–78.
- Selden, P. A. & Drygant, D. M. 1987. A new xiphosuran from the Silurian of Podolia, Ukraine, USSR. *Palaeontology*, 30: 537–542.

- Selden, P. A. & Gall, J.-C. 1992. A Triassic mygalomorph spider from the northern Vosges, France. *Palaeontology*, 35: 211–235.
- Selden, P.A. & Huang, D.-y. 2010. The oldest haplogyne spider (Araneae: Plectreuridae), from the Middle Jurassic of China. *Naturwissenschaften*, 97: 449–459
- Selden, P. A. & Penney, D. 2003. Lower Cretaceous spiders (Arthropoda: Arachnida: Araneae) from Spain. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 2003: 175–192.
- Selden, P. A. & Penney, D. 2009. A fossil spider (Araneae: Pisauridae) of Eocene age from Horsefly, British Columbia, Canada. *Contributions to Natural History*, 12: 1269–1282.
- Selden, P. A. & Shear, W. A. 1996. The first Mesozoic solifuge (Arachnida), from the Cretaceous of Brazil, and a redescription of the Palaeozoic solifuge. *Palaeontology*, 39: 583–604.
- Selden, P. A. & Siveter, D. J. 1987. The origin of the limuloids. *Lethaia*, 20: 383–392.
- Selden, P. A., Baker, A. S. & Phipps, K. J. 2008. An oribatid mite (Arachnida: Acari) from the Oxford Clay (Jurassic: Upper Callovian) of South Cave Station Quarry, Yorkshire, UK. *Palaeontology*, 51: 623–633.
- Selden, P. A., Casado, F. C. & Mesquita, M. V. 2006. Mygalomorph spiders (Araneae: Dipluridae) from the Lower Cretaceous Crato Lagerstätte, Araripe Basin, north-east Brazil. *Palaeontology*, 49: 817–826.
- Selden, P. A., Huang D.-y., Ren D. 2008. Palpimanoid spiders from the Jurassic of China. *Journal of Arachnology*, 36: 306–321.
- Selden, P. A., Shear, W. A. & Bonamo, P. M. 1991. A spider and other arachnids from the Devonian of New York, and reinterpretations of Devonian Araneae. *Palaeontology*, 34: 241–281.
- Selden, P. A., Shear, W. A. & Sutton, M. D. 2008. Fossil evidence for the origin of spider spinnerets, and a proposed arachnid order. *Proceedings of the National Academy of Sciences of the United States of America*, 105: 20781–20785.
- Selden, P. A., Anderson, J. M., Anderson, H. M. & Fraser, N. C. 1999. Fossil araneomorph spiders from the Triassic of South Africa and Virginia. *Journal of Arachnology*, 27: 401–414.
- Selden, P. A., Nam, K.-s., Kim, S. H. & Kim, H. J. 2012. A fossil spider from the Cretaceous of Korea. *Journal of Palaeontology*, 86: 1–6.
- Sellnick, M. 1918. Die Oribatiden der Bernsteinsammlung der Universität Königsberg I Pr. *Schriften der Physikalisch-Ökonomischen Gesellschaft zu Königsberg (1919)*, 59: 21–42.
- Sellnick, M. 1922. Milben der Sammlung des Deutschen Entomologischen Instituts. I. Oribatidae. *Entomologische Mitteilungen*, 11: 18–20.
- Sellnick, M. 1928. Formenkreis: Hornmilben, Oribatei. In Brohmer, P., Ehrmann, P. & Ulmer, G. (eds). *Die Tierwelt Mitteleuropas*, 3, 4(9): 1–42.
- Sellnick, M. 1931. Milben im Berstein. *Bersteinforschung*, 2: 148–180.
- Semper, M. 1898. Die Gigantostroken des älteren böhmischen Paläozoicum. *Beiträge zur Paläontologie Österreich-Ungarns und des Orients*, 2: 71–88.

- Shear, W. A., 1980. A review of the Cyphophthalmi of the United States and Mexico, with a proposed reclassification of the suborder (Arachnida, Opiliones). *American Museum Novitates*, 2705: 1–34.
- Shear, W. A., 1986. A cladistic analysis of the opilionid superfamily Ischyropsalidoidea, with description of the new family Ceratolasmatidae, the new genus *Acuclavella* and four new species. *American Museum Novitates*, 2844: 1–29.
- Shear, W. A., 1993. The genus *Troglosiro* and the new family Troglosironidae (Opiliones, Cyphophthalmi). *Journal of Arachnology*, 21: 81–90.
- Shear, W. A. 2000. *Gigantocharinus szatmaryi*, a new trigonotarbid arachnid from the Late Devonian of North America (Chelicerata, Arachnida, Trigonotarbida). *Journal of Paleontology*, 74: 25–31.
- Shear, W. A. 2010. New species and records of ortholasmatine harvestmen from México, Honduras, and the western United States (Opiliones, Nemastomatidae, Ortholasmatinae). *ZooKeys*, 52: 9–45.
- Shear, W. A., Selden, P. A., Rolfe, W. D. I., Bonamo, P. M. & Grierson, J. D. 1987. New terrestrial arachnids from the Devonian of Gilboa, New York. *American Museum Novitates*, 2901: 1–74.
- Sharma, P. P. & Giribet, G. 2011. The evolutionary and biogeographic history of the armoured harvestmen – Laniatores phylogeny based on ten molecular markers, with the description of two new families of Opiliones (Arachnida). *Invertebrate Systematics*, 25: 106–142.
- Sharma, P. P., Prieto, C. E. & Giribet, G. 2011. A new family of Laniatores (Arachnida: Opiliones) from the Afrotropics. *Invertebrate Systematics*, 25: 143–154.
- Shpinev, E. S. 2006. A new species of *Adelophthalmus* (Eurypterida) from the Lower Carboniferous of the Krasnoyarsk Region. *Paleontological Journal*, 40: 431–433. [English translation of Russian original]
- Shpinev, E. S. 2012. On some eurypterids (Eurypterida, Chelicerata) from the Devonian of South Siberia. *Paleontological Journal*, 46: 370–377. [English translation of Russian original]
- Shuler, E. W. 1915. A new Ordovician eurypterid. *American Journal of Science*, 4th Series, 39: 551–554.
- Sidorchuk, E. A. & Bertrand, M. 2013. New fossil labidostomatids (Acari: Labidostomatidae) from Eocene amber and presence of an apustulate species in Europe. *Acarologia*, 53: 25–39.
- Sidorchuk, E. A. & Klimov, P. B. 2011. Redescription of the mite *Glaesacarus rhombeus* (Koch & Berendt, 1854) from Baltic amber (Upper Eocene): evidence for female-controlled mating. *Journal of Systematic Palaeontology*, 9: 183–196.
- Sidorchuk, E. A. & Norton, R. A. 2011. The fossil mite family Archaeorchestidae (Acari, Oribatida) I: redescription of *Stieremaeus illibatus* and synonymy of *Strieremaeus* with *Archaeorchestes*. *Zootaxa*, 2993: 34–58.
- Siebold, C. T. E. von. 1850. Ueber *Eriophyes*. *Jahresbericht der Schlesischen Gesellschaft*, 28: 88–89.
- Siegfried, P. 1972. Ein Schwertschwanz (Merostomata, Xiphosurida) aus dem Oberkarbon von Ibbenburen/Westfalen. *Paläontologische Zeitschrift*, 46, 180–186.

- Šilhavý, V. 1973. Two new systematic groups of the gonyleptomorph phalangids from the Antillean-Caribbean Region. Agoristenidae fam. n. and Caribbantinae subfam. n. *Věstník Československé Společnosti Zoologické*, 37: 110–143.
- Šilhavý, V. 1979. New American representatives of the subfamily Samoinae (Opiliones, Phalangodidae, Arach.). *Annotationes zoologicae et botanicae, Bratislava*, 130: 1–27.
- Simon, E. 1864. *Histoire naturelle des Araignées (Aranéides)*. Paris, 540 pp.
- Simon, E. 1874. *Les arachnides de France. Tome 1*. Paris, 272 pp.
- Simon, E. 1876a. *Les Arachnides de France. Tome 3*. Paris, 360 pp.
- Simon, E. 1876b. Etude sur les Arachnides du Congo. *Bulletin de la Société zoologique de France*, 1: 12–15, 216–224.
- Simon, E. 1879a. *Les Arachnides de France VII. Contenant les ordres des Chernetes, Scorpiones et Opiliones*. Paris, 332 pp.
- Simon, E. 1879b. Essai d'une classification des Opiliones Mecostethi. Remarques synonymiques et descriptions d'espèces nouvelles. *Annales de la Société Entomologique de Belgique*, 22: 183–241.
- Simon, E. 1880. Études arachnologiques 12e Mémoire(1). XVII. Descriptions de Genres et Espèces de l'ordre des Scorpiones. *Annales de la Société Entomologique de France*, (5)10: 377–398.
- Simon, E. 1881. *Les Arachnides de France. Tome 5, 1^{re} partie*. Paris, 179 pp.
- Simon, E. 1882. Etudes arachnologiques. 13^e Mémoire. 20. Descriptions d'espèces et de genres nouveaux de la famille des Dysderidae. *Annales de la Société Entomologique de France*, (6) 2: 201–240.
- Simon, E. 1884a. Note synonymique sur les genres *Prodidomus* Hentz et *Miltia* E.S. *Annales de la Société Entomologique de Belgique*, 28: 302.
- Simon, E. 1884b. Note complémentaire sur la famille des Archaeidae. *Annali del Museo Civico di Storia Naturale di Genova*, 20: 373–380.
- Simon, E. 1884c. *Les Arachnides de France. Tome 5, 2^e et 3^e parties*. Paris, pp. 180–808.
- Simon, E. 1884d. Description d'une nouvelle famille de l'ordre des Araneae (Bradystichidae). *Annales de la Société Entomologique de Belgique*, 28: 297–301.
- Simon, E. 1885a. Etudes arachnologiques. 17e Mémoire. XXVI. Arachnides recueillis dans la vallée de Templé et sur le mont Ossa (Thessalie). *Annales de la Société Entomologique de France*, 5: 209–217.
- Simon, E. 1885b. Etude sur les Arachnides recueillis en Tunisie en 1883 et 1884 par MM. A. Letourneux, M. Sédillot et Valéry Mayet, membres de la Mission de l'Exploration scientifique de la Tunisie. *In Exploration scientifique de la Tunisie*, Paris, 55 pp.
- Simon, E. 1885c. Etudes arachnologiques. 18e Mémoire. XXVI. Matériaux pour servir à la fauna des Arachnides du Sénégal. (Suivi d'un appendice intitulé: Descriptions de plusieurs espèces africaines nouvelles). *Annales de la Société Entomologique de France*, 5: 345–396.

- Simon, E. 1887. Espèces et genres nouveaux de la famille des Sparassidae. *Bulletin de la Société zoologique de France*, 12: 466–474.
- Simon, E. 1888. Etudes arachnologiques. 21^e Mémoire. 29. Descriptions d'espèces et de genres nouveaux de l'Amérique centrale et des Antilles. *Annales de la Société Entomologique de France*, (6) 8: 203–216.
- Simon, E. 1889a. Etudes arachnologiques. 21^e Mémoire. 31. Descriptions d'espèces et de genres nouveaux de Madagascar et de Mayotte. *Annales de la Société Entomologique de France*, (6) 8: 223–236.
- Simon, E. 1889b. Arachnides. In Voyage de M. E. Simon au Venezuela (décembre 1887 – avril 1888). 4^e Mémoire. *Annales de la Société Entomologique de France*, (6) 9: 169–220.
- Simon, E. 1890. Etudes arachnologiques. 22^e Mémoire. 34. Etude sur les Arachnides de l'Yemen. *Annales de la Société Entomologique de France*, 10: 77–124.
- Simon, E. 1891a. Observations biologiques sur les Arachnides. I. Araignées sociables. In Voyage de M. E. Simon au Venezuela (Décembre 1887 – avril 1888). 11^e Mémoire. *Annales de la Société Entomologique de France*, 60: 5–14.
- Simon, E. 1891b. On the spiders of the Island of St. Vincent. Part I. *Proceedings of the Zoological Society of London*, 1891: 549–575.
- Simon, E. 1892a. Arachnides. In Raffray, A., Bolivar, I. & Simon, E. (eds) Etude sur les Arthropodes cavernicoles de île Luzon, Voyage de M. E. Simon aux îles Philippines (Mars et avril 1890). *Annales de la Société Entomologique de France*, 61: 35–52.
- Simon, E. 1892b. *Histoire naturelle des Araignées. Volume 1, part 1*. Roret, Paris, pp. 1–254.
- Simon, E. 1893. *Histoire naturelle des Araignées. Volume 1, part 2*. Roret, Paris, pp. 255–488.
- Simon, E. 1894. *Histoire naturelle des Araignées, Volume 1, part 3*. Roret, Paris, pp. 489–760.
- Simon, E. 1895. *Histoire naturelle des Araignées, Volume 1, part 4*. Roret, Paris, pp. 761–1084.
- Simon, E. 1896. Description d'un Arachnide cavernicole de l'Afrique australe. *Bulletin de la Société Entomologique de France*, 1869: 285–286.
- Simon, E. 1897a. *Histoire naturelle des Araignées, Volume 2, part 1*. Roret, Paris, 1–192.
- Simon, E. 1897b. On the Spiders of the Island of St. Vincent. Part III. *Proceedings of the Zoological Society of London*, 1897: 860–890.
- Simon, E. 1898a. *Histoire naturelle des Araignées, Volume 2, part 2*. Roret, Paris, 1–269.
- Simon, E. 1898b. Etude sur les Arachnides de la région des Maures (Var.) *Feuille des Jeunes Naturalistes*, (3) 29: 2–4.
- Simon, E. 1900. Descriptions d'arachnides nouveaux de la famille des Attidae. *Annales de la Société Entomologique de Belgique*, 44: 381–407.
- Simon, E. 1903. *Histoire naturelle des Araignées, Volume 2, part 4*. Roret, Paris, 669–1080.
- Simon, E. 1929. *Les Arachnides de France. Tome 6*. Paris, pp. 533–772.

- Simon, R. 1971. Neue Arthropodenfunde aus dem Stephan der Halleschen Mulde. *Bericht der Deutschen Gesellschaft für Geologische Wissenschaft, Reihe A: Geologie/Paläontologie*, 16: 53–62.
- Simonetta, A. M. & Delle Cave, L. 1978. Una possibile interpretazione filogenetica degli artropodi paleozoici. *Bollettino di zoologia*, 45: 87–90.
- Simpson, S. 1951. A new Eurypterid from the Upper Old Red Sandstone of Portishead. *Annals and Magazine of Natural History, series 12*, 4: 849–861.
- Siveter, D. J. & Selden, P. A. 1987. A new, giant xiphosurid from the lower Namurian of Weardale, County Durham. *Proceedings of the Yorkshire Geological Society*, 46: 153–168.
- Siveter, D. J., Sutton, M. D., Briggs, D. E. G. & Siveter, D. J. 2004. A Silurian sea spider. *Nature*, 431: 978–980.
- Sivhed, U. & Wallwork, J. A. 1978. An early Jurassic oribatid mite from southern Sweden. *Geologiska Föreningens I Stockholm Förhandlingar*, 100: 65–70.
- Smith, F. P. 1902. The spiders of Epping Forest. *Essex Naturalist*, 12: 181–201.
- Sørensen, W. E. 1884. Opiliones Laniatores (Gonyleptides W. S. Olim) Musei Hauniensis. *Naturhistorisk Tidsskrift, Kjøbenhavn, series 3*, 14: 555–646.
- Sørensen, W. 1886. Opiliones. pp. 53–86. In Koch, L. & Keyserling, E. (eds) *Die Arachniden Australiens nach der Natur Beschrieben und Abgebildet*. Bauer und Raspe, Nürnberg, xx pp.
- Sørensen, W. 1932. Descriptiones Laniatorum (Arachnidorum Opilionum Subordinis). (Opus posthum recognovit et editit Kai L. Henriksen). – *Kongelige Danske Videnskabernes Selskabs Skrifter - Naturvidenskab og Mathematisk Afdeling, København, ser. 9*, 3(4): 197–422.
- Soriano, C., Archer, M., Azar, D., Creaser, P., Delclòs, X., Godhelp, H., Hand, S., Jones, A., Nel, A., Nèraudeau, D., Ortega-Blanco, J., Pèrez-de la Fuente, R., Perrichot, V., Saupe, E., Solòrzano-Kraemer, M., Taffreau, P. 2010. Synchrotron X-ray imaging on inclusions in amber. *Comptes Rendus Palevol*, 9, 361–368.
- Southcott, R. V. 1957. Description of a new Australian raphignathoid mite, with remarks on the classification of the Trombidiformes (Acarina). *Proceedings of the Linnean Society of New South Wales*, 81(3): 306–312.
- Southcott, R. V. & Lange, R. T. 1971. Acarine and other microfossils from the Maslin eocene, South Australia. *Records of the South Australian Museum*, 16: 1–21.
- Stahnke, H. L. 1940. The scorpions of Arizona. *Iowa State College Journal of Science*, 15: 101–103. [Thesis abstract.]
- Sterzel, J.T. 1918. Die organischen Reste des Kulms und Rotliegenden der Gegend von Chemnitz. *Abhandlungen der Königlich Sächsischen Gesellschaft der Wissenschaften, Mathematisch-physikalische Klasse*, 35: 1–315.
- Stock, J. H. 1954. Papers from Dr. Th. Mortensen's Pacific expedition 1914–1916. LXXVII. Pycnogonida from Indo-West-Pacific, Australian, and New-Zealand waters. *Videnskabelige Meddelelser fra Dansk naturhistorisk Foreningen*, 116(1): 1–168.

- Stott, C. A., Tetlie, O. E., Braddy, S. J., Nowlan, G. S., Glasser, P. M. & Devereux, M. G. 2005. A new eurypterid (Chelicerata) from the Upper Ordovician of Manitoulin Island, Ontario, Canada. *Journal of Paleontology*, 79: 1166–1174.
- Stainier, X. 1917. On a new eurypterid from the Belgian Coal Measures. *Quarterly Journal of the Geological Society*, 71: 639–647.
- Størmer, L. 1934a. Downtonian Merostomata from Spitsbergen with remarks on the suborder Synziphosura. *Skrifter utgitt av Det Norske Videnskaps-Akademi I Oslo, I. Matem.-Naturvid. Klasse*, 1933(3): 1–26.
- Størmer, L. 1934b. Merostomata from the Downtonian Sandstones of Ringerike, Norway. *Skrifter utgitt av Det Norske Videnskaps-Akademi I Oslo, I. Matem.-Naturvid. Klasse*, 1933(10): 1–125.
- Størmer, L. 1934c. Über den neuen von W. Gross beschriebenen Eurypteriden aus dem Unterdevon von Overath im Rheinland. *Jahrbuch der Preussischen Geologischen Landesanstalt*, 55: 284–291.
- Størmer, L. 1934d. A new Eurypterid from the Saaremaa-(Oesel-)Beds in Estonia. *Publications of the Geological Institution of the University of Tartu*, 37: 1–8.
- Størmer, L. 1936a. Eurypteriden aus dem Rheinischen Unterdevon. *Abhandlungen der Preussischen Geologischen Landesanstalt, Neue Folge*, 175: 1–74.
- Størmer, L. 1936b. *Mixopterus dolichoshelus* (Laurie MS), a Downtonian eurypterid from Scotland. *Summary of Progress of the Geological Survey for 1934*: 41–46.
- Størmer, L. 1951. A new eurypterid from the Ordovician of Montgomeryshire, Wales. *Geological Magazine*, 88: 409–422.
- Størmer, L. 1952. Phylogeny and taxonomy of fossil horseshoe crabs. *Journal of Paleontology*, 26: 630–639.
- Størmer, L. 1963. *Gigantoscrapio willsi*, a new scorpion from the Lower Carboniferous of Scotland and its associated preying microorganisms. *Skrifter Utgitt av det Norske Videnskaps-Akademi I Oslo. Matematisk-Naturvidenskabelig Klasse*, 8: 1–171.
- Størmer, L. 1969. Eurypterids from the Lower Devonian of Willwerath, Eifel. *Senckenbergiana lethaea*, 50: 21–35.
- Størmer, L. 1970. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 1: Arachnida. *Senckenbergiana lethaea*, 51: 335–369.
- Størmer, L. 1972. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 2: Xiphosura. *Senckenbergiana lethaea*, 53: 1–29.
- Størmer, L. 1973. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 3: Eurypterida, Hughmilleridae. *Senckenbergiana lethaea*, 54: 119–205.
- Størmer, L. 1974. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 4: Eurypterida, Drepanopteridae, and other groups. *Senckenbergiana lethaea*, 54: 359–451.
- Størmer, L. 1976. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 5: Myriapoda and additional forms, with general remarks on the fauna and problems regarding invasion of land by arthropods. *Senckenbergiana lethaea*, 57: 87–183.

- Størmer, L. & Waterston, C. D. 1968. *Cyrtoctenus* gen. nov., a large late Palaeozoic arthropod with pectinate appendages. *Transactions of the Royal Society Edinburgh*, 68: 63–104.
- Strand, E. 1926. Miscellanea nomenclatorial zoologica et palaeontologica. I–II. *Archiv für Naturgeschichte A*, 92(8): 30–75.
- Strand, E. 1929. Zoological and palaeontological nomenclatorial notes. *Acta Universitatis Latviensis*, 20: 29 pp.
- Strand, E. 1932. Miscellanea nomenclatorialica zoologica et palaeontologica, III, IV. *Folia zoologica et hydrobiologica*, 4: 133–147, 193–196.
- Strand, E. 1942. Miscellanea nomenclatorialica zoologica et palaeontologica. *Folia Zoologica et Hydrobiologica*, 11: 386–402.
- Strenzke, K. 1963. Entwicklung und Verwandtschaftsbeziehungen der Oribatidengattung *Gehypochthonius* (Arach., Acari). *Senckenbergiana Biologica*, 44: 231–255.
- Stumm, E. C. & Kjellesvig-Waering, E. N. 1962. A new eurypterid from the Upper Silurian of southern Michigan. *Contributions from the Museum of Paleontology, The University of Michigan*, 17: 195–204.
- Stur, D. 1877. Die Culm-Flora der Ostrauer und Waldenburger Schichten. *Abhandlung der königliche geologische Reichanstalt*, 4: 5.
- Subías, L. S. 2004. Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo. *Graellsia* 60 (número extraordinario), 3–305. Available from: <http://www.ucm.es/info/zoo/Artropodos/Catalogo.pdf>.
- Subías, L. S. & Arillo, A. 2002. Oribatid mite fossils from the Upper Devonian of South Mountain, New York and the Lower Carboniferous of County Antrim, Northern Ireland (Acariformes, Oribatida). *Estudios del Museo de Ciencias Naturales de Alava*, 17: 93–106.
- Sundevall, J.C. 1833. *Conspectus Arachnidium*. C. F. Berling, Londini Gothorum, 39 pp.
- Swartz, C. K. 1923. Order Eurypterida. 716–778. In Swartz, C. K., Prouty, W. F., Ulrich, E. O. & Bassler, R. S. (eds). *Silurian Volume*. Maryland Geological Survey, 795 pp.
- Taczanowski, L. 1879. Les aranéides du Pérou central (suite). *Horae Societatis entomologicae Rossicae*, 15: 102–136.
- Tasch, P. 1961. Paleolimnology: part 2 – Harvey and Sedgwick counties, Kansas: stratigraphy and biota. *Journal of Paleontology*, 35: 836–865.
- Tasch, P. 1963. Paleolimnology: part 3 – Marion and Dickinson counties, Kansas, with additional sections in Harvey and Sedgwick counties: stratigraphy and biota. *Journal of Paleontology*, 37: 1233–1251.
- Tesakov, A. S. & Alekseev, A.S. 1992. Myriapod-like arthropods from the Lower Devonian of central Kazakhstan. *Paleontological Journal*, 26: 18–23.
- Tesakov, A. S. & Alekseev, A.S. 1998. *Maldybulakia* – new name for *Lophodesmus* Tesakov and Alekseev, 1992 (Arthropoda). *Paleontological Journal*, 32: 29.

- Tetlie, O. E. 2002. A new *Baltoeurypterus* (Eurypterida: Chelicerata) from the Wenlock of Norway. *Norwegian Journal of Geology*, 82: 37–44.
- Tetlie, O. E. 2006a. Two new Silurian species of *Eurypterus* (Chelicerata: Eurypterida) from Norway and Canada and the phylogeny of the genus. *Journal of Systematic Palaeontology* 4: 397–412.
- Tetlie, O. E. 2006b. Eurypterida (Chelicerata) from the Welsh Borderlands, England. *Geological Magazine*, 143: 723–735.
- Tetlie, O. E. & Braddy, S.J. 2004. The first Silurian chasmataspid, *Loganamaraspis dunlopi* gen. et sp. nov. (Chelicerata: Chasmataspidida) from Lesmahagow, Scotland, and its implications for eurypterid phylogeny. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, 94: 227–234.
- Tetlie, O. E. & Briggs, D. E. G. 2009. The origin of pterygotid eurypterids (Chelicerata: Eurypterida). *Palaeontology*, 52: 1141–1148.
- Tetlie, O. E. & Dunlop, J. A. 2008. *Geralinura carbonaria* (Arachnida; Uropygi) from Mazon Creek, Illinois, USA, and the origin of subchelate pedipalps in whip scorpions. *Journal of Paleontology*, 82: 299–312.
- Tetlie, O. E. & Van Roy, P. 2006. A reappraisal of *Eurypterus dumonti* Stainier, 1917 and its position within the Adelophthalmidae Tollerton, 1989. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre* 76: 79–90.
- Tetlie, O. E. & Poschmann, M. 2008. Phylogeny and palaeoecology of the Adelophthalmoidea (Arthropoda; Chelicerata; Eurypterida). *Journal of Systematic Palaeontology*, 6: 237–249.
- Tetlie, O. E., Selden, P. A. & Ren D. 2007. A new Silurian eurypterid (Arthropoda: Chelicerata) from China. *Palaeontology*, 50: 619–625.
- Tetlie O. E., Braddy, S. J., Butler, P.D. & Briggs, D.E.G. 2004. A new eurypterid (Chelicerata: Eurypterida) from the Upper Devonian Gogo Formation of Western Australia, with a review of the Rhenopteridae. *Palaeontology* 47: 801–809.
- Thevenin, A. 1901. Sur le découverte d'arachnides dans le Terrain Houiller de Commeny. *Bulletin de la Société Géologique de France*, 4^e Série, 1: 605–611.
- Thevenin, A. 1902. Sur une araignée du terrain houiller der Valenciennes. *Procès-Verbaux de la Société d'Histoire Naturelle de Autun*, 15: 195–203.
- Thompson, W. D'Arcy 1909. Pycnogonida. In Harmer, S. F. & Shipley, B. E. (eds). *The Cambridge Natural History*, pp. 501–542.
- Thor, S. 1900. Norske hydrachnider IV. – *Archiv för Matematik og Naturvidenskab* 23: 1-56.
- Thor, S. 1905. Eine interessante neue Milbengattung aus der schweizerischen Sammlung des Herrn Dr. W. Volz. *Zoologischer Anzeiger*, 28: 505–509.
- Thor, S. 1911a. *Lebertia*-Studien XXIV–XXV. *Zoologischer Anzeiger*, 37: 385–394.
- Thor, S. 1911b. Eine neue Acarinenfamilie (Teneriffiidae) und zwei neue Gattungen, die eine von Teneriffa, die andre aus Paraguay. *Zoologischer Anzeiger*, 38: 171–179.

- Thor, S. 1927. Acarinologische Notizen. *Zoologischer Anzeiger*, 72: 155–159.
- Thor, S. 1933. Über die prostigmatische Familie: Eupodidae C.L.Koch 1842 und über die Teilung dieser Familie, mit Definitionen der neuen Familien. *Zoologischer Anzeiger*, 101: 271–277.
- Thor, S. 1934. Neue Beiträge zur Kenntnis der invertebraten Fauna von Svalbard. (Nach Sammlungen von Garteninspektor L. Lange, Dozent B. Lynge und dem Verfasser.). *Zoologischer Anzeiger*, 107: 114–139.
- Thor, S. 1935. Übersicht und Einteilung der Familie Trombidiidae W.E. Leach 1814 in Unterfamilien. *Zoologischer Anzeiger*, 109: 107–112.
- Thor, S. 1937. Übersicht der norwegischen *Cryptostigmata* mit einzelnen Nebenbemerkungen. *Saertrykk av Nytt Magasin for Naturvidenskapene*, 77: 275–307.
- Thorell, T. 1856. Recensio critica Araneorum Suecicarum quas descripserunt Clerckius, Linnaeus, de Geerus. *Nova Acta Societas Scientiae Uppsalensis*, 2: 61–176.
- Thorell, T. 1869. On European spiders. Part I. Review of the European genera of spiders, preceded by some observations on zoological nomenclature. *Nova Acta Societas Scientiae Uppsalensis*, (3)7: 1–108.
- Thorell, T. 1870a. On European spiders. Part 2. *Nova Acta Societas Scientiae Uppsalensis*, (3)7: 109–242.
- Thorell, T. 1870b. *Remarks on synonyms of European spiders. Part I.* Uppsala, pp. 1–96.
- Thorell, T. 1873. *Remarks on synonyms of European spiders. Part IV.* Uppsala, pp. 375–645.
- Thorell, T. 1875. Diagnoses Araneorum Europaeorum aliquot novarum. *Tijdschrift voor Entomologie*, 18: 81–108.
- Thorell, T. 1876a. Études Scorpiologiques. *Atti della Società Italiana di Scienze Naturali*, 19: 75–272.
- Thorell, T. 1876b. On the classification of scorpions. *Annals and Magazine of Natural History, series 4*, 17: 1–15.
- Thorell, T. 1876c. Sopra alcuni Opilioni (Phalangidea) d'Europa e dell'Asia occidentale, con un quadro dei generi europei di quest'Ordine. *Annali del Museo Civico di Storia Naturale (Genoa) series 1*, 8: 452–508.
- Thorell, T. 1881. Studi sui Ragni Malesi e Papuani. III. Ragni dell'Austro Malesia e del Capo York, conservati nel Museo civico di storia naturale di Genova. *Annali del Museo Civico di Storia Naturale di Genova*, 17: 1–727.
- Thorell, T. 1882. Descrizione di Alcuni Aracnidi Inferiori dell' Arcipelago Malese. *Annali del Museo Civico di Storia Naturale di Genova*, 18: 21–69.
- Thorell, T. 1887. Viaggio di L. Fea in Birmania e regioni vicine. II. Primo saggio sui ragni birmani. *Annali del Museo Civico di Storia Naturale di Genova*, 25: 5–417.
- Thorell, T. 1888. Pedipalpi e Scorpioni dell'Arcipelago Malese conservati nel Museo Civico di Storia Naturale di Genova. *Annali del Museo Civico di Storia Naturale di Genova*, 26: 327–428.
- Thorell, T. 1889. Viaggio di Leonardo Fea in Birmania e regioni vicine. XXI. Aracnidi Artrogastri Birmani raccolti da L. Fea nel 1885–1887. *Annali del Museo Civico di Storia Naturale di Genova*, 27: 521–729.
- Thorell, T. 1890. Studi sui ragni Malesi e Papuani. Part IV, 1. *Annali del Museo Civico di Storia Naturale di Genova*, 28: 1–419.

- Thorell, T. 1891. Spindlar från Nikobarerna och andra delar af södra Asien. *Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar*, 24: 149 pp.
- Thorell, T. & Lindström, G. 1884. Discovery of a Silurian fossil scorpion. *The Glasgow Herald*, Dec. 19, 1884.
- Thorell, T. & Lindström, G. 1885. On a Silurian scorpion from Gotland. *Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar*, 21(9): 1–33.
- Tollerton, V. P., Jr. 1989. Morphology, taxonomy, and classification of the order Eurypterida Burmeister, 1843. *Journal of Paleontology*, 63: 642–657.
- Trägårdh, I. 1902. Beiträge zur Kenntnis der schwedischen Acaridenfauna. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar*, 28: 1–26.
- Trägårdh, I. 1915. Bidrag till kännedomen om spinnkvalstren (*Tetranychus* Duf.). *Centralanstalten för försöksväsendet på jordbruksområdet. Entomologiska avdelningen*, 20: 1–60.
- Trägårdh, I. 1946. Outlines of a new classification of the Mesostigmata (Acarina) based on comparative morphological data. *Lunds Universitets Arsskrift, N.F.* 42: ??
- Trägårdh, I. 1950. Description of a new species of *Heterocheylus* Lombardini from Africa, with notes on the classification of the Pseudocheyletidae. *Entomologisk tidskrift*, 71: 104–110.
- Travé, J. 1959. Sur le genre *Niphocephus* Balogh 1943. Les Niphocephidae, famille nouvelle (Acariens, Oribates). *Acarologia*, 1: 475–498.
- Travé, J. 1967. *Phyllochthonius aoutii* nov. gen., nov. spec., un Enarthronota (Acarien, Oribate) nouveau de Côte d'Ivoire, avec la création d'une superfamille nouvelle, Phyllochthonoidea. *Zoologische Mededelingen*, 42: 83–105.
- Treat, A. E. 1955. An ectoparasite (Acarina: Mesostigmata) from moths of the genus *Zale*. *Journal of Parasitology* 41: 555–561.
- Türk, E. 1963. A new tyroglyphid deutonymph in amber from Chiapas, Mexico. *University of California Publications in Entomology* 31: 49–51.
- Ubick, D. & Dunlop, J. A. 2005. On the placement of the Baltic amber harvestman *Gonyleptes nemastomoides* Koch & Berendt, 1854, with notes on the phylogeny of Cladonychiidae (Opiliones, Laniatores, Travunioidea). *Mitteilungen aus dem Musuem für Naturkunde Berlin, Geowissenschaftliche Reihe* 8: 75–82.
- Vachon, M. & Heyler, D. 1985. Description d'une nouvelle espèce de Scorpion: *Buthiscorpius pescei* (Stéphanien de Montceau-les-Mines, France). Remarques sur la classification des Scorpions (Arachnida) du Carbonifère. *Bulletin de la Société d'Histoire Naturelle d'Autun* 113: 29–47.
- Vandenbergh, A. 1960. *Pringlia demaisteri* nov. sp., un xiphosure (Chélicérate) du Stéphanien de la Loire. – *Bulletin de la Société géologique de France* 7: 687–689.

- Vercammen-Grandjean, P. H. 1973. Study of the "Erythraeidae, R.O.M. No. 8" of Ewing, 1937. 329–335. In Daniel, M. and Rosický, B. (eds). *Proceedings of the 3rd International Congress of Acarology*. Academia, Prague, 837 pp.
- Via Boada, L. & Villalta, J. F. de 1966. *Hetrolimulus gadeai*, nov. gen., nov. sp., représentant d'une nouvelle famille de Limulacés dans le Trias d'Espagne. *Comptes Rendues Sommaire Séances Société Géologique France*, 1966: 57–59.
- Viets, K. O. 1978. New water mites (Hydrachnellae: Acari) from Australia. *Australian Journal of Marine and Freshwater Research*, 29: 77–92.
- Villalta, J. F. 1957. Dos zoocecidias fósiles del Mioceno de Cerdaña (prov. de Lérida). *Cursillos y conferencias del Instituto Lucas Mallada*, 4: 63–64.
- Vitzthum, H. Graf 1931. Acari=Milben. In Kukenthal, W. (ed.) *Handbuch der Zoologie, Vol. III 2. 3*. Walter de Gruyter & Co., Berlin, pp. 1–160.
- Vitzthum, H. G. 1942. Acarina. In *Bronn's Klassen und Ordnungen des Tierreiches, IV. Abt., 5. Buch, 5. Lieferung* (1942), Leipzig, Akademische Verlagsgesellschaft Becker u. Erler: pp. 641–800.
- Wagner, W. A. 1887. Copulations organe des Männchens als Criterium für die Systematik der Spinnen. *Horae Societatis Entomologicae Rossicae*, 22: 3-132.
- Walcott, C. D. 1882. Description of a new genus of the order Eurypterida from the Utica Slate. *American Journal of Science, 3rd Series*, 23: 213–216.
- Walckenaer, C. A. 1802. *Faune parisienne. Insectes. Ou Histoire abrégée des Insectes des environs de Paris*. Paris, 2: 187–250.
- Walckenaer, C. A. 1805. *Tableau des Aranéides ou Caractères essentiels des tribus, genres, familles et races que renferme le genre Aranea de Linné, avec la désignation des espèces comprises dans chacune de ces divisions*. Paris, 88 pp.
- Walckenaer, C. A. 1826. Aranéides. In *Faune française...*, Paris: 96 pp.
- Walckenaer, C. A. 1837. *Histoire naturelle des insectes. Aptères. Vol. 1*. Librairie Encyclopédique de Roret, Paris, 682 pp.
- Walker, N. A. 1965. Euphthiracaroida of California Sequoia litter : with a reclassification of the families and genera of the world (Acarina: Oribatei). *Fort Hays Studies, New Series, Science Series*, 3: 154 pp.
- Walossek, D., Li, C.S. & Brauckmann, C. 1990. A scorpion from the Upper Devonian of Hubei Province, China (Arachnida, Scorpiones). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1990(3): 169–180.
- Waloszek, D. & Dunlop, J. A. 2002. A larval sea spider (Arthropoda: Pycnogonida) from the Upper Cambrian 'Orsten' of Sweden and the phylogenetic position of pycnogonids. *Palaeontology*, 45: 421–446.
- Walter, D. E. 1997. Heatherellidae – a new family of Mesostigmata (Acari: Parasitiformes) based on two new species from rainforest litter in Australia. *International Journal of Acarology*, 23: 167–175.

- Walter, D. E. 2000. A jumping mesostigmatan mite, *Saltiseius hunteri* n. g., n. sp. (Acari: Mesostigmata: Trigynaspida: Saltiseiidae, n. fam.) from Australia. *International Journal of Acarology*, 26: 25–31.
- Walter, D. E. & Gerson, U. 1998. Dasythyreidae, new family, and *Xanthodasythyreus* n. g. (Acari: Prostigmata: Raphignathoidea) from Australia. *International Journal of Acarology*, 24: 189–197.
- Walter, D. E. & Krantz, G. W. 1999. New early derivative mesostigmatans from Australia: *Nothogynus* n. g., Nothogynidae n. fam. (Mesostigmata: Microgyniina). *International Journal of Acarology*, 25: 67–76.
- Waterston, C. D. 1962. *Pagea sturrocki* gen. et sp. nov., a new eurypterid from the Old Red Sandstone of Scotland. *Palaeontology*, 5: 137–148.
- Waterston, C. D. 1964. Observations on pterygotid eurypterids. *Transactions of the Royal Society of Edinburgh*, 66: 9–33.
- Waterston, C. D. 1968. Further observations on the Scottish Carboniferous eurypterids. *Transactions of the Royal Society of Edinburgh*, 68: 1–20.
- Waterston, C. D. 1979. Problems of functional morphology and classification in stylonurid eurypterids (Chelicerata, Merostomata), with observations on the Scottish Stylonuroidea. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 70: 251–322.
- Waterston, C. D. 1985. Chelicerata from the Dinantian of Fouldon, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 76: 25–33.
- Waterston, C. D., Oelofsen, B. W. and Ooshuizen, R. D. F. 1985. *Cyrtoctenus wittebergensis* sp. nov. (Chelicerata: Eurypterida), a large sweep-feeder from the Carboniferous of South Africa. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 76: 339–358.
- Watson, D. M. S. 1909. *Limulus woodwardi*, sp. nov., from the Lower Oolite of England. *Geological Magazine, New Series*, (5) 6: 14–15.
- Waterlot, G. 1934. *Étude de la Faune continentale du Terrain houiller Sarro-Lorrain – Études des gîtes minéraux de la France. Bassin houiller de la Sarre et de la Lorraine II. Faune fossile*. Lille, 317 pp.
- Weidner, H. 1964. Eine Zecke, *Ixodes succineus* sp. n. im Batischen Bernstein. *Veöffentlichunge aus dem Überseemuseum Bremen*, 3: 143–151.
- Weitschat, W. & Wichard, W. 2002. *Atlas of plants and animals in Baltic amber*. Dr. F. Pfeil, Munich, 256 pp.
- Westring, N. 1851. Förteckning öfver de till närvarande tid Kände, i Sverige förekommande Spindlarter, utgörande ett antal af 253, deraf 132 äro nya för svenska Faunan. *Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhälles handlingar*, 2: 25–62.
- Westwood, J. O. 1835. Insectorum Arachnoidumque novorum Decades duo. *The Zoological Journal, London*, 5: 440–453.
- Westwood, J. O. 1874. *Thesaurus entomologicus oxoniensis*. Clarendon Press, Oxford, xx pp.
- Weyenbergh, H., Jr 1874. Notes sur quelques insectes du calcaire jurassique de la Bavière. *Archives Musée Teyler, Haarlem*, 3: 234–236.

- Weygoldt, P. 1996. Evolutionary morphology of whip spiders: towards a phylogenetic system (Chelicerata: Arachnida: Amblypygi). *Journal of Zoological Systematics and Evolutionary Research*, 34: 185–202.
- Weygoldt, P. & Paulus, H.F. 1979. Untersuchungen zur Morphologie, Taxonomie und Phylogenie der Chelicerata. *Zeitschrift für zoologische Systematik und Evolutionsforschung*, 17: 85–115, 177–200.
- White, D. 1908. Report on the fossil flora of the Coal Measures of Brazil. 377–607. In White, J. C. (ed.). *Final report on the coal measures and associated rocks of South Brazil*. Comissão de Estudos das Minas de Carvão de Pedra Do Brazil, Rio de Janeiro.
- Whiteaves, J. F. 1884. On some new, imperfectly characterized or previously unrecorded species of fossils from the Guelph Formations of Ontario. *Palaeozoic Fossils of Canada*, 3(1):1–43
- Whitfield, R. P. 1882. Descriptions of new species of fossils from Ohio, with remarks on some of the geological formations in which they occur. *Annals of the New York Academy of Science*, 2: 193–244.
- Whitfield, R. P. 1885a. An American Silurian scorpion. *Science*, 6: 87–88.
- Whitfield, R. P. 1885b. On a fossil scorpion from the Silurian rocks of America. *Bulletin of the American Museum of Natural History*, 1(9): 181–190.
- Wiles, P. R. 1996. A new family, genus and species of watermite (Acari: Hydrachnidia, Lebertioidea) from Brunei. *Quekett Journal of Microscopy*, 37: 692–695.
- Williams, H. 1915. An eurypterid horizon in the Niagara Formation of Ontario. *Geological Survey of Canada, Museum Bulletin*, 20: 1–9.
- Willmann, C. 1931. Oribatei (Acari), gesammelt von der Deutschen Limnologischen Sunda-Expedition. *Archiv für Hydrobiologie*, Supplement-Band IX: 240–305.
- Wills, L. J. 1910. On the fossiliferous Lower Keuper rocks of Worcestershire, with descriptions of some of the animals discovered therein. *Proceedings of the Geologists' Association*, 21: 249–331.
- Wills, L. J. 1947. *A monograph of the British Triassic scorpions*. The Palaeontographical Society, London, 100 & 101: 137 pp.
- Wills, L. J. 1959. The external anatomy of some Carboniferous “scorpions” Part 1. *Palaeontology*, 1: 261–282.
- Wills, L. J. 1960. The external anatomy of some Carboniferous “scorpions”. Part 2. *Palaeontology*, 3: 276–332.
- Wilson, E. B. 1878. Descriptions of two new genera of Pycnogonida. *American Journal of Science*, 15: 200–203.
- With, C. J. 1902. A new acaride *Opilioacarus segmentatus*. *Comptes Rendus du Congrès des Naturalistes et Médecins du Nord*, 20: 4–5.
- With, C. J. 1906. The Danish expedition to Siam 1899–1900. III. Chelonethi. An account of the Indian false-scorpions together with studies on the anatomy and classification of the order. *Oversight over det Kongelige Danske Videnskabernes Selskabs Forhandling*, 7(3): 1–214.
- Witaliński, W. 2000. *Aclerogamasus stenocornis* sp. n., a fossil mite from the Baltic amber (Acari: Gamasida: Parasitidae). *Genus*, 11: 619–626.

- Wolff, R.J. 1990. A new species of *Thiodina* (Araneae: Salticidae) from Dominican amber. *Acta Zoologica Fennica*, 190: 405–408.
- Womersley, H. 1956. On some new Acarina-Mesostigmata from Australia, New Zealand and New Guinea. *Zoological Journal of the Linnean Society of London*, 42: 505–599.
- Womersley, H. 1957. A fossil mite (*Acronothrus ramus* n.sp.) from Cainozoic resin at Allendale, Victoria. *Proceedings of the Royal Society of Victoria* 69: 21–23.
- Wood, T. G. 1969. The Homocaligidae, a new family of mites (Acari: Raphignathoidea), including a description of a new species from Malaya and the British Solomon Islands. *Acarologia*, 11: 711–729.
- Woodward, H. 1865. On a new genus of Eurypterida from the Lower Ludlow rock of Leintwardine, Shropshire. *Quarterly Journal of the Geological Society of London*, 21: 490–492.
- Woodward, H. 1868a. On a new limuloid crustacean (*Neolimulus falcatus*) from the Upper Silurian of Lesmahagow, Lanarkshire. *Geological Magazine*, 5: 1–3.
- Woodward, H. 1870. On *Necrogammarus salweyi* (H. Woodward), an amphipodus crustacean from the Lower Ludlow of Leintwardine. *Transactions of the Woolhope Naturalists Field Club*, 1870: 271–272.
- Woodward, H. 1871a. On the remains of a giant isopod *Praearcturus gigas*, (H. Woodward) from the Old Red Sandstone of Rowlestone Quarry, Herefordshire. *Transactions of the Woolhope Field Naturalist's Club*, 1871: 266–270.
- Woodward, H. 1871b. On the discovery of a new and very perfect Arachnide from the ironstone of the Dudley Coal-field. *Geological Magazine*, 8: 385–388.
- Woodward, H. 1872a. Notes on some British Palaeozoic Crustacea belonging to the order Merostomata. *Geological Magazine*, 9: 433–441.
- Woodward, H. 1872b. On a new Arachnide from the Coal-measures of Lancashire. *Geological Magazine*, 9: 385–387.
- Woodward, H. 1876. On the discovery of a fossil scorpion in the British Coal Measures. *Quarterly Journal of the Geological Society of London* 32: 57–59.
- Woodward, H. 1878b. Discovery of the remains of a fossil crab (Decapoda-Bracyura) in the Coal Measures of the Environs of Mons, Belgium. *Geological Magazine, new series, Decade 2*, 5: 433–436.
- Woodward, H. 1879. Contributions to the knowledge of fossil Crustacea. *Quarterly Journal of the Geological Society London*, 35: 549–555.
- Woodward, H. 1887. On a new species of *Eurypterus* from the Lower Carboniferous shales of Glencartholm, Eskdale, Scotland. *Geological Magazine, Decade 3*, 4: 481–484.
- Woodward, H. 1888. Note on *Eurypterus* from the Carboniferous. *Geological Magazine, Decade 3*, 5: 419–421.
- Woodward, H. 1907a. Two new species of *Eurypterus* from the Coal-Measures of Ilkeston, Derbyshire. *Geological Magazine*, 4: 277–282.

- Woodward, H. 1907*b*. Further notes on the Arthropoda of the British Coal Measures. *Geological Magazine*, 4: 539–549.
- Woodward, H. 1918. Fossil arthropods from the Carboniferous rocks of Cape Breton, Nova Scotia; and from the Upper Coal Measures, Sunderland, England. *Geological Magazine*, 5: 462–471.
- Woolley, T. A. 1969. Two new species of *Hydrozetes*, extant and fossil (Acari: Cryptostigmata, Hydrozetidae). *New York Entomological Society*, 77: 250–256.
- Woolley, T. A. 1971. Fossil oribatid mites in amber from Chiapas, Mexico (Acarina: Oribatei = Cryptostigmata). *University of California Publications in Entomology*, 63: 91–99.
- Woolley, T. A. & Higgins, H. G. 1968. Megeremaeidae: A New Family of Oribatid Mites (Acari: Cryptostigmata). *Great Basin Naturalist*, 28(4): 172–175.
- Wright, D. F. & Selden, P. A. 2011. A trigonotarbid arachnid from the Pennsylvanian of Kansas. *Journal of Paleontology*, 85: 871–876.
- Wunderlich, J. 1981. Fossile Zwergsechsaugenspinnen (Oonopidae) der Gattung *Orchestina* Simon, 1882 in Bernstein mit Anmerkungen zur Sexual-biologie (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 51: 83–113.
- Wunderlich, J. 1982. Die häufigsten Spinnen (Araneae) des Dominikanischen Bernsteins. *Neue Entomologische Nachrichten*, 1: 26–45.
- Wunderlich, J. 1985. Ein bisher unbekannte fossile Krabbenspinne aus dem Randecker Maar in Südwest-Deutschland (Arachnida: Araneae: Thomisidae). *Neue Entomologische Nachrichten*, 14: 4–13.
- Wunderlich, J. 1986. *Spinnenfauna Gestern und Heute. Fossile Spinnen in Bernstein und ihre heute lebenden Verwandten*. Erich Bauer Verlag bei Quelle und Meyer, Wiesbaden, 283 pp.
- Wunderlich, J. 1987. *Tama minor n. sp.*, eine fossile Spinnenart der Familie Hersiliidae in Dominikanischem Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 97: 93–96.
- Wunderlich, J. 1988. Die fossilen Spinnen im dominikanischen Bernstein. *Beiträge zur Araneologie*, 2: 1–378.
- Wunderlich, J. 1991. Beschreibung der ersten fossilen Spinne der Familie Leptonetidae: *Eoleptona kutscheri* n. gen., n. sp. in Sächsischem Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 101: 21–26.
- Wunderlich, J. 1993a. Die ersten fossilen Speispinnen (Fam. Scytodidae) im Baltischen Bernstein (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 75: 243–247.
- Wunderlich, J. 1993b. Die ersten fossilen Becherspinnen (Fam. Cyatholipidae) in Baltischem und Bitterfelder Bernstein (Arachnida: Araneae). *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, 75: 231–241.
- Wunderlich, J. 1998. Beschreibung der ersten fossilen Spinnen der Unterfamilien Mysmeninae (Anapidae) und Erigoninae (Linyphiidae) im Dominikanischen Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift*, 108: 363–367.

- Wunderlich, J. 2000. Zwei neue Arten der Familie Falltürspinnen (Araneae: Ctenizidae) aus dem Baltischen Bernstein. *Entomologische Zeitschrift*, 110: 345–348.
- Wunderlich, J. 2004a. Introduction, general findings and conclusions. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 5–329.
- Wunderlich, J. 2004b. The fossil mygalomorph spiders (Araneae) in Baltic and Dominican amber and about extant members of the family Micromygalidae. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 595–631.
- Wunderlich, J. 2004c. Fossil spiders (Araneae) of the superfamily Dysderoidea in Baltic and Dominican amber, with revised family diagnoses. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 633–746.
- Wunderlich, J. 2004d. Fossil and extant spiders (Araneae) of the superfamily Eresoidea s.l., with special reference to the Archaeidae and remarks on some higher taxa of the superfamily Araneoidea. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 747–808.
- Wunderlich, J. 2004e. On selected higher and lower taxa of fossil and extant spiders of the superfamily Oecobioidea, with a provisional Cladogram (Araneae: Hersiliidae and Oecobiidae). In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 809–848.
- Wunderlich, J. 2004f. Fossil spiders of the family Uloboridae (Araneae) in Baltic and Dominican amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 851–886.
- Wunderlich, J. 2004g. The fossil spiders of the family Deinopidae in Baltic and Dominican amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 887–897.
- Wunderlich, J. 2004h. The fossil spiders (Araneae) of the families Tetragnathidae and Zygellidae n. stat. in Baltic and Dominican amber, with notes on higher extant and fossil taxa. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 899–955.
- Wunderlich, J. 2004i. Fossil taxa of the family Araneidae (Araneae) inclusively Nephilinae in Baltic and Dominican amber, with the description of a new extinct subfamily and notes on selected extant taxa. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 956–997.
- Wunderlich, J. 2004j. The fossil Theridiosomatidae (Araneae) in Baltic and Dominican amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 998–1019.
- Wunderlich, J. 2004k. The fossil spiders of the family Anapidae s. l. (Aeaneae [sic] in Baltic, Dominican and Mexican amber and their extant relatives, with the description of a new subfamily Comarominae. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1020–1111.
- Wunderlich, J. 2004l. On the relationships of the families of the superfamily Araneoidea (Araneae) and their kin, with cladograms, remarks on the origin of the orb web and description of the new and extinct families Baltsuccinidae and Protheridiidae in Tertiary Baltic amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1112–1154.
- Wunderlich, J. 2004m. The fossil spiders (Araneae) of the family Cyatholipidae in Baltic amber. In Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1155–1188.

- Wunderlich, J. 2004n. The fossil spiders (Araneae) of the family Synotaxidae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1189–1239.
- Wunderlich, J. 2004o. Remarks on the fossil spiders (Araneae) of the family Nesticidae in amber, with the description of a new species in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1240–1244.
- Wunderlich, J. 2004p. Remarks on fossil spiders (Araneae) of the family Theridiidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1245–1248.
- Wunderlich, J. 2004q. Fossil pirate spiders (Araneae: Araneoidea: Mimetidae s. l.) in Baltic and Dominican amber, with notes on intrafamiliar higher taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1249–1278.
- Wunderlich, J. 2004r. Descriptions of the first fossil spiders (Araneae) of the family Pimoidae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1279–1297.
- Wunderlich, J. 2004s. The fossil spiders of the family Linyphiidae in Baltic and Dominican amber (Araneae: Linyphiidae). *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1298–1373.
- Wunderlich, J. 2004t. No proof of fossil spiders (Araneae) of the family Psecridae in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1375–1376.
- Wunderlich, J. 2004u. Fossil spiders of the family Amaurobiidae (Arachnida: Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1377–1379.
- Wunderlich, J. 2004v. Fossil spiders of the family Dictynidae s. l., including Cryphoecinae and Hahniinae in Baltic and Dominican amber and copal from Madagascar, and on selected extant Holarctic taxa, with new descriptions and diagnoses. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1380–1482.
- Wunderlich, J. 2004w. Fossil spiders (Araneae) of the family Agelenidae s. str. in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1483–1488.
- Wunderlich, J. 2004x. The fossil Zoropsidae in Baltic amber with revised diagnoses of the family Zoropsidae and its fossil and extant higher taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1489–1522.
- Wunderlich, J. 2004y. Spiders (Araneae) of the extinct family Insecutoridae Petrunkevitch 1942 in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1523–1531.
- Wunderlich, J. 2004z. Fossil spiders of the family Pisauridae (Araneae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1532–1541.
- Wunderlich, J. 2004aa. Members of the family Trechaleidae (Araneae) in Baltic and Dominican amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1542–1553.
- Wunderlich, J. 2004ab. Fossil spiders (Araneae) of the family Oxyopidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1554–1556.
- Wunderlich, J. 2004ac. Proof of presence of the family Lycosidae (Araneae) in Baltic and Dominican amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1557–1558.

- Wunderlich, J. 2004ad. Fossil spiders (Araneae) of the extinct family Ephalmatoridae Petrunkevitch 1950 in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1559–1577.
- Wunderlich, J. 2004ae. Fossil spiders (Araneae) of the family Zodariidae in Baltic amber, with remarks on their subfamilies including the Cryptothelinae and the Homalonychinae. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1578–1611.
- Wunderlich, J. 2004af. Fossil spiders (Araneae) of the families Clubionidae and Miturgidae (questionable) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1612–1622.
- Wunderlich, J. 2004ag. The fossil spiders of the family Liocranidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1623–1635.
- Wunderlich, J. 2004ah. Fossil spiders of the family Corinnidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1636–1680.
- Wunderlich, J. 2004ai. Fossil spiders (Araneae) of the family Gnaphosidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1681–1685.
- Wunderlich, J. 2004aj. Fossil spiders (Araneae) of the family Anyphaenidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1686–1688.
- Wunderlich, J. 2004ak. Members of the family Philodromidae (Araneae) in Baltic amber? *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1689–1693.
- Wunderlich, J. 2004al. Fossil spiders (Araneae) of the family Sparassidae in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1694–1698.
- Wunderlich, J. 2004am. Fossil spiders of the family Trochanteriidae (Araneae) in Baltic, Dominican and Mexican amber, with a revision of the genus *Sosybius* Koch and Berendt 1854. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1699–1732.
- Wunderlich, J. 2004an. Fossil spiders of the family Selenopidae in Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1733–1736.
- Wunderlich, J. 2004ao. The new spider (Araneae) family Borboropactidae from the tropics and fossil in Baltic amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1737–1746.
- Wunderlich, J. 2004ap. Fossil crab spiders (Araneae: Thomisidae) in Baltic and Dominican amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1747–1760.
- Wunderlich, J. 2004aq. Fossil jumping spiders (Araneae: Salticidae) in Baltic and Dominican amber, with remarks on Salticidae subfamilies. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1761–1819.
- Wunderlich, J. 2004ar. Fossil spiders (Araneae) in Early Tertiary amber from the Ukraine. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1821–1829.
- Wunderlich, J. 2004as. Subrecent spiders (Araneae) in copal from Madagascar, with description of new species. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1830–1853.

- Wunderlich, J. 2004at. Two new fossil spider species in Copal from Colombia (Araneae: Oonopidae and Dictynidae). *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1854–1859.
- Wunderlich, J. 2004au. Description of two fossil taxa of spiders (Araneae: Oonopidae, Pholcidae) in Chinese amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1860–1863.
- Wunderlich, J. 2004av. Report on spider (Araneae) of the families Araneidae and Zygellidae in Lebanese amber. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 3: 1864–1865.
- Wunderlich, J. 2006. *Spatiator martensi* n. sp., a second species of the extinct spider species Spatiatoridae in Eocene Baltic amber. *Zootaxa*, 1325: 313–318.
- Wunderlich, J. 2008a. Descriptions of fossil spider (Araneae) taxa mainly in Baltic amber, as well as certain related extant taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 5: 44–139.
- Wunderlich, J. 2008b. On extant and fossil (Eocene) European comb-footed spiders (Araneae: Theridiidae), with notes on their subfamilies, and with descriptions of new taxa. *In* Wunderlich, J. (ed.). *Beiträge zur Araneologie*, 5: 140–469.
- Wunderlich, J. 2008c. On extant and fossil members of the RTA-clade in Eocene European ambers of the families Borboropactidae, Corinnidae, Selenopidae, Sparassidae, Trochanteriidae, Zoridae s. l., and of the superfamily Lycosoidea. *In* Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 5: 470–523.
- Wunderlich, J. 2008d. The dominance of ancient spider families of the Araneae: Haplogyne in the Cretaceous, and the late diversification of advanced ecribellate spiders of the Entelegynae after the Cretaceous–Tertiary boundary extinction events, with descriptions of new families. *In* Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 5: 524–675.
- Wunderlich, J. 2011a. On extant European spiders of the tribe Mangorini (Araneae: Araneidae) and two doubtful taxa in Baltic amber. *Beiträge zur Araneologie*, 6: 9–18.
- Wunderlich, J. 2011b. Taxonomy of extant and fossil (Eocene) European ground spiders of the family Gnaphosidae (Araneae), with a key to the genera, and descriptions of new taxa. *Beiträge zur Araneologie*, 6: 19–97.
- Wunderlich, J. 2011c. Spiders of the family Prodidomidae (Araneae) from Europe and Madagascar. *Beiträge zur Araneologie*, 6: 98–107.
- Wunderlich, J. 2011d. On extant and fossil (Eocene) Holarctic sac spiders (Araneae: Clubionidae), with descriptions of new taxa. *Beiträge zur Araneologie*, 6: 121–157.
- Wunderlich, J. 2011e. New extant taxa of the spider family Theridiosomatidae (Araneae) from Laos and on some fossil taxa. *Beiträge zur Araneologie*, 6: 427–444.
- Wunderlich, J. 2011f. Some subrecent spiders (Araneae) in copal from Madagascar. *Beiträge zur Araneologie*, 6: 445–460.
- Wunderlich, J. 2011g. Some fossil spiders in Dominican amber (Araneae: Hersiliidae, Theridiidae, Gnaphosidae). *Beiträge zur Araneologie*, 6: 461–471.

- Wunderlich, J. 2011*h*. Some fossil spiders (Araneae) in Eocene European ambers. *Beiträge zur Araneologie*, 6: 472–538.
- Wunderlich, J. 2011*i*. Some fossil spiders (Araneae) in Cretaceous ambers. *Beiträge zur Araneologie*, 6: 539–557.
- Wunderlich, J. 2012*a*. New subrecent species in copal from Madagascar, and on the relationships of the Copaldictyninae Wunderlich 2004 (Araneae: Linyphiidae, Theridiidae, Dictynidae, and Titanoecidae). In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 75–88.
- Wunderlich, J. 2012*b*. New fossil spiders (Araneae) in Eocene amber from the Ukraine. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 89–93.
- Wunderlich, J. 2012*c*. New fossil spiders (Araneae) of eight families in Eocene Baltic amber, and revisions of selected taxa. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 94–149.
- Wunderlich, J. 2012*d*. On the fossil spider (Araneae) fauna in Cretaceous ambers, with descriptions of new taxa from Myanmar (Burma) and Jordan, and on the relationships of the superfamily Leptonetoidea. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 157–232.
- Wunderlich, J. 2012*e*. Description of the first fossil Ricinulei in amber from Burma (Myanmar), the first report of this arachnid order from the Mesozoic and from Asia, with notes on the related extinct order Trigonotarbida. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 233–244.
- Wunderlich, J. 2012*f*. Corrections and addenda to vol. 6 of the Beitr. Araneol. (2011). In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 7: 245–246.
- Wunderlich, J. & Milki, R. 2004. Description of the extinct new subfamily Microsegestriinae (Araneae: Segestriidae) in Cretaceous Lebanese Amber. In Wunderlich, J. (ed.) *Beiträge zur Araneologie*, 3: 1867–1873.
- Żabka, M. 1988. Fossil Eocene Salticidae (Araneae) from the collection of the Museum of Earth in Warsaw. *Annales Zoologici*, 41: 415–420.
- Zacharda, M. 1979. Strandmanniidae – a new family of Eupodoidea (Acarina : Prostigmata). *Vestník Československé Společnosti Zoologické*, 43: 76–81.
- Zacharda, M. & Krivoluckij, D. A. 1985. Prostigmatic mites (Acarina: Prostigmata) from the Upper Cretaceous and Paleogene amber of the USSR. *Věstník Československé Společnosti Zoologické*, 49: 147–152.
- Zachvatkin, A. A. 1952. [The division of the Acarina into orders and their position in the system of the Chelicerata.] *Parazitologičeskii Sbornik Zoologičeskii Institut Akademii Nauk SSSR*, 14: 5–46. [in Russian]
- Zapfe, H. 1955. Filogenia y función en *Austrochilus manni* Gertsch y Zapfe (Araneae-Hypochilidae). *Trabajos del Laboratorio de Zoología de la Universidad de Chile*, 2: 1–53.
- Zhang, J., Sun, B. & Zhang, X. 1994. *Miocene insects and spiders from Shanwang, Shandong*. Science Press, Beijing, 298 pp. [in Chinese with English Summary].

- Zhang, Q.-y., Hu, S.-x., Zhou, C.-y., Lv, T. & Bai, J.-k. (2009): [New occurrence of Xiphosura in China.] *Progress in Nature Science*, 19: 1090–1093. [in Chinese]
- Zhang, Z.-Q. 1998: An unusual early-derivative larva of Parasitengona (Acari: Prostigmata) and proposal of a new superfamily. *Systematic & applied acarology*, 3: 159–170.
- Zhang, Z.-Q. & Fan, Q.-H. 2007. Allotanaupodidae, a new family of early derivative Parasitengona (Acari: Prostigmata). *Zootaxa*, 1517: 1–52.
- Zinken, C. 1862. *Limulus Decheni* aus dem Braunkohlensandstein bei Teuchern. *Zeitschrift für die Gesammten Naturwissenschaften*, 19: 329–331.
- Zittel, K. A. 1885. *Handbuch der Palaeontologie. I. Abtheilung, Palaeozoologie, 2 [Mollusca und Arthropoda]*. R. Oldenbourg, München, Leipzig, 893 pp.
- Zittel, K. A. & Eastman, C. R. 1913. *Textbook of Palaeontology (2nd Ed.) 1*. Macmillan, London, 839 pp.